

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
19 May 2005 (19.05.2005)

PCT

(10) International Publication Number
WO 2005/044854 A2

(51) International Patent Classification⁷: **C07K 16/00**

CA 94662-8097 (US). **HURST, Deborah** [US/US]; Chiron Corporation, P.O.Box 8097, Emeryville, CA 94662-8097 (US).

(21) International Application Number:
PCT/US2004/037152

(74) **Agent: ALEXANDER, LiSA, E.**; Chiron Corporation, Intellectual Property Department, P.O.Box 8097, Emeryville, CA 94662-8097 (US).

(22) International Filing Date:
4 November 2004 (04.11.2004)

(25) Filing Language: English

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) **Priority Data:**
60/517,337 4 November 2003 (04.11.2003) US
60/525,579 26 November 2003 (26.11.2003) US
60/565,710 27 April 2004 (27.04.2004) US

(71) **Applicant** (*for all designated States except US*): **CHIRON CORPORATION** [US/US]; 4560 Horton Street, Emeryville, CA 94608-2916 (US).

(72) **Inventors; and**

(75) **Inventors/Applicants** (*for US only*): **LONG, Li** [CA/US]; Chiron Corporation, P.O. Box 8097, Emeryville, CA 94662-8097 (US). **LUQMAN, Mohammad** [US/US]; Chiron Corporation, P.O. Box 8097, Emeryville, CA 94662-8097 (US). **YABANNAVAR, Asha** [US/US]; Chiron Corporation, P.O. Box 8097, Emeryville, CA 94662-8097 (US). **ZAROR, Isabel** [US/US]; Chiron Corporation, P.O. Box 8097, Emeryville, CA 94662-8097 (US). **CHEN, Bao-Lu** [US/US]; Chiron Corporation, P.O. Box 8097, Emeryville, CA 94662-8097 (US). **LIU, Xiaofeng** [CN/US]; Chiron Corporation, P.O.Box 8097, Emeryville, CA 94662-8097 (US). **LEE, Sang, Hoon** [KR/US]; Chiron Corporation, P.O.Box 8097, Emeryville,

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) **Title:** ANTAGONIST ANTI-CD40 MONOCLONAL ANTIBODIES AND METHODS FOR THEIR USE

(57) **Abstract:** Methods of therapy for treating diseases mediated by stimulation of CD40 signaling on CD40-expressing cells are provided. The methods comprise administering a therapeutically effective amount of an antagonist anti-CD40 antibody or antigen-binding fragment thereof to a patient in need thereof. The antagonist anti-CD40 antibody or antigen-binding fragment thereof is free of significant agonist activity, but exhibits antagonist activity when the antibody binds a CD40 antigen on a human CD40-expressing cell. Antagonist activity of the anti-CD40 antibody or antigen-binding fragment thereof beneficially inhibits proliferation and/or differentiation of human CD40-expressing cells, such as B cells.



WO 2005/044854 A2

ANTAGONIST ANTI-CD40 MONOCLONAL ANTIBODIES AND METHODS FOR THEIR USE

FIELD OF THE INVENTION

The invention relates to human antibodies capable of binding to CD40, methods of using the antibodies, and methods for treatment of diseases mediated by stimulation of CD40 signaling on CD40-expressing cells.

BACKGROUND OF THE INVENTION

B cells play an important role during the normal *in vivo* immune response. A foreign antigen will bind to surface immunoglobulins on specific B cells, triggering a chain of events including endocytosis, processing, presentation of processed peptides on MHC-class II molecules, and up-regulation of the B7 antigen on the B cell surface. A specific T cell then binds to the B cell via T cell receptor (TCR) recognition of the processed antigen presented on the MHC-class II molecule. Stimulation through the TCR activates the T cell and initiates T-cell cytokine production. A second signal that further activates the T cell is an interaction between the CD28 antigen on T cells and the B7 antigen on B cells. When the above-mentioned signals are received, the CD40 ligand (CD40L or CD154), which is not expressed on resting human T cells, is up-regulated on the T-cell surface. Binding of the CD40 ligand to the CD40 antigen on the B cell surface stimulates the B cell, causing the B cell to mature into a plasma cell secreting high levels of soluble immunoglobulin.

CD40 is a 55 kDa cell-surface antigen present on the surface of both normal and neoplastic human B cells, dendritic cells, antigen presenting cells (APCs), endothelial cells, monocytic and epithelial cells. Transformed cells from patients with low- and high-grade B-cell lymphomas, B-cell acute lymphoblastic leukemia, multiple myeloma,

chronic lymphocytic leukemia, and Hodgkin's disease express CD40. CD40 expression is also detected in two-thirds of acute myeloblastic leukemia cases and 50% of AIDS-related lymphomas. Malignant B cells from several tumors of B-cell lineage express a high degree of CD40 and appear to depend on CD40 signaling for survival and proliferation.

Immunoblastic B-cell lymphomas frequently arise in immunocompromised individuals such as allograft recipients and others receiving long-term immunosuppressive therapy, AIDS patients, and patients with primary immunodeficiency syndromes such as X-linked lymphoproliferative syndrome or Wiscott-Aldrich syndrome (Thomas *et al.* (1991) *Adv. Cancer Res.* 57:329; Straus *et al.* (1993) *Ann. Intern. Med.* 118:45).

The CD40 antigen is related to the human nerve growth factor (NGF) receptor, tumor necrosis factor- α (TNF- α) receptor, and Fas, suggesting that CD40 is a receptor for a ligand with important functions in B-cell activation. CD40 expression on APCs plays an important co-stimulatory role in the activation of both T-helper and cytotoxic T lymphocytes. The CD40 receptor is expressed on activated T cells, activated platelets, and inflamed vascular smooth muscle cells. CD40 receptors can also be found on eosinophils, synovial membranes in rheumatoid arthritis, dermal fibroblasts, and other non-lymphoid cell types. Binding of CD40L to the CD40 receptor stimulates B-cell proliferation and differentiation, antibody production, isotype switching, and B-cell memory generation.

BRIEF SUMMARY OF THE INVENTION

Compositions and methods are provided for treating diseases mediated by stimulation of CD40 signaling on CD40-expressing cells, including lymphomas, autoimmune diseases, and transplant rejections. Compositions include monoclonal antibodies capable of binding to a human CD40 antigen located on the surface of a human CD40-expressing cell, wherein the binding prevents the growth or differentiation of the cell. Compositions also include monoclonal antibodies capable of specifically binding to a human CD40 antigen expressed on the surface of a human CD40-expressing cell, said monoclonal antibody being free of significant agonist activity, wherein

administration of said monoclonal antibody results in significantly less tumor volume than a similar concentration of the chimeric anti-CD20 monoclonal antibody IDEC-C2B8 in a staged nude mouse xenograft tumor model using the Daudi human B cell lymphoma cell line. Compositions also include antigen-binding fragments of these monoclonal antibodies, hybridoma cell lines producing these antibodies, and isolated nucleic acid molecules encoding the amino acid sequences of these monoclonal antibodies. The invention further includes pharmaceutical compositions comprising these anti-CD40 antibodies in a pharmaceutically acceptable carrier.

Methods are provided for preventing or treating a disease mediated by stimulation of CD40 signaling, comprising treating the patient with an anti-CD40 antibody or an antigen-binding fragment thereof that is free of significant agonist activity when bound to a CD40 antigen on a human CD40-expressing cell. Diseases mediated by stimulation of CD40-expressing cells include autoimmune diseases, cancers, and organ and tissue graft rejections. Lymphomas that can be treated or prevented by a method of the present invention include non-Hodgkin's lymphomas (high-grade lymphomas, intermediate-grade lymphomas, and low-grade lymphomas), Hodgkin's disease, acute lymphoblastic leukemias, myelomas, chronic lymphocytic leukemias, and myeloblastic leukemias.

Particular autoimmune diseases contemplated for treatment using the methods of the invention include systemic lupus erythematosus (SLE), rheumatoid arthritis, Crohn's disease, psoriasis, autoimmune thrombocytopenic purpura, multiple sclerosis, ankylosing spondylitis, myasthenia gravis, and pemphigus vulgaris. Such antibodies could also be used to prevent rejection of organ and tissue grafts by suppressing autoimmune responses, to treat lymphomas by depriving malignant B lymphocytes of the activating signal provided by CD40, and to deliver toxins to CD40-bearing cells in a specific manner.

Methods for inhibiting the growth, differentiation, and/or proliferation of human B cells and for inhibiting antibody production by B cells in a human patient are provided, as are methods for inhibiting the growth of cancer cells of a B-cell lineage. Methods for identifying antibodies that have antagonist activity toward CD40-expressing cells are also provided.

The monoclonal antibodies disclosed herein have a strong affinity for CD40 and are characterized by a dissociation equilibrium constant (K_D) of at least 10^{-6} M, preferably at least about 10^{-7} M to about 10^{-8} M, more preferably at least about 10^{-8} M to about 10^{-12} M. Monoclonal antibodies and antigen-binding fragments thereof that are

5 suitable for use in the methods of the invention are capable of specifically binding to a human CD40 antigen expressed on the surface of a human cell. They are free of significant agonist activity but exhibit antagonist activity when bound to CD40 antigen on human cells. In one embodiment, the anti-CD40 antibody or fragment thereof exhibits antagonist activity when bound to CD40 antigen on normal human B cells. In another

10 embodiment, the anti-CD40 antibody or fragment thereof exhibits antagonist activity when bound to CD40 antigen on malignant human B cells. Suitable monoclonal antibodies have human constant regions; preferably they also have wholly or partially humanized framework regions; and most preferably are fully human antibodies or antigen-binding fragments thereof. Examples of such monoclonal antibodies are the

15 antibodies designated herein as CHIR-5.9 and CHIR-12.12; the monoclonal antibodies produced by the hybridoma cell lines designated 131.2F8.5.9 (referred to herein as the cell line 5.9) and 153.8E2.D10.D6.12.12 (referred to herein as the cell line 12.12); a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID

20 NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8; a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID

25 NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5; a monoclonal antibody comprising an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3; and antigen-binding fragments of

30 these monoclonal antibodies that retain the capability of specifically binding to human CD40, and which are free of significant agonist activity but exhibit antagonist activity

when bound to CD40 antigen on human cells. Examples of such monoclonal antibodies also include a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 12.12; a monoclonal antibody that binds to an epitope comprising residues 82-87 of the amino acid sequence shown in SEQ ID NO:10 or SEQ ID NO:12; a monoclonal antibody that competes with the monoclonal antibody CHIR-12.12 in a competitive binding assay; and a monoclonal antibody that is an antigen-binding fragment of the CHIR-12.12 monoclonal antibody or any of the foregoing monoclonal antibodies, where the fragment retains the capability of specifically binding to the human CD40 antigen. Those skilled in the art recognize that the antagonist antibodies and antigen-binding fragments of these antibodies disclosed herein include antibodies and antigen-binding fragments thereof that are produced recombinantly using methods well known in the art and described herein below, and include, for example, monoclonal antibodies CHIR-5.9 and CHIR-12.12 that have been recombinantly produced.

In one embodiment of the invention, methods of treatment comprise administering to a patient a therapeutically effective dose of a pharmaceutical composition comprising suitable antagonist anti-CD40 antibodies or antigen-binding fragments thereof. A therapeutically effective dose of the anti-CD40 antibody or fragment thereof is in the range from about 0.01 mg/kg to about 40 mg/kg, from about 0.01 mg/kg to about 30 mg/kg, from about 0.1 mg/kg to about 30 mg/kg, from about 1 mg/kg to about 30 mg/kg, from about 3 mg/kg to about 30 mg/kg, from about 3 mg/kg to about 25 mg/kg, from about 3 mg/kg to about 20 mg/kg, from about 5 mg/kg to about 15 mg/kg, or from about 7 mg/kg to about 12 mg/kg. It is recognized that the method of treatment may comprise a single administration of a therapeutically effective dose or multiple administrations of a therapeutically effective dose of the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

The antagonist anti-CD40 antibodies identified herein as being suitable for use in the methods of the invention may be modified. Modifications of these antagonist anti-CD40 antibodies include, but are not limited to, immunologically active chimeric anti-CD40 antibodies, humanized anti-CD40 antibodies, and immunologically active murine anti-CD40 antibodies.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows binding of CHIR-5.9 and CHIR-12.12 monoclonal antibodies to CD40 on the surface of lymphoma cell line (Ramos).

Figures 2A and 2B illustrate binding properties of the CHIR-5.9 and CHIR-12.12 monoclonal anti-CD40 antibodies relative to CD40 ligand. Figure 2A shows that binding of CHIR-5.9 and CHIR-12.12 monoclonal antibodies to cell surface CD40 prevents subsequent CD40-ligand binding. Figure 2B shows that the CHIR-5.9 and CHIR-12.12 monoclonal antibodies can compete off CD40 ligand pre-bound to cell surface CD40.

Figures 3A and 3B show ADCC activity of the candidate monoclonal antibodies CHIR-5.9 and CHIR-12.12 against cancer cells from the lymph nodes of non-Hodgkin's lymphoma (NHL) patients. Enriched NK cells from a normal volunteer donor either fresh after isolation (Figure 3A) or after culturing overnight at 37°C (Figure 3B) were used as effector cells in this assay. As NHL cells also express CD20, the target antigen for rituximab (Rituxan®), ADCC activity of the candidate mAbs was compared with that of rituximab.

Figure 4 demonstrates *in vivo* anti-tumor activity of monoclonal antibodies CHIR-5.9 and CHIR-12.12 compared to that of rituximab using an unstaged nude mouse xenograft B cell lymphoma (Namalwa) model.

Figure 5 demonstrates *in vivo* anti-tumor activity of monoclonal antibodies CHIR-5.9 and CHIR-12.12 compared to that of rituximab using an unstaged nude mouse xenograft B cell lymphoma (Daudi) model. RC, resistance to tumor challenge.

Figure 6 demonstrates *in vivo* anti-tumor activity of monoclonal antibodies CHIR-5.9 and CHIR-12.12 compared to that of rituximab using a staged nude mouse xenograft B cell lymphoma (Daudi) model. CR, complete regression.

Figure 7 shows the protocol used for determining the number of CD20 and CD40 molecules on Namalwa and Daudi cells.

Figure 8 shows comparative ADCC of the mAb CHIR-12.12 and rituximab against Daudi lymphoma cells.

Figure 9 sets forth the amino acid sequences for the light and heavy chains of the mAb CHIR-12.12. The leader (residues 1-20 of SEQ ID NO:2), variable (residues 21-132 of SEQ ID NO:2), and constant (residues 133-239 of SEQ ID NO:2) regions of the

light chain are shown in Figure 9A. The leader (residues 1-19 of SEQ ID NO:4), variable (residues 20-139 of SEQ ID NO:4), and constant (residues 140-469 of SEQ ID NO:4) regions of the heavy chain are shown in Figure 9B. The alternative constant region for the heavy chain of the mAb CHIR-12.12 shown in Figure 9B reflects a substitution of a serine residue for the alanine residue at position 153 of SEQ ID NO:4. The complete sequence for this variant of the heavy chain of the mAb CHIR-12.12 is set forth in SEQ ID NO:5.

Figure 10 shows the coding sequence for the light chain (Figure 10A; SEQ ID NO:1) and heavy chain (Figure 10B; SEQ ID NO:3) for the mAb CHIR-12.12.

Figure 11 sets forth the amino acid sequences for the light and heavy chains of mAb CHIR-5.9. The leader (residues 1-20 of SEQ ID NO:6), variable (residues 21-132 of SEQ ID NO:6), and constant (residues 133-239 of SEQ ID NO:6) regions of the light chain are shown in Figure 11A. The leader (residues 1-19 of SEQ ID NO:7), variable (residues 20-144 of SEQ ID NO:7), and constant (residues 145-474 of SEQ ID NO:7) regions of the heavy chain are shown in Figure 11B. The alternative constant region for the heavy chain of the mAb CHIR-5.9 shown in Figure 11B reflects a substitution of a serine residue for the alanine residue at position 158 of SEQ ID NO:7. The complete sequence for this variant of the heavy chain of the mAb CHIR-5.9 is set forth in SEQ ID NO:8.

Figure 12 shows the coding sequence (Figure 12A; SEQ ID NO:9) for the short isoform of human CD40 (amino acid sequence shown in Figure 12B; SEQ ID NO:10), and the coding sequence (Figure 12C; SEQ ID NO:11) for the long isoform of human CD40 (amino acid sequence shown in Figure 12D).

Figure 13 shows thermal melting temperature of CHIR-12.12 in different pH formulations measured by differential scanning calorimetry (DSC).

DETAILED DESCRIPTION OF THE INVENTION

"Tumor," as used herein, refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues.

The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include but are not limited to lymphoma and leukemia.

5 "Antibodies" and "immunoglobulins" (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to an antigen, immunoglobulins include both antibodies and other antibody-like molecules that lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas.

10 The term "antibody" is used in the broadest sense and covers fully assembled antibodies, antibody fragments that can bind antigen (e.g., Fab', F'(ab)₂, Fv, single chain antibodies, diabodies), and recombinant peptides comprising the foregoing.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations
15 that may be present in minor amounts.

"Native antibodies" and "native immunoglobulins" are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy
20 chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain at one end (V_L) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light
25 chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are believed to form an interface between the light- and heavy-chain variable domains.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and
30 specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated

in three segments called complementarity determining regions (CDRs) or hypervariable regions both in the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework (FR) regions. The variable domains of native heavy and light chains each comprise four FR regions, largely
5 adopting a β -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming part of, the β -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.* (1991) *NIH Publ. No. 91-3242*, Vol. I, pages 647-669).

10 The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as Fc receptor (FcR) binding, participation of the antibody in antibody-dependent cellular toxicity, opsonization, initiation of complement dependent cytotoxicity, and mast cell degranulation.

The term "hypervariable region" when used herein refers to the amino acid
15 residues of an antibody that are responsible for antigen binding. The hypervariable region comprises amino acid residues from a "complementarity determining region" or "CDR" (i.e., residues 24-34 (L1); 50-56 (L2), and 89-97 (L3) in the light-chain variable domain and 31-35 (H1), 50-65 (H2), and 95-102 (H3) in the heavy-chain variable domain; Kabat *et al.* (1991) *Sequences of Proteins of Immunological Interest* (5th ed., Public Health
20 Service, National Institute of Health, Bethesda, MD) and/or those residues from a "hypervariable loop" (i.e., residues 26-32(L1), 50-52 (L2), and 91-96 (L3) in the light-chain variable domain and 26-32(H1), 53-55 (H2), and 96-101 (H3) in the heavy-chain variable domain; Clothia and Lesk (1987) *J. Mol. Biol.* 196:901-917). "Framework" or "FR" residues are those variable domain residues other than the hypervariable region
25 residues.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen-binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata *et al.* (1995) *Protein Eng.* 8(10):1057-1062); single-chain antibody molecules; and
30 multispecific antibodies formed from antibody fragments. Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a

single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment that contains a complete antigen
5 recognition and binding site. In a two-chain Fv species, this region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. In a single-chain Fv species, one heavy- and one light-chain variable domain can be covalently linked by flexible peptide linker such that the light and heavy chains can associate in a "dimeric" structure analogous to that in a two-chain Fv species. It is in this
10 configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

15 The Fab fragment also contains the constant domain of the light chain and the first constant domain (C_H1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy-chain C_H1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear
20 a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ),
25 based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of human immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and
30 IgA2. The heavy-chain constant domains that correspond to the different classes of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively. The

subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known. Different isotypes have different effector functions. For example, human IgG1 and IgG3 isotypes mediate antibody-dependent cell-mediated cytotoxicity (ADCC) activity.

5 The word "label" when used herein refers to a detectable compound or composition that is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable. Radionuclides that can serve
10 as detectable labels include, for example, I-131, I-123, I-125, Y-90, Re-188, Re-186, At-211, Cu-67, Bi-212, and Pd-109. The label might also be a non-detectable entity such as a toxin.

 The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native target
15 disclosed herein or the transcription or translation thereof.

 "Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers that are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include
20 buffers such as phosphate, citrate, succinate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose,
25 mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN, polyethylene glycol (PEG), and Pluronic. Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

30 A "host cell," as used herein, refers to a microorganism or a eukaryotic cell or cell line cultured as a unicellular entity that can be, or has been, used as a recipient for a

recombinant vector or other transfer polynucleotides, and include the progeny of the original cell that has been transfected. It is understood that the progeny of a single cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

5 "Human effector cells" are leukocytes that express one or more FcRs and perform effector functions. Preferably, the cells express at least Fc γ RIII and carry out antigen-dependent cell-mediated cytotoxicity (ADCC) effector function. Examples of human leukocytes that mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, macrophages, eosinophils, and neutrophils, with
10 PBMCs and NK cells being preferred. Antibodies that have ADCC activity are typically of the IgG1 or IgG3 isotype. Note that in addition to isolating IgG1 and IgG3 antibodies, such ADCC-mediating antibodies can be made by engineering a variable region from a non-ADCC antibody or variable region fragment to an IgG1 or IgG3 isotype constant region.

15 The terms "Fc receptor" or "FcR" are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native-sequence human FcR. Moreover, a preferred FcR is one that binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and Fc γ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc γ RII receptors include Fc γ RIIA (an
20 "activating receptor") and Fc γ RIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor Fc γ RIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor Fc γ RIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain (see Daeron (1997)
25 *Annu. Rev. Immunol.* 15:203-234). FcRs are reviewed in Ravetch and Kinet (1991) *Annu. Rev. Immunol.* 9:457-492 (1991); Capel *et al.* (1994) *Immunomethods* 4:25-34; and de Haas *et al.* (1995) *J. Lab. Clin. Med.* 126:330-341. Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal
30 IgGs to the fetus (Guyer *et al.* (1976) *J. Immunol.* 117:587 and Kim *et al.* (1994) *J. Immunol.* 24:249 (1994)).

There are a number of ways to make human antibodies. For example, secreting cells can be immortalized by infection with the Epstein-Barr virus (EBV). However, EBV-infected cells are difficult to clone and usually produce only relatively low yields of immunoglobulin (James and Bell (1987) *J. Immunol. Methods* 100:5-40). In the future, the immortalization of human B cells might possibly be achieved by introducing a defined combination of transforming genes. Such a possibility is highlighted by a recent demonstration that the expression of the telomerase catalytic subunit together with the SV40 large oncoprotein and an oncogenic allele of H-ras resulted in the tumorigenic conversion of normal human epithelial and fibroblast cells (Hahn *et al.* (1999) *Nature* 400:464-468). It is now possible to produce transgenic animals (e.g., mice) that are capable, upon immunization, of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production (Jakobovits *et al.* (1993) *Nature* 362:255-258; Lonberg and Huszar (1995) *Int. Rev. Immunol.* 13:65-93; Fishwild *et al.* (1996) *Nat. Biotechnol.* 14:845-851; Mendez *et al.* (1997) *Nat. Genet.* 15:146-156; Green (1999) *J. Immunol. Methods* 231:11-23; Tomizuka *et al.* (2000) *Proc. Natl. Acad. Sci. USA* 97:722-727; reviewed in Little *et al.* (2000) *Immunol. Today* 21:364-370). For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production (Jakobovits *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:2551-2555). Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice results in the production of human antibodies upon antigen challenge (Jakobovits *et al.* (1993) *Nature* 362:255-258). Mendez *et al.* (1997) (*Nature Genetics* 15:146-156) have generated a line of transgenic mice that, when challenged with an antigen, generates high affinity fully human antibodies. This was achieved by germ-line integration of megabase human heavy-chain and light-chain loci into mice with deletion into endogenous J_H segment as described above. These mice (XenoMouse[®] II technology (Abgenix; Fremont, California)) harbor 1,020 kb of human heavy-chain locus containing approximately 66 V_H genes, complete D_H and J_H regions, and three different constant regions, and also harbors 800 kb of human κ locus containing 32 $V\kappa$ genes, $J\kappa$ segments, and $C\kappa$ genes. The antibodies produced in these mice closely resemble that seen in humans in all respects, including gene rearrangement, assembly, and repertoire.

The human antibodies are preferentially expressed over endogenous antibodies due to deletion in endogenous segment that prevents gene rearrangement in the murine locus. Such mice may be immunized with an antigen of particular interest.

Sera from such immunized animals may be screened for antibody reactivity
5 against the initial antigen. Lymphocytes may be isolated from lymph nodes or spleen cells and may further be selected for B cells by selecting for CD138-negative and CD19-positive cells. In one aspect, such B cell cultures (BCCs) may be fused to myeloma cells to generate hybridomas as detailed above.

In another aspect, such B cell cultures may be screened further for reactivity
10 against the initial antigen, preferably. Such screening includes ELISA with the target/antigen protein, a competition assay with known antibodies that bind the antigen of interest, and in vitro binding to transiently transfected CHO or other cells that express the target antigen.

The present invention is directed to compositions and methods for treating human
15 patients with diseases mediated by stimulation of CD40 signaling on CD40-expressing cells. The methods involve treatment with an anti-CD40 antibody of the invention, or an antigen-binding fragment thereof, where administration of the antibody or antigen-binding fragment thereof promotes a positive therapeutic response within the patient undergoing this method of therapy. Anti-CD40 antibodies suitable for use in the methods
20 of the invention specifically bind a human CD40 antigen expressed on the surface of a human cell and are free of significant agonist activity, but exhibit antagonist activity when bound to the CD40 antigen on a human CD40-expressing cell. These anti-CD40 antibodies and antigen-binding fragments thereof are referred to herein as antagonist anti-CD40 antibodies. Such antibodies include, but are not limited to, the fully human
25 monoclonal antibodies CHIR-5.9 and CHIR-12.12 described below and monoclonal antibodies having the binding characteristics of monoclonal antibodies CHIR-5.9 and CHIR-12.12. Those skilled in the art recognize that the antagonist antibodies and antigen-binding fragments of these antibodies disclosed herein include antibodies and antigen-binding fragments thereof that are produced recombinantly using methods well
30 known in the art and described herein below, and include, for example, monoclonal antibodies CHIR-5.9 and CHIR-12.12 that have been recombinantly produced.

Antibodies that have the binding characteristics of monoclonal antibodies CHIR-5.9 and CHIR-12.12 include antibodies that competitively interfere with binding CD40 and/or bind the same epitopes as CHIR-5.9 and CHIR-12.12. One of skill could determine whether an antibody competitively interferes with CHIR-5.9 or CHIR-12.12 using standard methods.

When these antibodies bind CD40 displayed on the surface of human cells, such as human B cells, the antibodies are free of significant agonist activity; in some embodiments, their binding to CD40 displayed on the surface of human cells results in inhibition of proliferation and differentiation of these human cells. Thus, the antagonist anti-CD40 antibodies suitable for use in the methods of the invention include those monoclonal antibodies that can exhibit antagonist activity toward normal and malignant human cells expressing the cell-surface CD40 antigen.

In some embodiments, the anti-CD40 antibodies of the invention exhibit increased anti-tumor activity relative to the chimeric anti-CD20 monoclonal antibody IDEC-C2B8, where anti-tumor activity is assayed with equivalent amounts of these antibodies in a nude mouse xenograft tumor model using human lymphoma cell lines. IDEC-C2B8 (IDEC Pharmaceuticals Corp., San Diego, California; commercially available under the tradename Rituxan®, also referred to as rituximab) is a chimeric anti-CD20 monoclonal antibody containing human IgG1 and kappa constant regions with murine variable regions isolated from a murine anti-CD20 monoclonal antibody, IDEC-2B8 (Reff *et al.* (1994) *Blood* 83:435-445). Rituximab® is licensed for treatment of relapsed B cell low-grade or follicular non-Hodgkin's lymphoma (NHL). The discovery of antibodies with superior anti-tumor activity compared to Rituximab® could drastically improve methods of cancer therapy for B cell lymphomas, particularly B cell non-Hodgkin's lymphoma.

Suitable nude mouse xenograft tumor models include those using the human Burkitt's lymphoma cell lines known as Namalwa and Daudi. Preferred embodiments assay anti-tumor activity in a staged nude mouse xenograft tumor model using the Daudi human lymphoma cell line as described herein below in Example 17. A staged nude mouse xenograft tumor model using the Daudi lymphoma cell line is more effective at distinguishing the therapeutic efficacy of a given antibody than is an unstaged model, as

in the staged model antibody dosing is initiated only after the tumor has reached a measurable size. In the unstaged model, antibody dosing is initiated generally within about 1 day of tumor inoculation and before a palpable tumor is present. The ability of an antibody to outperform Rituxan® (i.e., to exhibit increased anti-tumor activity) in a staged model is a strong indication that the antibody will be more therapeutically effective than Rituxan®. Moreover, in the Daudi model, anti-CD20, the target for Rituxan® is expressed on the cell surface at a higher level than is CD40.

By "equivalent amount" of the anti-CD40 antibody of the invention and Rituxan® is intended the same mg dose is administered on a per weight basis. Thus, where the anti-CD40 antibody of the invention is dosed at 0.01 mg/kg body weight of the mouse used in the tumor model, Rituxan® is also dosed at 0.01 mg/kg body weight of the mouse. Similarly, where the anti-CD40 antibody of the invention is dosed at 0.1, 1, or 10 mg/kg body weight of the mouse used in the tumor model, the Rituxan® is also dosed at 0.1, 1, or 10 mg/kg, respectively, of the body weight of the mouse.

When administered in the nude mouse xenograft tumor model, some antibodies of the invention result in significantly less tumor volume than an equivalent amount of Rituxan®. Thus, for example, the fully human monoclonal antibody CHIR-12.12 exhibits at least a 20% increase in anti-tumor activity relative to that observed with an equivalent dose of Rituxan when assayed in the staged nude mouse xenograft tumor model using the Daudi human lymphoma cell line in the manner described in Examples herein below, and can exhibit as much as a 50% to 60% increase in anti-tumor activity in this assay. This increased anti-tumor activity is reflected in the greater reduction in tumor volume observed with the anti-CD40 antibody of the invention when compared to the equivalent dose of Rituxan®. Thus, for example, depending upon the length of time after tumor inoculation, the monoclonal antibody CHIR-12.12 can exhibit a tumor volume that is about one-third to about one-half that observed for an equivalent dose of Rituxan®.

Another difference in antibody efficacy is to measure *in vitro* the concentration of antibody needed to obtain the maximum lysis of tumor cells *in vitro* in the presence of NK cells. For example, the anti-CD40 antibodies of the invention reach maximum lysis

of Daudi cells at an EC₅₀ of less than 1/2, and preferably 1/4, and most preferably, 1/10 the concentration of Rituxan®.

In addition to the monoclonal antibody CHIR-12.12, other anti-CD40 antibodies that would share the characteristics of having significantly greater efficacy than equivalent amounts of Rituxan® in the assays described above include, but are not limited to: (1) the monoclonal antibody produced by the hybridoma cell line 12.12; (2) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence in SEQ ID NO:2, the sequence in SEQ ID NO:4, the sequence in SEQ ID NO:5, both the sequence in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence in SEQ ID NO:2 and SEQ ID NO:5; (3) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the nucleotide sequence in SEQ ID NO:1, the nucleotide sequence in SEQ ID NO:3, and both the sequence in SEQ ID NO:1 and SEQ ID NO:3; (4) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 12.12; (5) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the amino acid sequence in SEQ ID NO:10 or SEQ ID NO:12; (6) a monoclonal antibody that competes with the monoclonal antibody CHIR-12.12 in a competitive binding assay; and (7) a monoclonal antibody that is an antigen-binding fragment of the CHIR-12.12 monoclonal antibody or the foregoing monoclonal antibodies in preceding items (1)-(6), where the fragment retains the capability of specifically binding to the human CD40 antigen.

Antagonist Anti-CD40 Antibodies

The monoclonal antibodies CHIR-5.9 and CHIR-12.12 represent suitable antagonist anti-CD40 antibodies for use in the methods of the present invention. The CHIR-5.9 and CHIR-12.12 antibodies are fully human anti-CD40 monoclonal antibodies of the IgG₁ isotype produced from the hybridoma cell lines 131.2F8.5.9 (referred to herein as the cell line 5.9) and 153.8E2.D10.D6.12.12 (referred to herein as the cell line 12.12). These cell lines were created using splenocytes from immunized xenotypic mice containing the human IgG₁ heavy chain locus and the human κ chain locus (XenoMouse® technology; Abgenix; Fremont, California). The spleen cells were fused with the mouse

myeloma SP2/0 cells (Sierra BioSource). The resulting hybridomas were sub-cloned several times to create the stable monoclonal cell lines 5.9 and 12.12. Other antibodies of the invention may be prepared similarly using mice transgenic for human immunoglobulin loci or by other methods known in the art and/or described herein.

5 The nucleotide and amino acid sequences of the variable regions of the CHIR-12.12 antibody, and the amino acid sequences of the variable regions of the CHIR-5.9 antibody, are disclosed. More particularly, the amino acid sequences for the leader, variable, and constant regions for the light chain and heavy chain for mAb CHIR-12.12 are set forth in Figures 9A and 9B, respectively. See also SEQ ID NO:2 (complete
10 sequence for the light chain of mAb CHIR-12.12), SEQ ID NO:4 (complete sequence for the heavy chain for mAb CHIR-12.12), and SEQ ID NO:5 (complete sequence for a variant of the heavy chain for mAb CHIR-12.12 set forth in SEQ ID NO:4, where the variant comprises a serine substitution for the alanine residue at position 153 of SEQ ID NO:4). The nucleotide sequences encoding the light chain and heavy chain for mAb
15 CHIR-12.12 are set forth in Figures 11A and 11B, respectively. See also SEQ ID NO:1 (coding sequence for the light chain for mAb CHIR-12.12), and SEQ ID NO:3 (coding sequence for the heavy chain for mAb CHIR-12.12). The amino acid sequences for the leader, variable, and constant regions for the light chain and heavy chain of the CHIR-5.9 mAb are set forth in Figures 10A and 10B, respectively. See also SEQ ID NO:6
20 (complete sequence for the light chain of mAb CHIR-5.9), SEQ ID NO:7 (complete sequence for the heavy chain of mAb CHIR-5.9), and SEQ ID NO:8 (complete sequence for a variant of the heavy chain of mAb CHIR-5.9 set forth in SEQ ID NO:7, where the variant comprises a serine substitution for the alanine residue at position 158 of SEQ ID NO:7). Further, hybridomas expressing CHIR-5.9 and CHIR-12.12 antibodies have been
25 deposited with the ATCC with a patent deposit designation of PTA-5542 and PTA-5543, respectively.

 In addition to antagonist activity, it is preferable that anti-CD40 antibodies of this invention have another mechanism of action against a tumor cell. For example, native CHIR-5.9 and CHIR-12.12 antibodies have ADCC activity. Alternatively, the variable
30 regions of the CHIR-5.9 and CHIR-12.12 antibodies can be expressed on another antibody isotype that has ADCC activity. It is also possible to conjugate native forms,

recombinant forms, or antigen-binding fragments of CHIR-5.9 or CHIR-12.12 to a cytotoxin, a therapeutic agent, or a radioactive metal ion or radioisotope, as noted herein below.

The CHIR-5.9 and CHIR-12.12 monoclonal antibodies bind soluble CD40 in ELISA-type assays, prevent the binding of CD40-ligand to cell-surface CD40, and displace the pre-bound CD40-ligand, as determined by flow cytometric assays. Antibodies CHIR-5.9 and CHIR-12.12 compete with each other for binding to CD40 but not with 15B8, the anti-CD40 monoclonal antibody described in U.S. Provisional Application Serial No. 60/237,556, titled "*Human Anti-CD40 Antibodies*," filed October 2, 2000, and PCT International Application No. PCT/US01/30857, also titled "*Human Anti-CD40 Antibodies*," filed October 2, 2001 (Attorney Docket No. PP16092.003), both of which are herein incorporated by reference in their entirety. When tested *in vitro* for effects on proliferation of B cells from normal human subjects, CHIR-5.9 and CHIR-12.12 act as antagonist anti-CD40 antibodies. Furthermore, CHIR-5.9 and CHIR-12.12 do not induce strong proliferation of human lymphocytes from normal subjects. These antibodies are able to kill CD40-expressing target cells by antibody dependent cellular cytotoxicity (ADCC). The binding affinity of CHIR-5.9 for human CD40 is 1.2×10^{-8} M and the binding affinity of CHIR-12.12 is 5×10^{-10} M, as determined by the Biacore™ assay.

Suitable antagonist anti-CD40 antibodies for use in the methods of the present invention exhibit a strong single-site binding affinity for the CD40 cell-surface antigen. The monoclonal antibodies of the invention exhibit a dissociation equilibrium constant (K_D) for CD40 of at least 10^{-5} M, at least 3×10^{-5} M, preferably at least 10^{-6} M to 10^{-7} M, more preferably at least 10^{-8} M to about 10^{-12} M, measured using a standard assay such as Biacore™. Biacore analysis is known in the art and details are provided in the "BIAapplications handbook." Methods described in WO 01/27160 can be used to modulate the binding affinity.

By "CD40 antigen," "CD40 cell surface antigen," "CD40 receptor," or "CD40" is intended a transmembrane glycoprotein that belongs to the tumor necrosis factor (TNF) receptor family (see, for example, U.S. Patent Nos. 5,674,492 and 4,708,871; Stamenkovic *et al.* (1989) *EMBO* 8:1403; Clark (1990) *Tissue Antigens* 36:33; Barclay *et*

al. (1997) *The Leucocyte Antigen Facts Book* (2d ed.; Academic Press, San Diego)).

Two isoforms of human CD40, encoded by alternatively spliced transcript variants of this gene, have been identified. The first isoform (also known as the "long isoform" or "isoform 1") is expressed as a 277-amino-acid precursor polypeptide (SEQ ID NO:12 (first reported as GenBank Accession No. CAA43045, and identified as isoform 1 in GenBank Accession No. NP_001241), encoded by SEQ ID NO:11 (see GenBank Accession Nos. X60592 and NM_001250)), which has a signal sequence represented by the first 19 residues. The second isoform (also known as the "short isoform" or "isoform 2") is expressed as a 203-amino-acid precursor polypeptide (SEQ ID NO:10 (GenBank Accession No. NP_690593), encoded by SEQ ID NO:9 (GenBank Accession No. NM_152854)), which also has a signal sequence represented by the first 19 residues. The precursor polypeptides of these two isoforms of human CD40 share in common their first 165 residues (i.e., residues 1-165 of SEQ ID NO:10 and SEQ ID NO:12). The precursor polypeptide of the short isoform (shown in SEQ ID NO:10) is encoded by a transcript variant (SEQ ID NO:9) that lacks a coding segment, which leads to a translation frame shift; the resulting CD40 isoform contains a shorter and distinct C-terminus (residues 166-203 of SEQ ID NO:10) from that contained in the long isoform of CD40 (C-terminus shown in residues 166-277 of SEQ ID NO:12). For purposes of the present invention, the term "CD40 antigen," "CD40 cell surface antigen," "CD40 receptor," or "CD40" encompasses both the short and long isoforms of CD40. The anti-CD40 antibodies of the present invention bind to an epitope of human CD40 that resides at the same location within either the short isoform or long isoform of this cell surface antigen as noted herein below.

The CD40 antigen is displayed on the surface of a variety of cell types, as described elsewhere herein. By "displayed on the surface" and "expressed on the surface" is intended that all or a portion of the CD40 antigen is exposed to the exterior of the cell. The displayed or expressed CD40 antigen may be fully or partially glycosylated.

By "agonist activity" is intended that the substance functions as an agonist. An agonist combines with a receptor on a cell and initiates a reaction or activity that is similar to or the same as that initiated by the receptor's natural ligand. An agonist of CD40 induces any or all of, but not limited to, the following responses: B cell

proliferation and differentiation, antibody production, intercellular adhesion, B cell memory generation, isotype switching, up-regulation of cell-surface expression of MHC Class II and CD80/86, and secretion of pro-inflammatory cytokines such as IL-8, IL-12, and TNF. By "antagonist activity" is intended that the substance functions as an antagonist. An antagonist of CD40 prevents or reduces induction of any of the responses induced by binding of the CD40 receptor to an agonist ligand, particularly CD40L. The antagonist may reduce induction of any one or more of the responses to agonist binding by 5%, 10%, 15%, 20%, 25%, 30%, 35%, preferably 40%, 45%, 50%, 55%, 60%, more preferably 70%, 80%, 85%, and most preferably 90%, 95%, 99%, or 100%. Methods for measuring anti-CD40 antibody and CD40-ligand binding specificity and antagonist activity are known to one of skill in the art and include, but are not limited to, standard competitive binding assays, assays for monitoring immunoglobulin secretion by B cells, B cell proliferation assays, Banchereau-Like-B cell proliferation assays, T cell helper assays for antibody production, co-stimulation of B cell proliferation assays, and assays for up-regulation of B cell activation markers. See, for example, such assays disclosed in WO 00/75348 and U.S. Patent No. 6,087,329, herein incorporated by reference.

By "significant" agonist activity is intended an agonist activity of at least 30%, 35%, 40%, 45%, 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% greater than the agonist activity induced by a neutral substance or negative control as measured in an assay of a B cell response. Preferably, "significant" agonist activity is an agonist activity that is at least 2-fold greater or at least 3-fold greater than the agonist activity induced by a neutral substance or negative control as measured in an assay of a B cell response. Thus, for example, where the B cell response of interest is B cell proliferation, "significant" agonist activity would be induction of a level of B cell proliferation that is at least 2-fold greater or at least 3-fold greater than the level of B cell proliferation induced by a neutral substance or negative control. In one embodiment, a non-specific immunoglobulin, for example IgG1, that does not bind to CD40 serves as the negative control. A substance "free of significant agonist activity" would exhibit an agonist activity of not more than about 25% greater than the agonist activity induced by a neutral substance or negative control, preferably not more than about 20% greater, 15% greater, 10% greater, 5% greater, 1% greater, 0.5% greater, or even not more than about 0.1%

greater than the agonist activity induced by a neutral substance or negative control as measured in an assay of a B cell response. The antagonist anti-CD40 antibodies useful in the methods of the present invention are free of significant agonist activity as noted above when bound to a CD40 antigen on a human cell. In one embodiment of the invention, the antagonist anti-CD40 antibody is free of significant agonist activity in one B cell response. In another embodiment of the invention, the antagonist anti-CD40 antibody is free of significant agonist activity in assays of more than one B cell response (e.g., proliferation and differentiation, or proliferation, differentiation, and antibody production).

As used herein "anti-CD40 antibody" encompasses any antibody that specifically recognizes the CD40 B cell surface antigen, including polyclonal antibodies, monoclonal antibodies, single-chain antibodies, and fragments thereof such as Fab, F(ab')₂, F_v, and other fragments which retain the antigen binding function of the parent anti-CD40 antibody. Of particular interest to the present invention are the antagonist anti-CD40 antibodies disclosed herein that share the binding characteristics of the monoclonal antibodies CHIR-5.9 and CHIR-12.12 described above. Such antibodies include, but are not limited to the following: (1) the monoclonal antibodies produced by the hybridoma cell lines designated 131.2F8.5.9 (referred to herein as the cell line 5.9) and 153.8E2.D10.D6.12.12 (referred to herein as the cell line 12.12), deposited with the ATCC as Patent Deposit No. PTA-5542 and Patent Deposit No. PTA-5543, respectively; (2) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequences shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequences shown in SEQ ID NO:2 and SEQ ID NO:5; (3) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequences shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequences shown in SEQ ID NO:6 and SEQ ID NO:8; (4) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the nucleotide sequence shown in SEQ ID NO:1, the nucleotide sequence shown in SEQ ID

NO:3, and both the sequences shown in SEQ ID NO:1 and SEQ ID NO:3; (5) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or the hybridoma cell line 12.12; (6) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the amino acid sequence shown in SEQ ID NO:10 or SEQ ID NO:12; (7) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay; and (8) a monoclonal antibody that is an antigen-binding fragment of the CHIR-12.12 or CHIR-5.9 monoclonal antibody or the foregoing monoclonal antibodies in preceding items (1)-(7), where the fragment retains the capability of specifically binding to the human CD40 antigen.

Production of Antagonist Anti-CD40 Antibodies

The antagonist anti-CD40 antibodies disclosed herein and for use in the methods of the present invention can be produced using any antibody production method known to those of skill in the art. Thus, polyclonal sera may be prepared by conventional methods. In general, a solution containing the CD40 antigen is first used to immunize a suitable animal, preferably a mouse, rat, rabbit, or goat. Rabbits or goats are preferred for the preparation of polyclonal sera due to the volume of serum obtainable, and the availability of labeled anti-rabbit and anti-goat antibodies.

Polyclonal sera can be prepared in a transgenic animal, preferably a mouse bearing human immunoglobulin loci. In a preferred embodiment, Sf9 cells expressing CD40 are used as the immunogen. Immunization can also be performed by mixing or emulsifying the antigen-containing solution in saline, preferably in an adjuvant such as Freund's complete adjuvant, and injecting the mixture or emulsion parenterally (generally subcutaneously or intramuscularly). A dose of 50-200 µg/injection is typically sufficient. Immunization is generally boosted 2-6 weeks later with one or more injections of the protein in saline, preferably using Freund's incomplete adjuvant. One may alternatively generate antibodies by *in vitro* immunization using methods known in the art, which for the purposes of this invention is considered equivalent to *in vivo* immunization. Polyclonal antisera are obtained by bleeding the immunized animal into a glass or plastic container, incubating the blood at 25°C for one hour, followed by incubating at 4°C for 2-

18 hours. The serum is recovered by centrifugation (e.g., 1,000 x g for 10 minutes). About 20-50 ml per bleed may be obtained from rabbits.

Production of the Sf 9 (*Spodoptera frugiperda*) cells is disclosed in U.S. Patent No. 6,004,552, incorporated herein by reference. Briefly, sequences encoding human CD40 were recombined into a baculovirus using transfer vectors. The plasmids were co-transfected with wild-type baculovirus DNA into Sf 9 cells. Recombinant baculovirus-infected Sf 9 cells were identified and clonally purified.

Preferably the antibody is monoclonal in nature. By "monoclonal antibody" is intended an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. The term is not limited regarding the species or source of the antibody. The term encompasses whole immunoglobulins as well as fragments such as Fab, F(ab')₂, Fv, and others which retain the antigen binding function of the antibody. Monoclonal antibodies are highly specific, being directed against a single antigenic site, i.e., the CD40 cell surface antigen in the present invention. Furthermore, in contrast to conventional (polyclonal) antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.* (1975) *Nature* 256:495, or may be made by recombinant DNA methods (see, e.g., U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in, for example, Clackson *et al.* (1991) *Nature* 352:624-628; Marks *et al.* (1991) *J. Mol. Biol.* 222:581-597; and U.S. Patent No. 5,514,548.

By "epitope" is intended the part of an antigenic molecule to which an antibody is produced and to which the antibody will bind. Epitopes can comprise linear amino acid residues (i.e., residues within the epitope are arranged sequentially one after another in a

linear fashion), nonlinear amino acid residues (referred to herein as "nonlinear epitopes"; these epitopes are not arranged sequentially), or both linear and nonlinear amino acid residues.

Monoclonal antibodies can be prepared using the method of Kohler *et al.* (1975) *Nature* 256:495-496, or a modification thereof. Typically, a mouse is immunized with a solution containing an antigen. Immunization can be performed by mixing or emulsifying the antigen-containing solution in saline, preferably in an adjuvant such as Freund's complete adjuvant, and injecting the mixture or emulsion parenterally. Any method of immunization known in the art may be used to obtain the monoclonal antibodies of the invention. After immunization of the animal, the spleen (and optionally, several large lymph nodes) are removed and dissociated into single cells. The spleen cells may be screened by applying a cell suspension to a plate or well coated with the antigen of interest. The B cells expressing membrane bound immunoglobulin specific for the antigen bind to the plate and are not rinsed away. Resulting B cells, or all dissociated spleen cells, are then induced to fuse with myeloma cells to form hybridomas, and are cultured in a selective medium. The resulting cells are plated by serial dilution and are assayed for the production of antibodies that specifically bind the antigen of interest (and that do not bind to unrelated antigens). The selected monoclonal antibody (mAb)-secreting hybridomas are then cultured either *in vitro* (e.g., in tissue culture bottles or hollow fiber reactors), or *in vivo* (as ascites in mice).

Where the antagonist anti-CD40 antibodies of the invention are to be prepared using recombinant DNA methods, the DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells described herein serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese Hamster Ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. Review articles on recombinant expression in bacteria of DNA encoding the antibody include Skerra *et al.* (1993) *Curr. Opinion in Immunol.* 5:256 and

Phickthun (1992) *Immunol. Revs.* 130:151. Alternatively, antibody can be produced in a cell line such as a CHO cell line, as disclosed in U.S. Patent Nos. 5,545,403; 5,545,405; and 5,998,144; incorporated herein by reference. Briefly the cell line is transfected with vectors capable of expressing a light chain and a heavy chain, respectively. By
5 transfecting the two proteins on separate vectors, chimeric antibodies can be produced. Another advantage is the correct glycosylation of the antibody.

In some embodiments, the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 antibody, or antigen-binding fragment thereof is produced in CHO cells using the GS gene expression system (Lonza Biologics, Portsmouth, New
10 Hampshire), which uses glutamine synthetase as a marker. See, also U.S. Patent Nos. 5,122,464; 5,591,639; 5,658,759; 5,770,359; 5,827,739; 5,879,936; 5,891,693; and 5,981,216; the contents of which are herein incorporated by reference in their entirety.

Monoclonal antibodies to CD40 are known in the art. See, for example, the sections dedicated to B-cell antigen in McMichael, ed. (1987; 1989) *Leukocyte Typing III and IV* (Oxford University Press, New York); U.S. Patent Nos. 5,674,492; 5,874,082; 5,677,165; 6,056,959; WO 00/63395; International Publication Nos. WO 02/28905 and WO 02/28904; Gordon *et al.* (1988) *J. Immunol.* 140:1425; Valle *et al.* (1989) *J. Immunol.* 19:1463; Clark *et al.* (1986) *PNAS* 83:4494; Paulie *et al.* (1989) *J. Immunol.* 142:590; Gordon *et al.* (1987) *Eur. J. Immunol.* 17:1535; Jabara *et al.* (1990) *J. Exp.*
20 *Med.* 172:1861; Zhang *et al.* (1991) *J. Immunol.* 146:1836; Gascan *et al.* (1991) *J. Immunol.* 147:8; Banachereau *et al.* (1991) *Clin. Immunol. Spectrum* 3:8; and Banachereau *et al.* (1991) *Science* 251:70; all of which are herein incorporated by reference. Of particular interest to the present invention are the antagonist anti-CD40 antibodies disclosed herein that share the binding characteristics of the monoclonal antibodies
25 CHIR-5.9 and CHIR-12.12 described above.

The term "CD40-antigen epitope" as used herein refers to a molecule that is capable of immunoreactivity with the anti-CD40 monoclonal antibodies of this invention, excluding the CD40 antigen itself. CD40-antigen epitopes may comprise proteins, protein fragments, peptides, carbohydrates, lipids, and other molecules, but for the
30 purposes of the present invention are most commonly proteins, short oligopeptides, oligopeptide mimics (i.e., organic compounds which mimic the antibody binding

properties of the CD40 antigen), or combinations thereof. Suitable oligopeptide mimics are described, inter alia, in PCT application US 91/04282.

Additionally, the term "anti-CD40 antibody" as used herein encompasses chimeric anti-CD40 antibodies; such chimeric anti-CD40 antibodies for use in the methods of the invention have the binding characteristics of the CHIR-5.9 and CHIR-12.12 monoclonal antibodies described herein. By "chimeric" antibodies is intended antibodies that are most preferably derived using recombinant deoxyribonucleic acid techniques and which comprise both human (including immunologically "related" species, e.g., chimpanzee) and non-human components. Thus, the constant region of the chimeric antibody is most preferably substantially identical to the constant region of a natural human antibody; the variable region of the chimeric antibody is most preferably derived from a non-human source and has the desired antigenic specificity to the CD40 cell-surface antigen. The non-human source can be any vertebrate source that can be used to generate antibodies to a human CD40 cell-surface antigen or material comprising a human CD40 cell-surface antigen. Such non-human sources include, but are not limited to, rodents (e.g., rabbit, rat, mouse, etc.; see, for example, U.S. Patent No. 4,816,567, herein incorporated by reference) and non-human primates (e.g., Old World Monkey, Ape, etc.; see, for example, U.S. Patent Nos. 5,750,105 and 5,756,096; herein incorporated by reference). As used herein, the phrase "immunologically active" when used in reference to chimeric anti-CD40 antibodies means a chimeric antibody that binds human CD40.

Chimeric and humanized anti-CD40 antibodies are also encompassed by the term anti-CD40 antibody as used herein. Chimeric antibodies comprise segments of antibodies derived from different species. Rituxan® is an example of a chimeric antibody with a murine variable region and a human constant region.

By "humanized" is intended forms of anti-CD40 antibodies that contain minimal sequence derived from non-human immunoglobulin sequences. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region (also known as complementarity determining region or CDR) of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit, or nonhuman primate having

the desired specificity, affinity, and capacity. The phrase "complementarity determining region" refers to amino acid sequences which together define the binding affinity and specificity of the natural Fv region of a native immunoglobulin binding site. See, e.g., Chothia *et al* (1987) *J. Mol. Biol.* 196:901-917; Kabat *et al* (1991) U. S. Dept. of Health and Human Services, NIH Publication No. 91-3242). The phrase "constant region" refers to the portion of the antibody molecule that confers effector functions. In previous work directed towards producing non-immunogenic antibodies for use in therapy of human disease, mouse constant regions were substituted by human constant regions. The constant regions of the subject humanized antibodies were derived from human immunoglobulins. However, these humanized antibodies still elicited an unwanted and potentially dangerous immune response in humans and there was a loss of affinity. Humanized anti-CD40 antibodies for use in the methods of the present invention have binding characteristics similar to those exhibited by the CHIR-5.9 and CHIR-12.12 monoclonal antibodies described herein.

Humanization can be essentially performed following the method of Winter and co-workers (Jones *et al.* (1986) *Nature* 321:522-525; Riechmann *et al.* (1988) *Nature* 332:323-327; Verhoeven *et al.* (1988) *Science* 239:1534-1536), by substituting rodent or mutant rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. See also U.S. Patent Nos. 5,225,539; 5,585,089; 5,693,761; 5,693,762; 5,859,205; herein incorporated by reference. In some instances, residues within the framework regions of one or more variable regions of the human immunoglobulin are replaced by corresponding non-human residues (see, for example, U.S. Patent Nos. 5,585,089; 5,693,761; 5,693,762; and 6,180,370). Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance (e.g., to obtain desired affinity). In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For

further details see Jones *et al.* (1986) *Nature* 331:522-525; Riechmann *et al.* (1988) *Nature* 332:323-329; and Presta (1992) *Curr. Op. Struct. Biol.* 2:593-596; herein incorporated by reference. Accordingly, such "humanized" antibodies may include antibodies wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some framework residues are substituted by residues from analogous sites in rodent antibodies. See, for example, U.S. Patent Nos. 5,225,539; 5,585,089; 5,693,761; 5,693,762; 5,859,205. See also U.S. Patent No. 6,180,370, and International Publication No. WO 01/27160, where humanized antibodies and techniques for producing humanized antibodies having improved affinity for a predetermined antigen are disclosed.

Also encompassed by the term anti-CD40 antibodies are xenogeneic or modified anti-CD40 antibodies produced in a non-human mammalian host, more particularly a transgenic mouse, characterized by inactivated endogenous immunoglobulin (Ig) loci. In such transgenic animals, competent endogenous genes for the expression of light and heavy subunits of host immunoglobulins are rendered non-functional and substituted with the analogous human immunoglobulin loci. These transgenic animals produce human antibodies in the substantial absence of light or heavy host immunoglobulin subunits. See, for example, U.S. Patent Nos. 5,877,397 and 5,939,598, herein incorporated by reference.

Preferably, fully human antibodies to CD40 are obtained by immunizing transgenic mice. One such mouse is obtained using XenoMouse[®] technology (Abgenix; Fremont, California), and is disclosed in U.S. Patent Nos. 6,075,181, 6,091,001, and 6,114,598, all of which are incorporated herein by reference. To produce the antibodies disclosed herein, mice transgenic for the human Ig G₁ heavy chain locus and the human κ light chain locus were immunized with Sf 9 cells expressing human CD40. Mice can also be transgenic for other isotypes. Fully human antibodies useful in the methods of the present invention are characterized by binding properties similar to those exhibited by the CHIR-5.9 and CHIR-12.12 monoclonal antibodies disclosed herein.

Fragments of the anti-CD40 antibodies are suitable for use in the methods of the invention so long as they retain the desired affinity of the full-length antibody. Thus, a

fragment of an anti-CD40 antibody will retain the ability to bind to the CD40 B cell surface antigen. Such fragments are characterized by properties similar to the corresponding full-length antagonist anti-CD40 antibody, that is, the fragments will specifically bind a human CD40 antigen expressed on the surface of a human cell, and are free of significant agonist activity but exhibit antagonist activity when bound to a CD40 antigen on a human CD40-expressing cell. Such fragments are referred to herein as "antigen-binding" fragments.

Suitable antigen-binding fragments of an antibody comprise a portion of a full-length antibody, generally the antigen-binding or variable region thereof. Examples of antibody fragments include, but are not limited to, Fab, F(ab')₂, and Fv fragments and single-chain antibody molecules. By "Fab" is intended a monovalent antigen-binding fragment of an immunoglobulin that is composed of the light chain and part of the heavy chain. By F(ab')₂ is intended a bivalent antigen-binding fragment of an immunoglobulin that contains both light chains and part of both heavy chains. By "single-chain Fv" or "sFv" antibody fragments is intended fragments comprising the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain. See, for example, U.S. Patent Nos. 4,946,778, 5,260,203, 5,455,030, and 5,856,456, herein incorporated by reference. Generally, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains that enables the sFv to form the desired structure for antigen binding. For a review of sFv see Pluckthun (1994) in *The Pharmacology of Monoclonal Antibodies*, Vol. 113, ed. Rosenberg and Moore (Springer-Verlag, New York), pp. 269-315. Antigen-binding fragments of the antagonist anti-CD40 antibodies disclosed herein can also be conjugated to a cytotoxin to effect killing of the target cancer cells, as described herein below.

Antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in, for example, McCafferty *et al.* (1990) *Nature* 348:552-554 (1990) and U.S. Patent No. 5,514,548. Clackson *et al.* (1991) *Nature* 352:624-628 and Marks *et al.* (1991) *J. Mol. Biol.* 222:581-597 describe the isolation of murine and human antibodies, respectively, using phage libraries.

Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks *et al.* (1992) *Bio/Technology* 10:779-783), as well

as combinatorial infection and *in vivo* recombination as a strategy for constructing very large phage libraries (Waterhouse *et al.* (1993) *Nucleic Acids Res.* 21:2265-2266). Thus, these techniques are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

5 Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived *via* proteolytic digestion of intact antibodies (see, e.g., Morimoto *et al.* (1992) *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan *et al.* (1985) *Science* 229:81). However, these fragments can now be produced directly by recombinant host cells. For example, the
10 antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')₂ fragments (Carter *et al.* (1992) *Bio/Technology* 10:163-167). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments
15 will be apparent to the skilled practitioner.

Antagonist anti-CD40 antibodies useful in the methods of the present invention include the CHIR-5.9 and CHIR-12.12 monoclonal antibodies disclosed herein as well as antibodies differing from this antibody but retaining the CDRs; and antibodies with one or more amino acid addition(s), deletion(s), or substitution(s), wherein the antagonist
20 activity is measured by inhibition of B-cell proliferation and/or differentiation. The invention also encompasses de-immunized antagonist anti-CD40 antibodies, which can be produced as described in, for example, International Publication Nos. WO 98/52976 and WO 0034317; herein incorporated by reference. In this manner, residues within the antagonist anti-CD40 antibodies of the invention are modified so as to render the
25 antibodies non- or less immunogenic to humans while retaining their antagonist activity toward human CD40-expressing cells, wherein such activity is measured by assays noted elsewhere herein. Also included within the scope of the claims are fusion proteins comprising an antagonist anti-CD40 antibody of the invention, or a fragment thereof, which fusion proteins can be synthesized or expressed from corresponding
30 polynucleotide vectors, as is known in the art. Such fusion proteins are described with reference to conjugation of antibodies as noted below.

The antibodies of the present invention can have sequence variations produced using methods described in, for example, Patent Publication Nos. EP 0 983 303 A1, WO 00/34317, and WO 98/52976, incorporated herein by reference. For example, it has been shown that sequences within the CDR can cause an antibody to bind to MHC Class II and trigger an unwanted helper T-cell response. A conservative substitution can allow the antibody to retain binding activity yet lose its ability to trigger an unwanted T-cell response. Any such conservative or non-conservative substitutions can be made using art-recognized methods, such as those noted elsewhere herein, and the resulting antibodies will fall within the scope of the invention. The variant antibodies can be routinely tested for antagonist activity, affinity, and specificity using methods described herein.

An antibody produced by any of the methods described above, or any other method not disclosed herein, will fall within the scope of the invention if it possesses at least one of the following biological activities: inhibition of immunoglobulin secretion by normal human peripheral B cells stimulated by T cells; inhibition of survival and/or proliferation of normal human peripheral B cells stimulated by Jurkat T cells; inhibition of survival and/or proliferation of normal human peripheral B cells stimulated by CD40L-expressing cells or soluble CD40 ligand (sCD40L); inhibition of "survival" anti-apoptotic intracellular signals in any cell stimulated by sCD40L or solid-phase CD40L; inhibition of CD40 signal transduction in any cell upon ligation with sCD40L or solid-phase CD40L; and inhibition of proliferation of human malignant B cells as noted below. These assays can be performed as described in the Examples herein. See also the assays described in Schultze *et al.* (1998) *Proc. Natl. Acad. Sci. USA* 92:8200-8204; Denton *et al.* (1998) *Pediatr. Transplant.* 2:6-15; Evans *et al.* (2000) *J. Immunol.* 164:688-697; Noelle (1998) *Agents Actions Suppl.* 49:17-22; Lederman *et al.* (1996) *Curr. Opin. Hematol.* 3:77-86; Coligan *et al.* (1991) *Current Protocols in Immunology* 13:12; Kwekkeboom *et al.* (1993) *Immunology* 79:439-444; and U.S. Patent Nos. 5,674,492 and 5,847,082; herein incorporated by reference.

A representative assay to detect antagonist anti-CD40 antibodies specific to the CD40-antigen epitopes identified herein is a "competitive binding assay." Competitive binding assays are serological assays in which unknowns are detected and quantitated by

their ability to inhibit the binding of a labeled known ligand to its specific antibody. This is also referred to as a competitive inhibition assay. In a representative competitive binding assay, labeled CD40 polypeptide is precipitated by candidate antibodies in a sample, for example, in combination with monoclonal antibodies raised against one or more epitopes of the monoclonal antibodies of the invention. Anti-CD40 antibodies that specifically react with an epitope of interest can be identified by screening a series of antibodies prepared against a CD40 protein or fragment of the protein comprising the particular epitope of the CD40 protein of interest. For example, for human CD40, epitopes of interest include epitopes comprising linear and/or nonlinear amino acid residues of the short isoform of human CD40 (see GenBank Accession No. NP_690593) set forth in Figure 12B (SEQ ID NO:10), encoded by the sequence set forth in Figure 12A (SEQ ID NO:9; see also GenBank Accession No. NM_152854), or of the long isoform of human CD40 (see GenBank Accession Nos. CAA43045 and NP_001241) set forth in Figure 12D (SEQ ID NO:12), encoded by the sequence set forth in Figure 12C (SEQ ID NO:11; see GenBank Accession Nos. X60592 and NM_001250). Alternatively, competitive binding assays with previously identified suitable antagonist anti-CD40 antibodies could be used to select monoclonal antibodies comparable to the previously identified antibodies.

Antibodies employed in such immunoassays may be labeled or unlabeled. Unlabeled antibodies may be employed in agglutination; labeled antibodies may be employed in a wide variety of assays, employing a wide variety of labels. Detection of the formation of an antibody-antigen complex between an anti-CD40 antibody and an epitope of interest can be facilitated by attaching a detectable substance to the antibody. Suitable detection means include the use of labels such as radionuclides, enzymes, coenzymes, fluorescers, chemiluminescers, chromogens, enzyme substrates or co-factors, enzyme inhibitors, prosthetic group complexes, free radicals, particles, dyes, and the like. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a

luminescent material is luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S , or ^3H . Such labeled reagents may be used in a variety of well-known assays, such as radioimmunoassays, enzyme immunoassays, e.g., ELISA, fluorescent immunoassays, and the like. See for example, U.S. Patent Nos. 3,766,162; 3,791,932; 3,817,837; and 4,233,402.

Any of the previously described antagonist anti-CD40 antibodies or antibody fragments thereof may be conjugated prior to use in the methods of the present invention. Methods for producing conjugated antibodies are known in the art. Thus, the anti-CD40 antibody may be labeled using an indirect labeling or indirect labeling approach. By "indirect labeling" or "indirect labeling approach" is intended that a chelating agent is covalently attached to an antibody and at least one radionuclide is inserted into the chelating agent. See, for example, the chelating agents and radionuclides described in Srivagtava and Mease (1991) *Nucl. Med. Bio.* 18:589-603, herein incorporated by reference. Suitable labels include fluorophores, chromophores, radioactive atoms (particularly ^{32}P and ^{125}I), electron-dense reagents, enzymes, and ligands having specific binding partners. Enzymes are typically detected by their activity. For example, horseradish peroxidase is usually detected by its ability to convert 3,3',5,5'-tetramethylbenzidine (TMB) to a blue pigment, quantifiable with a spectrophotometer. "Specific binding partner" refers to a protein capable of binding a ligand molecule with high specificity, as for example in the case of an antigen and a monoclonal antibody specific therefore. Other specific binding partners include biotin and avidin or streptavidin, Ig G and protein A, and the numerous receptor-ligand couples known in the art. It should be understood that the above description is not meant to categorize the various labels into distinct classes, as the same label may serve in several different modes. For example, ^{125}I may serve as a radioactive label or as an electron-dense reagent. HRP may serve as enzyme or as antigen for a mAb. Further, one may combine various labels for desired effect. For example, mAbs and avidin also require labels in the practice of this invention: thus, one might label a mAb with biotin, and detect its presence with avidin labeled with ^{125}I , or with an anti-biotin mAb labeled with HRP. Other

permutations and possibilities will be readily apparent to those of ordinary skill in the art, and are considered as equivalents within the scope of the instant invention.

Alternatively, the anti-CD40 antibody may be labeled using "direct labeling" or a "direct labeling approach," where a radionuclide is covalently attached directly to an antibody (typically via an amino acid residue). Preferred radionuclides are provided in
5 Srivagtava and Mease (1991) *supra*. The indirect labeling approach is particularly preferred. See also, for example, International Publication Nos. WO 00/52031 and WO 00/52473, where a linker is used to attach a radioactive label to antibodies; and the labeled forms of anti-CD40 antibodies described in U.S. Patent No. 6,015,542; herein
10 incorporated by reference.

Further, an antibody (or fragment thereof) may be conjugated to a therapeutic moiety such as a cytotoxin, a therapeutic agent, or a radioactive metal ion or radioisotope. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin,
15 etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine,
20 cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly
25 actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine). Radioisotopes include, but are not limited to, I-131, I-123, I-125, Y-90, Re-188, Re-186, At-211, Cu-67, Bi-212, Bi-213, Pd-109, Tc-99, In-111, and the like. The conjugates of the invention can be used for modifying a given biological response; the drug moiety is not to be construed as limited to classical
30 chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for

example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, interferon-alpha, interferon-beta, nerve growth factor, platelet derived growth factor, tissue plasminogen activator; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"),
 5 interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Techniques for conjugating such therapeutic moiety to antibodies are well known. See, for example, Arnon *et al.* (1985) "Monoclonal Antibodies for Immunotargeting of
 10 Drugs in Cancer Therapy," in *Monoclonal Antibodies and Cancer Therapy*, ed. Reisfeld *et al.* (Alan R. Liss, Inc.), pp. 243-256; ed. Hellstrom *et al.* (1987) "Antibodies for Drug Delivery," in *Controlled Drug Delivery*, ed. Robinson *et al.* (2d ed; Marcel Dekker, Inc.), pp. 623-653; Thorpe (1985) "Antibody Carriers of Cytotoxic Agents in Cancer Therapy: A Review," in *Monoclonal Antibodies '84: Biological and Clinical Applications*, ed.
 15 Pinchera *et al.* pp. 475-506 (Editrice Kurtis, Milano, Italy, 1985); "Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody in Cancer Therapy," in *Monoclonal Antibodies for Cancer Detection and Therapy*, ed. Baldwin *et al.* (Academic Press, New York, 1985), pp. 303-316; and Thorpe *et al.* (1982) *Immunol. Rev.* 62:119-158.

20 Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described in U.S. Patent No. 4,676,980. In addition, linkers may be used between the labels and the antibodies of the invention (see U.S. Patent No. 4,831,175). Antibodies or, antigen-binding fragments thereof may be directly labeled with radioactive iodine, indium, yttrium, or other radioactive particle known in the art
 25 (U.S. Patent No. 5,595,721). Treatment may consist of a combination of treatment with conjugated and nonconjugated antibodies administered simultaneously or subsequently (WO 00/52031 and WO 00/52473).

Variants of Antagonist Anti-CD40 Antibodies

30 Suitable biologically active variants of the antagonist anti-CD40 antibodies can be used in the methods of the present invention. Such variants will retain the desired

binding properties of the parent antagonist anti-CD40 antibody. Methods for making antibody variants are generally available in the art.

For example, amino acid sequence variants of an antagonist anti-CD40 antibody, for example, the CHIR-5.9 or CHIR-12.12 monoclonal antibody described herein, can be prepared by mutations in the cloned DNA sequence encoding the antibody of interest. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York); Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods Enzymol.* 154:367-382; Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor, New York); U.S. Patent No. 4,873,192; and the references cited therein; herein incorporated by reference. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the polypeptide of interest may be found in the model of Dayhoff *et al.* (1978) in *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferred. Examples of conservative substitutions include, but are not limited to, Gly \leftrightarrow Ala, Val \leftrightarrow Ile \leftrightarrow Leu, Asp \leftrightarrow Glu, Lys \leftrightarrow Arg, Asn \leftrightarrow Gln, and Phe \leftrightarrow Trp \leftrightarrow Tyr.

In constructing variants of the antagonist anti-CD40 antibody polypeptide of interest, modifications are made such that variants continue to possess the desired activity, i.e., similar binding affinity and are capable of specifically binding to a human CD40 antigen expressed on the surface of a human cell, and being free of significant agonist activity but exhibiting antagonist activity when bound to a CD40 antigen on a human CD40-expressing cell. Obviously, any mutations made in the DNA encoding the variant polypeptide must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. See EP Patent Application Publication No. 75,444.

In addition, the constant region of an antagonist anti-CD40 antibody can be mutated to alter effector function in a number of ways. For example, see U.S. Patent No. 6,737,056B1 and U.S. Patent Application Publication No. 2004/0132101A1, which disclose Fc mutations that optimize antibody binding to Fc receptors.

Preferably, variants of a reference antagonist anti-CD40 antibody have amino acid sequences that have at least 70% or 75% sequence identity, preferably at least 80% or 85% sequence identity, more preferably at least 90%, 91%, 92%, 93%, 94% or 95% sequence identity to the amino acid sequence for the reference antagonist anti-CD40 antibody molecule, for example, the CHIR-5.9 or CHIR-12.12 monoclonal antibody described herein, or to a shorter portion of the reference antibody molecule. More preferably, the molecules share at least 96%, 97%, 98% or 99% sequence identity. For purposes of the present invention, percent sequence identity is determined using the Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 2, BLOSUM matrix of 62. The Smith-Waterman homology search algorithm is taught in Smith and Waterman (1981) *Adv. Appl. Math.* 2:482-489. A variant may, for example, differ from the reference antagonist anti-CD40 antibody by as few as 1 to 15 amino acid residues, as few as 1 to 10 amino acid residues, such as 6-10, as few as 5, as few as 4, 3, 2, or even 1 amino acid residue.

With respect to optimal alignment of two amino acid sequences, the contiguous segment of the variant amino acid sequence may have additional amino acid residues or deleted amino acid residues with respect to the reference amino acid sequence. The contiguous segment used for comparison to the reference amino acid sequence will include at least 20 contiguous amino acid residues, and may be 30, 40, 50, or more amino acid residues. Corrections for sequence identity associated with conservative residue substitutions or gaps can be made (see Smith-Waterman homology search algorithm).

The precise chemical structure of a polypeptide capable of specifically binding CD40 and retaining antagonist activity, particularly when bound to CD40 antigen on malignant B cells, depends on a number of factors. As ionizable amino and carboxyl groups are present in the molecule, a particular polypeptide may be obtained as an acidic or basic salt, or in neutral form. All such preparations that retain their biological activity when placed in suitable environmental conditions are included in the definition of antagonist anti-CD40 antibodies as used herein. Further, the primary amino acid sequence of the polypeptide may be augmented by derivatization using sugar moieties (glycosylation) or by other supplementary molecules such as lipids, phosphate, acetyl groups and the like. It may also be augmented by conjugation with saccharides. Certain

aspects of such augmentation are accomplished through post-translational processing systems of the producing host; other such modifications may be introduced *in vitro*. In any event, such modifications are included in the definition of an anti-CD40 antibody used herein so long as the antagonist properties of the anti-CD40 antibody are not destroyed. It is expected that such modifications may quantitatively or qualitatively affect the activity, either by enhancing or diminishing the activity of the polypeptide, in the various assays. Further, individual amino acid residues in the chain may be modified by oxidation, reduction, or other derivatization, and the polypeptide may be cleaved to obtain fragments that retain activity. Such alterations that do not destroy antagonist activity do not remove the polypeptide sequence from the definition of anti-CD40 antibodies of interest as used herein.

The art provides substantial guidance regarding the preparation and use of polypeptide variants. In preparing the anti-CD40 antibody variants, one of skill in the art can readily determine which modifications to the native protein nucleotide or amino acid sequence will result in a variant that is suitable for use as a therapeutically active component of a pharmaceutical composition used in the methods of the present invention.

Methods of Therapy Using the Antagonist Anti-CD40 Antibodies of the Invention

Methods of the invention are directed to the use of antagonist anti-CD40 antibodies to treat patients having a disease mediated by stimulation of CD40 signaling on CD40-expressing cells. By "CD40-expressing cell" is intended normal and malignant B cells expressing CD40 antigen. Methods for detecting CD40 expression in cells are well known in the art and include, but are not limited to, PCR techniques, immunohistochemistry, flow cytometry, Western blot, ELISA, and the like. By "malignant" B cell is intended any neoplastic B cell, including but not limited to B cells derived from lymphomas including low-, intermediate-, and high-grade B cell lymphomas, immunoblastic lymphomas, non-Hodgkin's lymphomas, Hodgkin's disease, Epstein-Barr Virus (EBV) induced lymphomas, and AIDS-related lymphomas, as well as B cell acute lymphoblastic leukemias, myelomas, chronic lymphocytic leukemias, acute myeloblastic leukemias, and the like.

"Treatment" is herein defined as the application or administration of an antagonist anti-CD40 antibody or antigen-binding fragment thereof to a patient, or application or administration of an antagonist anti-CD40 antibody or fragment thereof to an isolated tissue or cell line from a patient, where the patient has a disease, a symptom of a disease, or a predisposition toward a disease, where the purpose is to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptoms of the disease, or the predisposition toward the disease. By "treatment" is also intended the application or administration of a pharmaceutical composition comprising the antagonist anti-CD40 antibodies or fragments thereof to a patient, or application or administration of a pharmaceutical composition comprising the anti-CD40 antibodies or fragments thereof to an isolated tissue or cell line from a patient, who has a disease, a symptom of a disease, or a predisposition toward a disease, where the purpose is to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptoms of the disease, or the predisposition toward the disease.

By "anti-tumor activity" is intended a reduction in the rate of malignant CD40-expressing cell proliferation or accumulation, and hence a decline in growth rate of an existing tumor or in a tumor that arises during therapy, and/or destruction of existing neoplastic (tumor) cells or newly formed neoplastic cells, and hence a decrease in the overall size of a tumor during therapy. Therapy with at least one anti-CD40 antibody (or antigen-binding fragment thereof) causes a physiological response that is beneficial with respect to treatment of disease states associated with stimulation of CD40 signaling on CD40-expressing cells in a human.

The methods of the invention find use in the treatment of non-Hodgkin's lymphomas related to abnormal, uncontrollable B cell proliferation or accumulation. For purposes of the present invention, such lymphomas will be referred to according to the *Working Formulation* classification scheme, that is those B cell lymphomas categorized as low grade, intermediate grade, and high grade (see "The Non-Hodgkin's Lymphoma Pathologic Classification Project," *Cancer* 49(1982):2112-2135). Thus, low-grade B cell lymphomas include small lymphocytic, follicular small-cleaved cell, and follicular mixed small-cleaved and large cell lymphomas; intermediate-grade lymphomas include follicular large cell, diffuse small cleaved cell, diffuse mixed small and large cell, and

diffuse large cell lymphomas; and high-grade lymphomas include large cell immunoblastic, lymphoblastic, and small non-cleaved cell lymphomas of the Burkitt's and non-Burkitt's type.

It is recognized that the methods of the invention are useful in the therapeutic treatment of B cell lymphomas that are classified according to the Revised European and American Lymphoma Classification (REAL) system. Such B cell lymphomas include, but are not limited to, lymphomas classified as precursor B cell neoplasms, such as B lymphoblastic leukemia/lymphoma; peripheral B cell neoplasms, including B cell chronic lymphocytic leukemia/small lymphocytic lymphoma, lymphoplasmacytoid lymphoma/immunocytoma, mantle cell lymphoma (MCL), follicle center lymphoma (follicular) (including diffuse small cell, diffuse mixed small and large cell, and diffuse large cell lymphomas), marginal zone B cell lymphoma (including extranodal, nodal, and splenic types), hairy cell leukemia, plasmacytoma/ myeloma, diffuse large cell B cell lymphoma of the subtype primary mediastinal (thymic), Burkitt's lymphoma, and Burkitt's like high grade B cell lymphoma; acute leukemias; acute lymphocytic leukemias; myeloblastic leukemias; acute myelocytic leukemias; promyelocytic leukemia; myelomonocytic leukemia; monocytic leukemia; erythroleukemia; granulocytic leukemia (chronic myelocytic leukemia); chronic lymphocytic leukemia; polycythemia vera; multiple myeloma; Waldenstrom's macroglobulinemia; heavy chain disease; and unclassifiable low-grade or high-grade B cell lymphomas.

It is recognized that the methods of the invention may be useful in preventing further tumor outgrowths arising during therapy. The methods of the invention are particularly useful in the treatment of subjects having low-grade B cell lymphomas, particularly those subjects having relapses following standard chemotherapy. Low-grade B cell lymphomas are more indolent than the intermediate- and high-grade B cell lymphomas and are characterized by a relapsing/remitting course. Thus, treatment of these lymphomas is improved using the methods of the invention, as relapse episodes are reduced in number and severity.

The antagonist anti-CD40 antibodies described herein may also find use in the treatment of inflammatory diseases and deficiencies or disorders of the immune system including, but not limited to, systemic lupus erythematosus, psoriasis, scleroderma,

CREST syndrome, inflammatory myositis, Sjogren's syndrome, mixed connective tissue disease, rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease, acute respiratory distress syndrome, pulmonary inflammation, idiopathic pulmonary fibrosis, osteoporosis, delayed type hypersensitivity, asthma, primary biliary cirrhosis, and
5 idiopathic thrombocytopenic purpura.

In accordance with the methods of the present invention, at least one antagonist anti-CD40 antibody (or antigen-binding fragment thereof) as defined elsewhere herein is used to promote a positive therapeutic response with respect to a malignant human B cell. By "positive therapeutic response" with respect to cancer treatment is intended an
10 improvement in the disease in association with the anti-tumor activity of these antibodies or fragments thereof, and/or an improvement in the symptoms associated with the disease. That is, an anti-proliferative effect, the prevention of further tumor outgrowths, a reduction in tumor size, a reduction in the number of cancer cells, and/or a decrease in one or more symptoms mediated by stimulation of CD40-expressing cells can be
15 observed. Thus, for example, an improvement in the disease may be characterized as a complete response. By "complete response" is intended an absence of clinically detectable disease with normalization of any previously abnormal radiographic studies, bone marrow, and cerebrospinal fluid (CSF). Such a response must persist for at least one month following treatment according to the methods of the invention. Alternatively,
20 an improvement in the disease may be categorized as being a partial response. By "partial response" is intended at least about a 50% decrease in all measurable tumor burden (i.e., the number of tumor cells present in the subject) in the absence of new lesions and persisting for at least one month. Such a response is applicable to measurable tumors only.

Tumor response can be assessed for changes in tumor morphology (i.e., overall tumor burden, tumor size, and the like) using screening techniques such as magnetic resonance imaging (MRI) scan, x-radiographic imaging, computed tomographic (CT) scan, bioluminescent imaging, for example, luciferase imaging, bone scan imaging, and tumor biopsy sampling including bone marrow aspiration (BMA). In addition to these
25 positive therapeutic responses, the subject undergoing therapy with the antagonist anti-CD40 antibody or antigen-binding fragment thereof may experience the beneficial effect
30

of an improvement in the symptoms associated with the disease. Thus for B cell tumors, the subject may experience a decrease in the so-called B symptoms, i.e., night sweats, fever, weight loss, and/or urticaria.

By "therapeutically effective dose or amount" or "effective amount" is intended
5 an amount of antagonist anti-CD40 antibody or antigen-binding fragment thereof that, when administered brings about a positive therapeutic response with respect to treatment of a patient with a disease comprising stimulation of CD40-expressing cells. In some
10 embodiments of the invention, a therapeutically effective dose of the anti-CD40 antibody or fragment thereof is in the range from about 0.01 mg/kg to about 40 mg/kg, from about 0.01 mg/kg to about 30 mg/kg, from about 0.1 mg/kg to about 30 mg/kg, from about 1 mg/kg to about 30 mg/kg, from about 3 mg/kg to about 30 mg/kg, from about 3 mg/kg to about 25 mg/kg, from about 3 mg/kg to about 20 mg/kg, from about 5 mg/kg to about 15 mg/kg, or from about 7 mg/kg to about 12 mg/kg. It is recognized that the method of
15 treatment may comprise a single administration of a therapeutically effective dose or multiple administrations of a therapeutically effective dose of the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

A further embodiment of the invention is the use of antagonist anti-CD40 antibodies for diagnostic monitoring of protein levels in tissue as part of a clinical testing procedure, e.g., to determine the efficacy of a given treatment regimen. Detection can be
20 facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include
25 streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S , or ^3H .

30 The anti-CD40 antibodies described herein can further be used to provide reagents, e.g., labeled antibodies that can be used, for example, to identify cells

expressing CD40. This can be very useful in determining the cell type of an unknown sample. Panels of monoclonal antibodies can be used to identify tissue by species and/or by organ type. In a similar fashion, these anti-CD40 antibodies can be used to screen tissue culture cells for contamination (i.e., screen for the presence of a mixture of CD40-expressing and non-CD40 expressing cells in a culture).

The antagonist anti-CD40 antibodies can be used in combination with known chemotherapeutics and cytokines for the treatment of disease states comprising stimulated CD40-expressing cells. For example, the anti-CD40 antibodies of the invention can be used in combination with cytokines such as interleukin-2. In another embodiment, the anti-CD40 antibodies of the invention can be used in combination with rituximab (IDEC-C2B8; Rituxan®; IDEC Pharmaceuticals Corp., San Diego, California).

In this manner, the antagonist anti-CD40 antibodies described herein, or antigen-binding fragments thereof, are administered in combination with at least one other cancer therapy, including, but not limited to, surgery or surgical procedures (e.g. splenectomy, hepatectomy, lymphadenectomy, leukopheresis, bone marrow transplantation, and the like); radiation therapy; chemotherapy, optionally in combination with autologous bone marrow transplant, where suitable chemotherapeutic agents include, but are not limited to, fludarabine or fludarabine phosphate, chlorambucil, vincristine, pentostatin, 2-chlorodeoxyadenosine (cladribine), cyclophosphamide, doxorubicin, prednisone, and combinations thereof, for example, anthracycline-containing regimens such as CAP (cyclophosphamide, doxorubicin plus prednisone), CHOP (cyclophosphamide, vincristine, prednisone plus doxorubicin), VAD (vincristine, doxorubicin, plus dexamethasone), MP (melphalan plus prednisone), and other cytotoxic and/or therapeutic agents used in chemotherapy such as mitoxantrone, daunorubicin, idarubicin, asparaginase, and antimetabolites, including, but not limited to, cytarabine, methotrexate, 5-fluorouracil decarbazine, 6-thioguanine, 6-mercaptopurine, and nelarabine; other anti-cancer monoclonal antibody therapy (for example, alemtuzumab (Campath®) or other anti-CD52 antibody targeting the CD52 cell-surface glycoprotein on malignant B cells; rituximab (Rituxan®), the fully human antibody HuMax-CD20, R-1594, IMMU-106, TRU-015, AME-133, tositumomab/I-131 tositumomab (Bexxar®), ibritumomab tiuxetan (Zevalin®), or any other therapeutic anti-CD20 antibody targeting the CD20 antigen on

malignant B cells; anti-CD19 antibody (for example, MT103, a bispecific antibody); anti-CD22 antibody (for example, the humanized monoclonal antibody epratuzumab); bevacizumab (Avastin®) or other anti-cancer antibody targeting human vascular endothelial growth factor; anti-CD22 antibody targeting the CD22 antigen on malignant B cells (for example, the monoclonal antibody BL-22, an alphaCD22 toxin); α -M-CSF antibody targeting macrophage colony stimulating factor; antibodies targeting the receptor activator of nuclear factor-kappaB (RANK) and its ligand (RANKL), which are overexpressed in multiple myeloma; anti-CD23 antibody targeting the CD23 antigen on malignant B cells (for example, IDEC-152); anti-CD80 antibody targeting the CD80 antigen (for example, IDEC-114); anti-CD38 antibody targeting the CD38 antigen on malignant B cells; antibodies targeting major histocompatibility complex class II receptors (anti-MHC antibodies) expressed on malignant B cells; other anti-CD40 antibodies (for example, SGN-40) targeting the CD40 antigen on malignant B cells; and antibodies targeting tumor necrosis factor-related apoptosis-inducing ligand receptor 1 (TRAIL-R1) (for example, the agonistic human monoclonal antibody HGS-ETR1) and TRAIL-R2 expressed on a number of solid tumors and tumors of hematopoietic origin); small molecule-based cancer therapy, including, but not limited to, microtubule and/or topoisomerase inhibitors (for example, the mitotic inhibitor dolastatin and dolastatin analogues; the tubulin-binding agent T900607; XL119; and the topoisomerase inhibitor aminocamptothecin), SDX-105 (bendamustine hydrochloride), ixabepilone (an epothilone analog, also referred to as BMS-247550), protein kinase C inhibitors, for example, midostaurin ((PKC-412, CGP 41251, N-benzoylstaurosporine), pixantrone, eloxatin (an antineoplastic agent), ganite (gallium nitrate), Thalomid® (thalidomide), immunomodulatory derivatives of thalidomide (for example, revlimid (formerly revimid)), Affinitak™ (antisense inhibitor of protein kinase C-alpha), SDX-101 (R-etodolac, inducing apoptosis of malignant lymphocytes), second-generation purine nucleoside analogs such as clofarabine, inhibitors of production of the protein Bcl-2 by cancer cells (for example, the antisense agents oblimersen and Genasense®), proteasome inhibitors (for example, Velcade™ (bortezomib)), small molecule kinase inhibitors (for example, CHIR-258), small molecule VEGF inhibitors (for example, ZD-6474), small molecule inhibitors of heat shock protein (HSP) 90 (for example, 17-AAG), small

molecule inhibitors of histone deacetylases (for example, hybrid/polar cytodifferentiation HPC) agents such as suberanilohydroxamic acid (SAHA), and FR-901228) and apoptotic agents such as Trisenox[®] (arsenic trioxide) and Xcytrin[®] (motexafin gadolinium); vaccine /immunotherapy-based cancer therapies, including, but not limited to, vaccine approaches (for example, Id-KLH, oncopophage, vitaethine), personalized immunotherapy or active idiotypic immunotherapy (for example, MyVax[®] Personalized Immunotherapy, formally designated GTOP-99), Promune[®] (CpG 7909, a synthetic agonist for toll-like receptor 9 (TLR9)), interferon-alpha therapy, interleukin-2 (IL-2) therapy, IL-12 therapy, IL-15 therapy, and IL-21 therapy; steroid therapy; or other cancer therapy; where the additional cancer therapy is administered prior to, during, or subsequent to the antagonist anti-CD40 antibody therapy. Thus, where the combined therapies comprise administration of an antagonist anti-CD40 antibody or antigen-binding fragment thereof in combination with administration of another therapeutic agent, as with chemotherapy, radiation therapy, other anti-cancer antibody therapy, small molecule-based cancer therapy, or vaccine/immunotherapy-based cancer therapy, the methods of the invention encompass coadministration, using separate formulations or a single pharmaceutical formulation, or and consecutive administration in either order. Where the methods of the present invention comprise combined therapeutic regimens, these therapies can be given simultaneously, i.e., the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered concurrently or within the same time frame as the other cancer therapy (i.e., the therapies are going on concurrently, but the antagonist anti-CD40 antibody or antigen-binding fragment thereof is not administered precisely at the same time as the other cancer therapy). Alternatively, the antagonist anti-CD40 antibody of the present invention or antigen-binding fragment thereof may also be administered prior to or subsequent to the other cancer therapy. Sequential administration of the different cancer therapies may be performed regardless of whether the treated subject responds to the first course of therapy to decrease the possibility of remission or relapse. Where the combined therapies comprise administration of the antagonist anti-CD40 antibody or antigen-binding fragment thereof in combination with administration of a cytotoxic agent, preferably the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered prior to administering the cytotoxic agent.

In some embodiments of the invention, the antagonist anti-CD40 antibodies described herein, or antigen-binding fragments thereof, are administered in combination with chemotherapy, and optionally in combination with autologous bone marrow transplantation, wherein the antibody and the chemotherapeutic agent(s) may be administered sequentially, in either order, or simultaneously (i.e., concurrently or within the same time frame). Examples of suitable chemotherapeutic agents include, but are not limited to, fludarabine or fludarabine phosphate, chlorambucil, vincristine, pentostatin, 2-chlorodeoxyadenosine (cladribine), cyclophosphamide, doxorubicin, prednisone, and combinations thereof, for example, anthracycline-containing regimens such as CAP (cyclophosphamide, doxorubicin plus prednisone), CHOP (cyclophosphamide, vincristine, prednisone plus doxorubicin), VAD (vincristine, doxorubicin, plus dexamethasone), MP (melphalan plus prednisone), and other cytotoxic and/or therapeutic agents used in chemotherapy such as mitoxantrone, daunorubicin, idarubicin, asparaginase, and antimetabolites, including, but not limited to, cytarabine, methotrexate, 5-fluorouracil decarbazine, 6-thioguanine, 6-mercaptopurine, and nelarabine. In some embodiments, the antagonist anti-CD40 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or an antigen-binding fragment thereof is administered prior to treatment with the chemotherapeutic agent. In alternative embodiments, the antagonist anti-CD40 antibody is administered after treatment with the chemotherapeutic agent. In yet other embodiments, the chemotherapeutic agent is administered simultaneously with the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

Thus, for example, in some embodiments, the antagonist anti-CD40 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof is administered in combination with fludarabine or fludarabine phosphate. In one such embodiment, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered prior to administration of fludarabine or fludarabine phosphate. In alternative embodiments, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered after treatment with fludarabine or fludarabine phosphate. In yet other embodiments, the fludarabine or fludarabine phosphate is administered simultaneously with the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

In other embodiments of the invention, chlorambucil, an alkylating drug, is administered in combination with an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or an antigen-binding fragment thereof. In one such embodiment, the antagonist anti-CD40 antibody or
5 antigen-binding fragment thereof is administered prior to administration of chlorambucil. In alternative embodiments, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered after treatment with chlorambucil. In yet other embodiments, the chlorambucil is administered simultaneously with the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

10 In yet other embodiments, anthracycline-containing regimens such as CAP (cyclophosphamide, doxorubicin plus prednisone) and CHOP (cyclophosphamide, vincristine, prednisone plus doxorubicin) may be combined with administration of an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof. In one such embodiment,
15 the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered prior to administration of anthracycline-containing regimens. In other embodiments, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered after treatment with anthracycline-containing regimens. In yet other embodiments, the anthracycline-containing regimen is administered simultaneously with the antagonist
20 anti-CD40 antibody or antigen-binding fragment thereof.

In alternative embodiments, an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or an antigen-binding fragment thereof, is administered in combination with alemtuzumab (Campath[®]; distributed by Berlex Laboratories, Richmond, California). Alemtuzumab is a
25 recombinant humanized monoclonal antibody (Campath-1H) that targets the CD52 antigen expressed on malignant B cells. In one such embodiment, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered prior to administration of alemtuzumab. In other embodiments, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered after treatment with
30 alemtuzumab. In yet other embodiments, the alemtuzumab is administered

simultaneously with the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

In alternative embodiments, an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding
5 fragment thereof, is administered in combination with a therapeutic anti-CD20 antibody targeting the CD20 antigen on malignant B cells, for example, rituximab (Rituxan[®]), the fully human antibody HuMax-CD20, R-1594, IMMU-106, TRU-015, AME-133, tositumomab/I-131 tositumomab (Bexxar[®]), or ibritumomab tiuxetan (Zevalin[®]). In one such embodiment, the antagonist anti-CD40 antibody or antigen-binding fragment thereof
10 is administered prior to administration of the anti-CD20 antibody. In other embodiments, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered after treatment with the anti-CD20 antibody. In yet other embodiments, the anti-CD20 antibody is administered simultaneously with the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

In alternative embodiments, an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding
15 fragment thereof, is administered in combination with a small molecule-based cancer therapy, including, but not limited to, microtubule and/or topoisomerase inhibitors (for example, the mitotic inhibitor dolastatin and dolastatin analogues; the tubulin-binding
20 agent T900607; XL119; and the topoisomerase inhibitor aminocamptothecin), SDX-105 (bendamustine hydrochloride), ixabepilone (an epothilone analog, also referred to as BMS-247550), protein kinase C inhibitors, for example, midostaurin ((PKC-412, CGP 41251, N-benzoylstauroporine), pixantrone, eloxatin (an antineoplastic agent), ganite (gallium nitrate), Thalomid[®] (thalidomide), immunomodulatory derivatives of
25 thalidomide (for example, revlimid (formerly revimid)), Affinitak[™] (antisense inhibitor of protein kinase C- α), SDX-101 (R-etodolac, inducing apoptosis of malignant lymphocytes), second-generation purine nucleoside analogs such as clofarabine, inhibitors of production of the protein Bcl-2 by cancer cells (for example, the antisense agents oblimersen and Genasense[®]), proteasome inhibitors (for example, Velcade[™]
30 (bortezomib)), small molecule kinase inhibitors (for example, CHIR-258), small molecule VEGF inhibitors (for example, ZD-6474), small molecule inhibitors of heat

shock protein (HSP) 90 (for example, 17-AAG), small molecule inhibitors of histone deacetylases (for example, hybrid/polar cytodifferentiation HPC) agents such as suberanilohydroxamic acid (SAHA), and FR-901228) and apoptotic agents such as Trisenox[®] (arsenic trioxide) and Xcytrin[®] (motexafin gadolinium). In one such
5 embodiment, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered prior to administration of the small molecule-based cancer therapy. In other embodiments, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered after treatment with the small molecule-based cancer therapy. In yet other
10 embodiments, the small molecule-based cancer therapy is administered simultaneously with the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

In yet other embodiments, an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9 or an antigen-binding fragment thereof, can be used in combination with vaccine/immunotherapy-based cancer therapy, including, but not limited to, vaccine approaches (for example, Id-KLH,
15 oncopophage, vitaethine), personalized immunotherapy or active idiotype immunotherapy (for example, MyVax[®] Personalized Immunotherapy, formally designated GTOP-99), Promune[®] (CpG 7909, a synthetic agonist for toll-like receptor 9 (TLR9)), interferon-alpha therapy, interleukin-2 (IL-2) therapy, IL-12 therapy, IL-15 therapy, or IL-21
20 therapy; or steroid therapy. In one such embodiment, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered prior to administration of the vaccine/immunotherapy-based cancer therapy. In other embodiments, the antagonist anti-CD40 antibody or antigen-binding fragment thereof is administered after treatment with the vaccine/immunotherapy-based cancer therapy. In yet other embodiments, the
25 vaccine/immunotherapy-based cancer therapy is administered simultaneously with the antagonist anti-CD40 antibody or antigen-binding fragment thereof.

In one such embodiment, an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or an antigen-binding fragment thereof, can be used in combination with IL-2. IL-2, an agent known to expand the number of natural killer (NK) effector cells in treated patients, can be administered
30 prior to, or concomitantly with, the antagonist anti-CD40 antibody of the invention or antigen-binding fragment thereof. This expanded number of NK effector cells may lead

to enhanced ADCC activity of the administered antagonist anti-CD40 antibody or antigen-binding fragment thereof. In other embodiments, IL-21 serves as the immunotherapeutic agent to stimulate NK cell activity when administered in combination with an antagonist anti-CD40 antibody described herein, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or an antigen-binding fragment thereof.

Further, combination therapy with two or more therapeutic agents and an antagonist anti-CD40 antibody described herein can also be used for treatment of a treatment of disease states comprising stimulated CD40-expressing cells, for example, B cell-related cancers, and autoimmune and/or inflammatory disorders. Without being limiting, examples include triple combination therapy, where two chemotherapeutic agents are administered in combination with an antagonist anti-CD40 antibody described herein, and where a chemotherapeutic agent and another anti-cancer monoclonal antibody (for example, alemtuzumab, rituximab, or anti-CD23 antibody) are administered in combination with an antagonist anti-CD40 antibody described herein. Examples of such combinations include, but are not limited to, combinations of fludarabine, cyclophosphamide, and the antagonist anti-CD40 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9 or an antigen-binding fragment thereof; and combinations of fludarabine, an anti-CD20 antibody, for example, rituximab (Rituxan[®]; IDEC Pharmaceuticals Corp., San Diego, California), and the antagonist anti-CD40 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9 or an antigen-binding fragment thereof.

Pharmaceutical Formulations and Modes of Administration

The antagonist anti-CD40 antibodies of this invention are administered at a concentration that is therapeutically effective to prevent or treat CD40-expressing cell-mediated diseases such as SLE, PBC, ITP, multiple sclerosis, psoriasis, Crohn's disease, graft rejection, and B-cell lymphoma. To accomplish this goal, the antibodies may be formulated using a variety of acceptable excipients known in the art. Typically, the antibodies are administered by injection, either intravenously or intraperitoneally. Methods to accomplish this administration are known to those of ordinary skill in the art.

It may also be possible to obtain compositions which may be topically or orally administered, or which may be capable of transmission across mucous membranes.

Intravenous administration occurs preferably by infusion over a period of about 1 to about 10 hours, more preferably over about 1 to about 8 hours, even more preferably over about 2 to about 7 hours, still more preferably over about 4 to about 6 hours, depending upon the anti-CD40 antibody being administered. The initial infusion with the pharmaceutical composition may be given over a period of about 4 to about 6 hours with subsequent infusions delivered more quickly. Subsequent infusions may be administered over a period of about 1 to about 6 hours, including, for example, about 1 to about 4 hours, about 1 to about 3 hours, or about 1 to about 2 hours.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of possible routes of administration include parenteral, (e.g., intravenous (IV), intramuscular (IM), intradermal, subcutaneous (SC), or infusion), oral and pulmonary (e.g., inhalation), nasal, transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes, or multiple dose vials made of glass or plastic.

The anti-CD40 antibodies are typically provided by standard technique within a pharmaceutically acceptable buffer, for example, sterile saline, sterile buffered water, propylene glycol, combinations of the foregoing, etc. Methods for preparing parenterally administrable agents are described in *Remington's Pharmaceutical Sciences* (18th ed.; Mack Publishing Company, Eaton, Pennsylvania, 1990), herein incorporated by reference. See also, for example, WO 98/56418, which describes stabilized antibody pharmaceutical formulations suitable for use in the methods of the present invention.

The amount of at least one anti-CD40 antibody or fragment thereof to be administered is readily determined by one of ordinary skill in the art without undue experimentation. Factors influencing the mode of administration and the respective amount of at least one antagonist anti-CD40 antibody (or fragment thereof) include, but
5 are not limited to, the the severity of the disease, the history of the disease, and the age, height, weight, health, and physical condition of the individual undergoing therapy. Similarly, the amount of antagonist anti-CD40 antibody or fragment thereof to be administered will be dependent upon the mode of administration and whether the subject will undergo a single dose or multiple doses of this anti-tumor agent. Generally, a higher
10 dosage of anti-CD40 antibody or fragment thereof is preferred with increasing weight of the subject undergoing therapy. The dose of anti-CD40 antibody or fragment thereof to be administered is in the range from about 0.003 mg/kg to about 50 mg/kg, preferably in the range of 0.01 mg/kg to about 40 mg/kg. Thus, for example, the dose can be 0.01 mg/kg, 0.03 mg/kg, 0.1 mg/kg, 0.3 mg/kg, 0.5 mg/kg, 1 mg/kg, 1.5 mg/kg, 2 mg/kg, 2.5
15 mg/kg, 3 mg/kg, 5 mg/kg, 7 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 30 mg/kg, 35 mg/kg, 40 mg/kg, 45 mg/kg, or 50 mg/kg.

In another embodiment of the invention, the method comprises administration of multiple doses of antagonist anti-CD40 antibody or fragment thereof. The method may comprise administration of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, or more
20 therapeutically effective doses of a pharmaceutical composition comprising an antagonist anti-CD40 antibody or fragment thereof. The frequency and duration of administration of multiple doses of the pharmaceutical compositions comprising anti-CD40 antibody or fragment thereof can be readily determined by one of skill in the art without undue experimentation. Moreover, treatment of a subject with a therapeutically effective
25 amount of an antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with antagonist anti-CD40 antibody or antigen-binding fragment thereof in the range of between about 0.1 to 20 mg/kg body weight, once per week for between about 1 to 10 weeks, preferably between about 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more
30 preferably for about 4, 5, or 6 weeks. Treatment may occur annually to prevent relapse or upon indication of relapse. It will also be appreciated that the effective dosage of

antibody or antigen-binding fragment thereof used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result and become apparent from the results of diagnostic assays as described herein.

Thus, in one embodiment, the dosing regimen includes a first administration of a therapeutically effective dose of at least one anti-CD40 antibody or fragment thereof on days 1, 7, 14, and 21 of a treatment period. In another embodiment, the dosing regimen includes a first administration of a therapeutically effective dose of at least one anti-CD40 antibody or fragment thereof on days 1, 2, 3, 4, 5, 6, and 7 of a week in a treatment period. Further embodiments include a dosing regimen having a first administration of a therapeutically effective dose of at least one anti-CD40 antibody or fragment thereof on days 1, 3, 5, and 7 of a week in a treatment period; a dosing regimen including a first administration of a therapeutically effective dose of at least one anti-CD40 antibody or fragment thereof on days 1 and 3 of a week in a treatment period; and a preferred dosing regimen including a first administration of a therapeutically effective dose of at least one anti-CD40 antibody or fragment thereof on day 1 of a week in a treatment period. The treatment period may comprise 1 week, 2 weeks, 3 weeks, a month, 3 months, 6 months, or a year. Treatment periods may be subsequent or separated from each other by a day, a week, 2 weeks, a month, 3 months, 6 months, or a year.

In some embodiments, the therapeutically effective doses of antagonist anti-CD40 antibody or antigen-binding fragment thereof ranges from about 0.003 mg/kg to about 50 mg/kg, from about 0.01 mg/kg to about 40 mg/kg, from about 0.01 mg/kg to about 30 mg/kg, from about 0.1 mg/kg to about 30 mg/kg, from about 0.5 mg/kg to about 30 mg/kg, from about 1 mg/kg to about 30 mg/kg, from about 3 mg/kg to about 30 mg/kg, from about 3 mg/kg to about 25 mg/kg, from about 3 mg/kg to about 20 mg/kg, from about 5 mg/kg to about 15 mg/kg, or from about 7 mg/kg to about 12 mg/kg. Thus, for example, the dose of any one antagonist anti-CD40 antibody or antigen-binding fragment thereof, for example the anti-CD40 monoclonal antibody CHIR-12.12 or CHIR-5.9 or antigen-binding fragment thereof, can be 0.003 mg/kg, 0.01 mg/kg, 0.03 mg/kg, 0.1 mg/kg, 0.3 mg/kg, 0.5 mg/kg, 1 mg/kg, 1.5 mg/kg, 2 mg/kg, 2.5 mg/kg, 3 mg/kg, 5 mg/kg, 7 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 30 mg/kg, 35 mg/kg, 40 mg/kg, 45 mg/kg, 50 mg/kg, or other such doses falling within the range of about 0.003

mg/kg to about 50 mg/kg. The same therapeutically effective dose of an antagonist anti-CD40 antibody or antigen-binding fragment thereof can be administered throughout each week of antibody dosing. Alternatively, different therapeutically effective doses of an antagonist anti-CD40 antibody or antigen-binding fragment thereof can be used over the course of a treatment period.

In other embodiments, the initial therapeutically effective dose of an antagonist anti-CD40 antibody or antigen-binding fragment thereof as defined elsewhere herein can be in the lower dosing range (i.e., about 0.003 mg/kg to about 20 mg/kg) with subsequent doses falling within the higher dosing range (i.e., from about 20 mg/kg to about 50 mg/kg).

In alternative embodiments, the initial therapeutically effective dose of an antagonist anti-CD40 antibody or antigen-binding fragment thereof as defined elsewhere herein can be in the upper dosing range (i.e., about 20 mg/kg to about 50 mg/kg) with subsequent doses falling within the lower dosing range (i.e., 0.003 mg/kg to about 20 mg/kg). Thus, in one embodiment, the initial therapeutically effective dose of the antagonist anti-CD40 antibody or antigen-binding fragment thereof is about 20 mg/kg to about 35 mg/kg, including about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, and about 35 mg/kg, and subsequent therapeutically effective doses of the antagonist anti-CD40 antibody or antigen binding fragment thereof are about 5 mg/kg to about 15 mg/kg, including about 5 mg/kg, 8 mg/kg, 10 mg/kg, 12 mg/kg, and about 15 mg/kg.

In some embodiments of the invention, antagonist anti-CD40 antibody therapy is initiated by administering a "loading dose" of the antibody or antigen-binding fragment thereof to the subject in need of antagonist anti-CD40 antibody therapy. By "loading dose" is intended an initial dose of the antagonist anti-CD40 antibody or antigen-binding fragment thereof that is administered to the subject, where the dose of the antibody or antigen-binding fragment thereof administered falls within the higher dosing range (i.e., from about 20 mg/kg to about 50 mg/kg). The "loading dose" can be administered as a single administration, for example, a single infusion where the antibody or antigen-binding fragment thereof is administered IV, or as multiple administrations, for example, multiple infusions where the antibody or antigen-binding fragment thereof is administered IV, so long as the complete "loading dose" is administered within about a

24-hour period. Following administration of the "loading dose," the subject is then administered one or more additional therapeutically effective doses of the antagonist anti-CD40 antibody or antigen-binding fragment thereof. Subsequent therapeutically effective doses can be administered, for example, according to a weekly dosing schedule, or once every two weeks, once every three weeks, or once every four weeks. In such embodiments, the subsequent therapeutically effective doses generally fall within the lower dosing range (i.e., 0.003 mg/kg to about 20 mg/kg).

Alternatively, in some embodiments, following the "loading dose," the subsequent therapeutically effective doses of the antagonist anti-CD40 antibody or antigen-binding fragment thereof are administered according to a "maintenance schedule," wherein the therapeutically effective dose of the antibody or antigen-binding fragment thereof is administered once a month, once every 6 weeks, once every two months, once every 10 weeks, once every three months, once every 14 weeks, once every four months, once every 18 weeks, once every five months, once every 22 weeks, once every six months, once every 7 months, once every 8 months, once every 9 months, once every 10 months, once every 11 months, or once every 12 months. In such embodiments, the therapeutically effective doses of the antagonist anti-CD40 antibody or antigen-binding fragment thereof fall within the lower dosing range (i.e., 0.003 mg/kg to about 20 mg/kg), particularly when the subsequent doses are administered at more frequent intervals, for example, once every two weeks to once every month, or within the higher dosing range (i.e., from about 20 mg/kg to about 50 mg/kg), particularly when the subsequent doses are administered at less frequent intervals, for example, where subsequent doses are administered about one month to about 12 months apart.

The antagonist anti-CD40 antibodies present in the pharmaceutical compositions described herein for use in the methods of the invention may be native or obtained by recombinant techniques, and may be from any source, including mammalian sources such as, e.g., mouse, rat, rabbit, primate, pig, and human. Preferably such polypeptides are derived from a human source, and more preferably are recombinant, human proteins from hybridoma cell lines.

The pharmaceutical compositions useful in the methods of the invention may comprise biologically active variants of the antagonist anti-CD40 antibodies of the

invention. Such variants should retain the desired biological activity of the native polypeptide such that the pharmaceutical composition comprising the variant polypeptide has the same therapeutic effect as the pharmaceutical composition comprising the native polypeptide when administered to a subject. That is, the variant anti-CD40 antibody will
5 serve as a therapeutically active component in the pharmaceutical composition in a manner similar to that observed for the native antagonist antibody, for example CHIR-5.9 or CHIR-12.12 as expressed by the hybridoma cell line 5.9 or 12.12, respectively. Methods are available in the art for determining whether a variant anti-CD40 antibody retains the desired biological activity, and hence serves as a therapeutically active
10 component in the pharmaceutical composition. Biological activity of antibody variants can be measured using assays specifically designed for measuring activity of the native antagonist antibody, including assays described in the present invention.

Any pharmaceutical composition comprising an antagonist anti-CD40 antibody having the binding properties described herein as the therapeutically active component
15 can be used in the methods of the invention. Thus liquid, lyophilized, or spray-dried compositions comprising one or more of the antagonist anti-CD40 antibodies of the invention may be prepared as an aqueous or nonaqueous solution or suspension for subsequent administration to a subject in accordance with the methods of the invention. Each of these compositions will comprise at least one of the antagonist anti-CD40
20 antibodies of the present invention as a therapeutically or prophylactically active component. By "therapeutically or prophylactically active component" is intended the anti-CD40 antibody is specifically incorporated into the composition to bring about a desired therapeutic or prophylactic response with regard to treatment, prevention, or diagnosis of a disease or condition within a subject when the pharmaceutical composition
25 is administered to that subject. Preferably the pharmaceutical compositions comprise appropriate stabilizing agents, bulking agents, or both to minimize problems associated with loss of protein stability and biological activity during preparation and storage.

Formulants may be added to pharmaceutical compositions comprising an antagonist anti-CD40 antibody of the invention. These formulants may include, but are
30 not limited to, oils, polymers, vitamins, carbohydrates, amine acids, salts, buffers, albumin, surfactants, or bulking agents. Preferably carbohydrates include sugar or sugar

alcohols such as mono-, di-, or polysaccharides, or water soluble glucans. The saccharides or glucans can include fructose, glucose, mannose, sorbose, xylose, maltose, sucrose, dextran, pullulan, dextrin, α and β cyclodextrin, soluble starch, hydroxyethyl starch, and carboxymethylcellulose, or mixtures thereof. "Sugar alcohol" is defined as a C₄ to C₈ hydrocarbon having a hydroxyl group and includes galactitol, inositol, mannitol, xylitol, sorbitol, glycerol, and arabitol. These sugars or sugar alcohols may be used individually or in combination. The sugar or sugar alcohol concentration is between 1.0% and 7% w/v., more preferably between 2.0% and 6.0% w/v. Preferably amino acids include levorotary (L) forms of carnitine, arginine, and betaine; however, other amino acids may be added. Preferred polymers include polyvinylpyrrolidone (PVP) with an average molecular weight between 2,000 and 3,000, or polyethylene glycol (PEG) with an average molecular weight between 3,000 and 5,000. Surfactants that can be added to the formulation are shown in EP Nos. 270,799 and 268,110.

Additionally, antibodies can be chemically modified by covalent conjugation to a polymer to increase their circulating half-life, for example. Preferred polymers, and methods to attach them to peptides, are shown in U.S. Patent Nos. 4,766,106; 4,179,337; 4,495,285; and 4,609,546; which are all hereby incorporated by reference in their entireties. Preferred polymers are polyoxyethylated polyols and polyethylene glycol (PEG). PEG is soluble in water at room temperature and has the general formula: R(O--CH₂--CH₂)_nO--R where R can be hydrogen, or a protective group such as an alkyl or alkanol group. Preferably, the protective group has between 1 and 8 carbons, more preferably it is methyl. The symbol n is a positive integer, preferably between 1 and 1,000, more preferably between 2 and 500. The PEG has a preferred average molecular weight between 1,000 and 40,000, more preferably between 2,000 and 20,000, most preferably between 3,000 and 12,000. Preferably, PEG has at least one hydroxy group, more preferably it is a terminal hydroxy group. It is this hydroxy group which is preferably activated to react with a free amino group on the inhibitor. However, it will be understood that the type and amount of the reactive groups may be varied to achieve a covalently conjugated PEG/antibody of the present invention.

Water-soluble polyoxyethylated polyols are also useful in the present invention. They include polyoxyethylated sorbitol, polyoxyethylated glucose, polyoxyethylated

glycerol (POG), and the like. POG is preferred. One reason is because the glycerol backbone of polyoxyethylated glycerol is the same backbone occurring naturally in, for example, animals and humans in mono-, di-, triglycerides. Therefore, this branching would not necessarily be seen as a foreign agent in the body. The POG has a preferred molecular weight in the same range as PEG. The structure for POG is shown in Knauf *et al.* (1988) *J. Bio. Chem.* 263:15064-15070, and a discussion of POG/IL-2 conjugates is found in U.S. Patent No. 4,766,106, both of which are hereby incorporated by reference in their entireties.

Another drug delivery system for increasing circulatory half-life is the liposome. Methods of preparing liposome delivery systems are discussed in Gabizon *et al.* (1982) *Cancer Research* 42:4734; Cafiso (1981) *Biochem Biophys Acta* 649:129; and Szoka (1980) *Ann. Rev. Biophys. Eng.* 9:467. Other drug delivery systems are known in the art and are described in, *e.g.*, Poznansky *et al.* (1980) *Drug Delivery Systems* (R.L. Juliano, ed., Oxford, N.Y.) pp. 253-315; Poznansky (1984) *Pharm Revs* 36:277.

The formulants to be incorporated into a pharmaceutical composition should provide for the stability of the antagonist anti-CD40 antibody or antigen-binding fragment thereof. That is, the antagonist anti-CD40 antibody or antigen-binding fragment thereof should retain its physical and/or chemical stability and have the desired biological activity, *i.e.*, one or more of the antagonist activities defined herein above, including, but not limited to, inhibition of immunoglobulin secretion by normal human peripheral B cells stimulated by T cells; inhibition of survival and/or proliferation of normal human peripheral B cells stimulated by Jurkat T cells; inhibition of survival and/or proliferation of normal human peripheral B cells stimulated by CD40L-expressing cells or soluble CD40 ligand (sCD40L); inhibition of "survival" anti-apoptotic intracellular signals in any cell stimulated by sCD40L or solid-phase CD40L; inhibition of CD40 signal transduction in any cell upon ligation with sCD40L or solid-phase CD40L; and inhibition of proliferation of human malignant B cells as noted elsewhere herein.

Methods for monitoring protein stability are well known in the art. See, for example, Jones (1993) *Adv. Drug Delivery Rev.* 10:29-90; Lee, ed. (1991) *Peptide and Protein Drug Delivery* (Marcel Dekker, Inc., New York, New York); and the stability assays disclosed herein below. Generally, protein stability is measured at a chosen

temperature for a specified period of time. In preferred embodiments, a stable antibody pharmaceutical formulation provides for stability of the antagonist anti-CD40 antibody or antigen-binding fragment thereof when stored at room temperature (about 25°C) for at least 1 month, at least 3 months, or at least 6 months, and/or is stable at about 2-8°C for at least 6 months, at least 9 months, at least 12 months, at least 18 months, at least 24 months.

A protein such as an antibody, when formulated in a pharmaceutical composition, is considered to retain its physical stability at a given point in time if it shows no visual signs (i.e., discoloration or loss of clarity) or measurable signs (for example, using size-exclusion chromatography (SEC) or UV light scattering) of precipitation, aggregation, and/or denaturation in that pharmaceutical composition. With respect to chemical stability, a protein such as an antibody, when formulated in a pharmaceutical composition, is considered to retain its chemical stability at a given point in time if measurements of chemical stability are indicative that the protein (i.e., antibody) retains the biological activity of interest in that pharmaceutical composition. Methods for monitoring changes in chemical stability are well known in the art and include, but are not limited to, methods to detect chemically altered forms of the protein such as result from clipping, using, for example, SDS-PAGE, SEC, and/or matrix-assisted laser desorption ionization/time of flight mass spectrometry; and degradation associated with changes in molecular charge (for example, associated with deamidation), using, for example, ion-exchange chromatography. See, for example, the methods disclosed herein below.

An antagonist anti-CD40 antibody or antigen-binding fragment thereof, when formulated in a pharmaceutical composition, is considered to retain a desired biological activity at a given point in time if the desired biological activity at that time is within about 30%, preferably within about 20% of the desired biological activity exhibited at the time the pharmaceutical composition was prepared as determined in a suitable assay for the desired biological activity. Assays for measuring the desired biological activity of the antagonist anti-CD40 antibodies disclosed herein, and antigen-binding fragments thereof, can be performed as described in the Examples herein. See also the assays described in Schultze *et al.* (1998) *Proc. Natl. Acad. Sci. USA* 92:8200-8204; Denton *et al.* (1998)

Pediatr. Transplant. 2:6-15; Evans *et al.* (2000) *J. Immunol.* 164:688-697; Noelle (1998) *Agents Actions Suppl.* 49:17-22; Lederman *et al.* (1996) *Curr. Opin. Hematol.* 3:77-86; Coligan *et al.* (1991) *Current Protocols in Immunology* 13:12; Kwekkeboom *et al.* (1993) *Immunology* 79:439-444; and U.S. Patent Nos. 5,674,492 and 5,847,082; herein

5 incorporated by reference.

In some embodiments of the invention, the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof is formulated in a liquid pharmaceutical formulation. The antagonist anti-CD40 antibody or antigen binding fragment thereof can be prepared using any
10 method known in the art, including those methods disclosed herein above. In one embodiment, the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof is recombinantly produced in a CHO cell line.

Following its preparation and purification, the antagonist anti-CD40 antibody or
15 antigen-binding fragment thereof can be formulated as a liquid pharmaceutical formulation in the manner set forth herein. Where the antagonist anti-CD40 antibody or antigen-binding fragment thereof is to be stored prior to its formulation, it can be frozen, for example, at $\leq -20^{\circ}\text{C}$, and then thawed at room temperature for further formulation. The liquid pharmaceutical formulation comprises a therapeutically effective amount of
20 the antagonist anti-CD40 antibody or antigen-binding fragment thereof. The amount of antibody or antigen-binding fragment thereof present in the formulation takes into consideration the route of administration and desired dose volume.

In this manner, the liquid pharmaceutical composition comprises the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 antibody, or antigen-
25 binding fragment thereof at a concentration of about 0.1 mg/ml to about 50.0 mg/ml, about 0.5 mg/ml to about 40.0 mg/ml, about 1.0 mg/ml to about 30.0 mg/ml, about 5.0 mg/ml to about 25.0 mg/ml, about 5.0 mg/ml to about 20.0 mg/ml, or about 15.0 mg/ml to about 25.0 mg/ml. In some embodiments, the liquid pharmaceutical composition comprises the antagonist anti-CD40 antibody or antigen-binding fragment thereof at a
30 concentration of about 0.1 mg/ml to about 5.0 mg/ml, about 5.0 mg/ml to about 10.0 mg/ml, about 10.0 mg/ml to about 15.0 mg/ml, about 15.0 mg/ml to about 20.0 mg/ml,

about 20.0 mg/ml to about 25.0 mg/ml, about 25.0 mg/ml to about 30.0 mg/ml, about 30.0 mg/ml to about 35.0 mg/ml, about 35.0 mg/ml to about 40.0 mg/ml, about 40.0 mg/ml to about 45.0 mg/ml, or about 45.0 mg/ml to about 50.0 mg/ml. In other embodiments, the liquid pharmaceutical composition comprises the antagonist anti-CD40 antibody or antigen-binding fragment thereof at a concentration of about 15.0 mg/ml, about 16.0 mg/ml, about 17.0 mg/ml, about 18.0 mg/ml, about 19.0 mg/ml, about 20.0 mg/ml, about 21.0 mg/ml, about 22.0 mg/ml, about 23.0 mg/ml, about 24.0 mg/ml, or about 25.0 mg/ml. The liquid pharmaceutical composition comprises the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 antibody, or antigen-binding fragment thereof and a buffer that maintains the pH of the formulation in the range of about pH 5.0 to about pH 7.0, including about pH 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7.0, and other such values within the range of about pH 5.0 to about pH 7.0. In some embodiments, the buffer maintains the pH of the formulation in the range of about pH 5.0 to about pH 6.5, about pH 5.0 to about pH 6.0, about pH 5.0 to about pH 5.5, about pH 5.5 to about 7.0, about pH 5.5 to about pH 6.5, or about pH 5.5 to about pH 6.0.

Any suitable buffer that maintains the pH of the liquid anti-CD40 antibody formulation in the range of about pH 5.0 to about pH 7.0 can be used in the formulation, so long as the physicochemical stability and desired biological activity of the antibody are retained as noted herein above. Suitable buffers include, but are not limited to, conventional acids and salts thereof, where the counter ion can be, for example, sodium, potassium, ammonium, calcium, or magnesium. Examples of conventional acids and salts thereof that can be used to buffer the pharmaceutical liquid formulation include, but are not limited to, succinic acid or succinate, citric acid or citrate, acetic acid or acetate, tartaric acid or tartarate, phosphoric acid or phosphate, gluconic acid or gluconate, glutamic acid or glutamate, aspartic acid or aspartate, maleic acid or maleate, and malic acid or malate buffers. The buffer concentration within the formulation can be from about 1 mM to about 50 mM, including about 1 mM, 2 mM, 5 mM, 8 mM, 10 mM, 15 mM, 20 mM, 25 mM, 30 mM, 35 mM, 40 mM, 45 mM, 50 mM, or other such values within the range of about 1 mM to about 50 mM. In some embodiments, the buffer concentration within the formulation is from about 5 mM to about 15 mM, including

about 5 mM, 6 mM, 7 mM, 8 mM, 9 mM, 10 mM, 11 mM, 12 mM, 13 mM, 14 mM, 15 mM, or other such values within the range of about 5 mM to about 15 mM.

In some embodiments of the invention, the liquid pharmaceutical formulation comprises a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof and succinate buffer or citrate buffer at a concentration that maintains the pH of the formulation in the range of about pH 5.0 to about pH 7.0, preferably about pH 5.0 to about pH 6.5. By "succinate buffer" or "citrate buffer" is intended a buffer comprising a salt of succinic acid or a salt of citric acid, respectively. In a preferred embodiment, the succinate or citrate counterion is the sodium cation, and thus the buffer is sodium succinate or sodium citrate, respectively. However, any cation is expected to be effective. Other possible succinate or citrate cations include, but are not limited to, potassium, ammonium, calcium, and magnesium. As noted above, the succinate or citrate buffer concentration within the formulation can be from about 1 mM to about 50 mM, including about 1 mM, 2 mM, 5 mM, 8 mM, 10 mM, 15 mM, 20 mM, 25 mM, 30 mM, 35 mM, 40 mM, 45 mM, 50 mM, or other such values within the range of about 1 mM to about 50 mM. In some embodiments, the buffer concentration within the formulation is from about 5 mM to about 15 mM, including about 5 mM, 6 mM, 7 mM, 8 mM, 9 mM, 10 mM, 11 mM, 12 mM, 13 mM, 14 mM, or about 15 mM. In other embodiments, the liquid pharmaceutical formulation comprises the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof at a concentration of about 0.1 mg/ml to about 50.0 mg/ml, or about 5.0 mg/ml to about 25.0 mg/ml, and succinate or citrate buffer, for example, sodium succinate or sodium citrate buffer, at a concentration of about 1 mM to about 20 mM, about 5 mM to about 15 mM, preferably about 10 mM.

Where it is desirable for the liquid pharmaceutical formulation to be near isotonic, the liquid pharmaceutical formulation comprising a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, and a buffer to maintain the pH of the formulation within the range of about pH 5.0 to about pH 7.0 can further comprise an amount of an isotonicizing agent sufficient to render the formulation near

isotonic. By "near isotonic" is intended the aqueous formulation has an osmolarity of about 240 mmol/kg to about 360 mmol/kg, preferably about 240 to about 340 mmol/kg, more preferably about 250 to about 330 mmol/kg, even more preferably about 260 to about 320 mmol/kg, still more preferably about 270 to about 310 mmol/kg. Methods of determining the isotonicity of a solution are known to those skilled in the art. See, for example, Setnikar *et al.* (1959) *J. Am. Pharm. Assoc.* 48:628.

Those skilled in the art are familiar with a variety of pharmaceutically acceptable solutes useful in providing isotonicity in pharmaceutical compositions. The isotonicizing agent can be any reagent capable of adjusting the osmotic pressure of the liquid pharmaceutical formulation of the present invention to a value nearly equal to that of a body fluid. It is desirable to use a physiologically acceptable isotonicizing agent. Thus, the liquid pharmaceutical formulation comprising a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, and a buffer to maintain the pH of the formulation within the range of about pH 5.0 to about pH 7.0, can further comprise components that can be used to provide isotonicity, for example, sodium chloride; amino acids such as alanine, valine, and glycine; sugars and sugar alcohols (polyols), including, but not limited to, glucose, dextrose, fructose, sucrose, maltose, mannitol, trehalose, glycerol, sorbitol, and xylitol; acetic acid, other organic acids or their salts, and relatively minor amounts of citrates or phosphates. The ordinary skilled person would know of additional agents that are suitable for providing optimal tonicity of the liquid formulation.

In some preferred embodiments, the liquid pharmaceutical formulation comprising a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, and a buffer to maintain the pH of the formulation within the range of about pH 5.0 to about pH 7.0, further comprises sodium chloride as the isotonicizing agent. The concentration of sodium chloride in the formulation will depend upon the contribution of other components to tonicity. In some embodiments, the concentration of sodium chloride is about 50 mM to about 300 mM, about 50 mM to about 250 mM, about 50 mM to about 200 mM, about 50 mM to about 175 mM, about 50 mM to about 150 mM, about 75 mM to about 175 mM, about 75 mM to about 150 mM, about 100 mM to

about 175 mM, about 100 mM to about 200 mM, about 100 mM to about 150 mM, about 125 mM to about 175 mM, about 125 mM to about 150 mM, about 130 mM to about 170 mM, about 130 mM to about 160 mM, about 135 mM to about 155 mM, about 140 mM to about 155 mM, or about 145 mM to about 155 mM. In one such embodiment, the concentration of sodium chloride is about 150 mM. In other such embodiments, the concentration of sodium chloride is about 150 mM, the buffer is sodium succinate or sodium citrate buffer at a concentration of about 5 mM to about 15 mM, the liquid pharmaceutical formulation comprises a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, and the formulation has a pH of about pH 5.0 to about pH 7.0, about pH 5.0 to about pH 6.0, or about pH 5.5 to about pH 6.5. In other embodiments, the liquid pharmaceutical formulation comprises the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, at a concentration of about 0.1 mg/ml to about 50.0 mg/ml or about 5.0 mg/ml to about 25.0 mg/ml, about 150 mM sodium chloride, and about 10 mM sodium succinate or sodium citrate, at a pH of about pH 5.5.

Protein degradation due to freeze thawing or mechanical shearing during processing of a liquid pharmaceutical formulations of the present invention can be inhibited by incorporation of surfactants into the formulation in order to lower the surface tension at the solution-air interface. Thus, in some embodiments, the liquid pharmaceutical formulation comprises a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, a buffer to maintain the pH of the formulation within the range of about pH 5.0 to about pH 7.0, and further comprises a surfactant. In other embodiments, the liquid pharmaceutical formulation comprises a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, a buffer to maintain the pH of the formulation within the range of about pH 5.0 to about pH 7.0, an isotonicizing agent such as sodium chloride at a concentration of about 50 mM to about 300 mM, and further comprises a surfactant.

Typical surfactants employed are nonionic surfactants, including polyoxyethylene sorbitol esters such as polysorbate 80 (Tween 80) and polysorbate 20 (Tween 20); polyoxypropylene-polyoxyethylene esters such as Pluronic F68; polyoxyethylene alcohols such as Brij 35; simethicone; polyethylene glycol such as PEG400;

5 lysophosphatidylcholine; and polyoxyethylene-p-t-octylphenol such as Triton X-100. Classic stabilization of pharmaceuticals by surfactants or emulsifiers is described, for example, in Levine *et al.* (1991) *J. Parenteral Sci. Technol.* 45(3):160-165, herein incorporated by reference. A preferred surfactant employed in the practice of the present invention is polysorbate 80. Where a surfactant is included, it is typically added in an
10 amount from about 0.001 % to about 1.0% (w/v), about 0.001% to about 0.5%, about 0.001% to about 0.4%, about 0.001% to about 0.3%, about 0.001% to about 0.2%, about 0.005% to about 0.5%, about 0.005% to about 0.2%, about 0.01% to about 0.5%, about 0.01% to about 0.2%, about 0.03% to about 0.5%, about 0.03% to about 0.3%, about 0.05% to about 0.5%, or about 0.05% to about 0.2%.

15 Thus, in some embodiments, the liquid pharmaceutical formulation comprises a therapeutically effective amount of the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, the buffer is sodium succinate or sodium citrate buffer at a concentration of about 1 mM to about 50 mM, about 5 mM to about 25 mM, or about 5 mM to about 15 mM; the
20 formulation has a pH of about pH 5.0 to about pH 7.0, about pH 5.0 to about pH 6.0, or about pH 5.5 to about pH 6.5; and the formulation further comprises a surfactant, for example, polysorbate 80, in an amount from about 0.001% to about 1.0% or about 0.001% to about 0.5%. Such formulations can optionally comprise an isotonicizing agent, such as sodium chloride at a concentration of about 50 mM to about 300 mM, about 50
25 mM to about 200 mM, or about 50 mM to about 150 mM. In other embodiments, the liquid pharmaceutical formulation comprises the antagonist anti-CD40 antibody, for example, the CHIR-12.12 or CHIR-5.9 monoclonal antibody, or antigen-binding fragment thereof, at a concentration of about 0.1 mg/ml to about 50.0 mg/ml or about 5.0 mg/ml to about 25.0 mg/ml, including about 20.0 mg/ml; about 50 mM to about 200 mM
30 sodium chloride, including about 150 mM sodium chloride; sodium succinate or sodium citrate at about 5 mM to about 20 mM, including about 10 mM sodium succinate or

sodium citrate; sodium chloride at a concentration of about 50 mM to about 200 mM, including about 150 mM; and optionally a surfactant, for example, polysorbate 80, in an amount from about 0.001% to about 1.0%, including about 0.001% to about 0.5%; where the liquid pharmaceutical formulation has a pH of about pH 5.0 to about pH 7.0, about
5 pH 5.0 to about pH 6.0, about pH 5.0 to about pH 5.5, about pH 5.5 to about pH 6.5, or about pH 5.5 to about pH 6.0.

The liquid pharmaceutical formulation can be essentially free of any preservatives and other carriers, excipients, or stabilizers noted herein above. Alternatively, the formulation can include one or more preservatives, for example, antibacterial agents,
10 pharmaceutically acceptable carriers, excipients, or stabilizers described herein above provided they do not adversely affect the physicochemical stability of the antagonist anti-CD40 antibody or antigen-binding fragment thereof. Examples of acceptable carriers, excipients, and stabilizers include, but are not limited to, additional buffering agents, co-solvents, surfactants, antioxidants including ascorbic acid and methionine, chelating
15 agents such as EDTA, metal complexes (for example, Zn-protein complexes), and biodegradable polymers such as polyesters. A thorough discussion of formulation and selection of pharmaceutically acceptable carriers, stabilizers, and isomolytes can be found in *Remington's Pharmaceutical Sciences* (18th ed.; Mack Publishing Company, Eaton, Pennsylvania, 1990), herein incorporated by reference.

20 After the liquid pharmaceutical formulation or other pharmaceutical composition described herein is prepared, it can be lyophilized to prevent degradation. Methods for lyophilizing liquid compositions are known to those of ordinary skill in the art. Just prior to use, the composition may be reconstituted with a sterile diluent (Ringer's solution, distilled water, or sterile saline, for example) that may include additional ingredients.
25 Upon reconstitution, the composition is preferably administered to subjects using those methods that are known to those skilled in the art.

Use of Antagonist Anti-CD40 Antibodies in the Manufacture of Medicaments

The present invention also provides for the use of an antagonist anti-CD40
30 antibody or antigen-binding fragment thereof in the manufacture of a medicament for treating a subject for a cancer characterized by neoplastic B cell growth, wherein the

medicament is coordinated with treatment with at least one other cancer therapy. Cancers characterized by neoplastic B cell growth include, but are not limited to, the B cell-related cancers discussed herein above, for example, non-Hodgkin's lymphoma, chronic lymphocytic leukemia, multiple myeloma, B cell lymphoma, high-grade B cell

5 lymphoma, intermediate-grade B cell lymphoma, low-grade B cell lymphoma, B cell acute lymphoblastic leukemia, myeloblastic leukemia, Hodgkin's disease, plasmacytoma, follicular lymphoma, follicular small cleaved lymphoma, follicular large cell lymphoma, follicular mixed small cleaved lymphoma, diffuse small cleaved cell lymphoma, diffuse small lymphocytic lymphoma, prolymphocytic leukemia,
10 lymphoplasmacytic lymphoma, marginal zone lymphoma, mucosal associated lymphoid tissue lymphoma, monocytoid B cell lymphoma, splenic lymphoma, hairy cell leukemia, diffuse large cell lymphoma, mediastinal large B cell lymphoma, lymphomatoid granulomatosis, intravascular lymphomatosis, diffuse mixed cell lymphoma, diffuse large cell lymphoma, immunoblastic lymphoma, Burkitt's lymphoma, AIDS-related
15 lymphoma, and mantle cell lymphoma.

By "coordinated" is intended the medicament comprising the antagonist anti-CD40 antibody or antigen-binding fragment thereof is to be used either prior to, during, or after treatment of the subject with at least one other cancer therapy. Examples of other cancer therapies include, but are not limited to, surgery; radiation therapy; chemotherapy,
20 optionally in combination with autologous bone marrow transplant, where suitable chemotherapeutic agents include, but are not limited to, fludarabine or fludarabine phosphate, chlorambucil, vincristine, pentostatin, 2-chlorodeoxyadenosine (cladribine), cyclophosphamide, doxorubicin, prednisone, and combinations thereof, for example, anthracycline-containing regimens such as CAP (cyclophosphamide, doxorubicin plus
25 prednisone), CHOP (cyclophosphamide, vincristine, prednisone plus doxorubicin), VAD (vincristine, doxorubicin, plus dexamethasone), MP (melphalan plus prednisone), and other cytotoxic and/or therapeutic agents used in chemotherapy such as mitoxantrone, daunorubicin, idarubicin, asparaginase, and antimetabolites, including, but not limited to, cytarabine, methotrexate, 5-fluorouracil decarbazine, 6-thioguanine, 6-mercaptopurine,
30 and nelarabine; other anti-cancer monoclonal antibody therapy (for example, alemtuzumab (Campath[®]) or other anti-CD52 antibody targeting the CD52 cell-surface

glycoprotein on malignant B cells; rituximab (Rituxan[®]), the fully human antibody HuMax-CD20, R-1594, IMMU-106, TRU-015, AME-133, tositumomab/I-131 tositumomab (Bexxar[®]), ibritumomab tiuxetan (Zevalin[®]), or any other therapeutic anti-CD20 antibody targeting the CD20 antigen on malignant B cells; anti-CD19 antibody (for example, MT103, a bispecific antibody); anti-CD22 antibody (for example, the humanized monoclonal antibody epratuzumab); bevacizumab (Avastin[®]) or other anti-cancer antibody targeting human vascular endothelial growth factor; anti-CD22 antibody targeting the CD22 antigen on malignant B cells (for example, the monoclonal antibody BL-22, an alphaCD22 toxin); α -M-CSF antibody targeting macrophage colony stimulating factor; antibodies targeting the receptor activator of nuclear factor-kappaB (RANK) and its ligand (RANKL), which are overexpressed in multiple myeloma; anti-CD23 antibody targeting the CD23 antigen on malignant B cells (for example, IDEC-152); anti-CD38 antibody targeting the CD38 antigen on malignant B cells; antibodies targeting major histocompatibility complex class II receptors (anti-MHC antibodies) expressed on malignant B cells; other anti-CD40 antibodies (for example, SGN-40) targeting the CD40 antigen on malignant B cells; and antibodies targeting tumor necrosis factor-related apoptosis-inducing ligand receptor 1 (TRAIL-R1) (for example, the agonistic human monoclonal antibody HGS-ETR1) expressed on a number of solid tumors and tumors of hematopoietic origin); small molecule-based cancer therapy, including, but not limited to, microtubule and/or topoisomerase inhibitors (for example, the mitotic inhibitor dolastatin and dolastatin analogues; the tubulin-binding agent T900607; XL119; and the topoisomerase inhibitor aminocamptothecin), SDX-105 (bendamustine hydrochloride), ixabepilone (an epothilone analog, also referred to as BMS-247550), protein kinase C inhibitors, for example, midostaurin ((PKC-412, CGP 41251, N-benzoylstauroporine), pixantrone, eloxatin (an antineoplastic agent), ganite (gallium nitrate), Thalomid[®] (thalidomide), immunomodulatory derivatives of thalidomide (for example, revlimid (formerly revimid)), Affinitak[™] (antisense inhibitor of protein kinase C- α), SDX-101 (R-etodolac, inducing apoptosis of malignant lymphocytes), second-generation purine nucleoside analogs such as clofarabine, inhibitors of production of the protein Bcl-2 by cancer cells (for example, the antisense agents oblimersen and Genasense[®]), proteasome inhibitors (for example, Velcade[™]

(bortezomib)), small molecule kinase inhibitors (for example, CHIR-258), small molecule VEGF inhibitors (for example, ZD-6474), small molecule inhibitors of heat shock protein (HSP) 90 (for example, 17-AAG), small molecule inhibitors of histone deacetylases (for example, hybrid/polar cytodifferentiation HPC) agents such as
5 suberanilohydroxamic acid (SAHA), and FR-901228) and apoptotic agents such as Trisenox[®] (arsenic trioxide) and Xcytrin[®] (motexafin gadolinium); vaccine /immunotherapy-based cancer therapies, including, but not limited to, vaccine approaches (for example, Id-KLH, oncoplage, vitalethine), personalized immunotherapy or active idiotype immunotherapy (for example, MyVax[®] Personalized Immunotherapy, formally
10 designated GTOP-99), Promune[®] (CpG 7909, a synthetic agonist for toll-like receptor 9 (TLR9)), interferon-alpha therapy, interleukin-2 (IL-2) therapy, IL-12 therapy; IL-15 therapy, and IL-21 therapy; steroid therapy; or other cancer therapy; where treatment with the additional cancer therapy, or additional cancer therapies, occurs prior to, during, or subsequent to treatment of the subject with the medicament comprising the antagonist
15 anti-CD40 antibody or antigen-binding fragment thereof, as noted herein above.

In some embodiments, the present invention provides for the use of the anti-CD40 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof in the manufacture of a medicament for treating a B cell lymphoma, for example non-Hodgkin's lymphoma, in a subject, wherein the medicament
20 is coordinated with treatment with at least one other cancer therapy selected from the group consisting of chemotherapy, anti-cancer antibody therapy, small molecule-based cancer therapy, and vaccine/immunotherapy-based cancer therapy, wherein the medicament is to be used either prior to, during, or after treatment of the subject with the other cancer therapy or, in the case of multiple combination therapies, either prior to,
25 during, or after treatment of the subject with the other cancer therapies.

Thus, for example, in some embodiments, the invention provides for the use of the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof, in the manufacture of a medicament for treating a B cell lymphoma, for example, non-Hodgkin's lymphoma, in a subject, wherein the medicament is coordinated with
30 treatment with chemotherapy, where the chemotherapeutic agent is selected from the group consisting of cytoxan, doxorubicin, vincristine, prednisone, and combinations

thereof, for example CHOP. In other embodiments, the invention provides for the use of the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof, in the manufacture of a medicament for treating a B cell lymphoma, for example non-Hodgkin's lymphoma, in a subject, wherein the medicament is coordinated with

5 treatment with at least one other anti-cancer antibody selected from the group consisting of alemtuzumab (Campath®) or other anti-CD52 antibody targeting the CD52 cell-surface glycoprotein on malignant B cells; rituximab (Rituxan®), the fully human antibody HuMax-CD20, R-1594, IMMU-106, TRU-015, AME-133, tositumomab/I-131 tositumomab (Bexxar®), ibritumomab tiuxetan (Zevalin®), or any other therapeutic anti-
10 CD20 antibody targeting the CD20 antigen on malignant B cells; anti-CD19 antibody (for example, MT103, a bispecific antibody); anti-CD22 antibody (for example, the humanized monoclonal antibody epratuzumab); bevacizumab (Avastin®) or other anti-cancer antibody targeting human vascular endothelial growth factor; and any combinations thereof; wherein the medicament is to be used either prior to, during, or
15 after treatment of the subject with the other cancer therapy or, in the case of multiple combination therapies, either prior to, during, or after treatment of the subject with the other cancer therapies.

In yet other embodiments, the present invention provides for the use of the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof, in
20 the manufacture of a medicament for treating a B cell lymphoma, for example non-Hodgkin's lymphoma, in a subject, wherein the medicament is coordinated with treatment with at least one other small molecule-based cancer therapy selected from the group consisting of microtubule and/or topoisomerase inhibitors (for example, the mitotic inhibitor dolastatin and dolastatin analogues; the tubulin-binding agent T900607; XL119;
25 and the topoisomerase inhibitor aminocamptothecin), SDX-105 (bendamustine hydrochloride), ixabepilone (an epothilone analog, also referred to as BMS-247550), protein kinase C inhibitors, for example, midostaurin ((PKC-412, CGP 41251, N-benzoylstaurosporine), pixantrone, eloxatin (an antineoplastic agent), ganite (gallium nitrate), Thalomid® (thalidomide), an apoptotic agent such as Xcytrin® (motexafin
30 gadolinium), inhibitors of production of the protein Bcl-2 by cancer cells (for example, the antisense agents oblimersen and Genasense®), nelarabine, and any combinations

thereof; wherein the medicament is to be used either prior to, during, or after treatment of the subject with the other cancer therapy or, in the case of multiple combination therapies, either prior to, during, or after treatment of the subject with the other cancer therapies.

5 In still other embodiments, the present invention provides for the use of the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof, in the manufacture of a medicament for treating a B cell lymphoma, for example non-Hodgkin's lymphoma, in a subject, wherein the medicament is coordinated with treatment with at least one other vaccine/immunotherapy-based cancer therapy selected
10 from the group consisting of vaccine approaches (for example, Id-KLH, oncopophage, vitalethine), personalized immunotherapy or active idiotype immunotherapy (for example, MyVax[®] Personalized Immunotherapy, formally designated GTOP-99), Promune[®] (CpG 7909, a synthetic agonist for toll-like receptor 9 (TLR9)), interleukin-2 (IL-2) therapy, IL-12 therapy; IL-15 therapy, and IL-21 therapy, and any combinations
15 thereof; wherein the medicament is to be used either prior to, during, or after treatment of the subject with the other cancer therapy or, in the case of multiple combination therapies, either prior to, during, or after treatment of the subject with the other cancer therapies.

In some embodiments, the present invention provides for the use of the anti-CD40
20 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof in the manufacture of a medicament for treating a B cell-related leukemia, for example B-cell acute lymphocytic leukemia (B-ALL), in a subject, wherein the medicament is coordinated with treatment with at least one other cancer therapy selected from the group consisting of chemotherapy and anti-metabolite therapy, wherein
25 the medicament is to be used either prior to, during, or after treatment of the subject with the other cancer therapy or, in the case of multiple combination therapies, either prior to, during, or after treatment of the subject with the other cancer therapies. Examples of such embodiments include, but are not limited to, those instances where the medicament comprising the antagonist anti-CD40 antibody, for example, the monoclonal antibody
30 CHIR-12.12 or CHIR-5.9, or antigen-binding fragment thereof is coordinated with treatment with a chemotherapeutic agent or anti-metabolite selected from the group

consisting of cytoxan, doxorubicin, vincristine, prednisone, cytarabine, mitoxantrone, idarubicin, asparaginase, methotrexate, 6-thioguanine, 6-mercaptopurine, and combinations thereof; wherein the medicament is to be used either prior to, during, or after treatment of the subject with the other cancer therapy or, in the case of multiple combination therapies, either prior to, during, or after treatment of the subject with the other cancer therapies. In one such example, the medicament is coordinated with treatment with cytarabine plus daunorubicin, cytarabine plus mitoxantrone, and/or cytarabine plus idarubicin; wherein the medicament is to be used either prior to, during, or after treatment of the B-ALL subject with the other cancer therapy or, in the case of multiple combination therapies, either prior to, during, or after treatment of the subject with the other cancer therapies.

The invention also provides for the use of an antagonist anti-CD40 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9 disclosed herein, or antigen-binding fragment thereof in the manufacture of a medicament for treating a subject for a cancer characterized by neoplastic B cell growth, including the B cell-related cancers described herein above, wherein the medicament is used in a subject that has been pretreated with at least one other cancer therapy. By “pretreated” or “pretreatment” is intended the subject has received one or more other cancer therapies (i.e., been treated with at least one other cancer therapy) prior to receiving the medicament comprising the antagonist anti-CD40 antibody or antigen-binding fragment thereof. “Pretreated” or “pretreatment” includes subjects that have been treated with at least one other cancer therapy within 2 years, within 18 months, within 1 year, within 6 months, within 2 months, within 6 weeks, within 1 month, within 4 weeks, within 3 weeks, within 2 weeks, within 1 week, within 6 days, within 5 days, within 4 days, within 3 days, within 2 days, or even within 1 day prior to initiation of treatment with the medicament comprising the antagonist anti-CD40 antibody, for example, the monoclonal antibody CHIR-12.12 or CHIR-5.9 disclosed herein, or antigen-binding fragment thereof. It is not necessary that the subject was a responder to pretreatment with the prior cancer therapy, or prior cancer therapies. Thus, the subject that receives the medicament comprising the antagonist anti-CD40 antibody or antigen-binding fragment thereof could have responded, or could have failed to respond (i.e. the cancer was refractory), to

pretreatment with the prior cancer therapy, or to one or more of the prior cancer therapies where pretreatment comprised multiple cancer therapies. Examples of other cancer therapies for which a subject can have received pretreatment prior to receiving the medicament comprising the antagonist anti-CD40 antibody or antigen-binding fragment thereof include, but are not limited to, surgery; radiation therapy; chemotherapy, optionally in combination with autologous bone marrow transplant, where suitable chemotherapeutic agents include, but are not limited to, those listed herein above; other anti-cancer monoclonal antibody therapy, including, but not limited to, those anti-cancer antibodies listed herein above; small molecule-based cancer therapy, including, but not limited to, the small molecules listed herein above; vaccine/immunotherapy-based cancer therapies, including, but limited to, those listed herein above; steroid therapy; other cancer therapy; or any combination thereof.

“Treatment” in the context of coordinated use of a medicament described herein with one or more other cancer therapies is herein defined as the application or administration of the medicament or of the other cancer therapy to a subject, or application or administration of the medicament or other cancer therapy to an isolated tissue or cell line from a subject, where the subject has a cancer characterized by neoplastic B cell growth, a symptom associated with such a cancer, or a predisposition toward development of such a cancer, where the purpose is to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the cancer, any associated symptoms of the cancer, or the predisposition toward the development of the cancer.

The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

The antagonist anti-CD40 antibodies used in the examples below are CHIR-5.9 and CHIR-12.12. The CHIR-5.9 and CHIR-12.12 anti-CD40 antibodies are human IgG₁ subtype anti-human CD40 monoclonal antibodies (mAbs) generated by immunization of transgenic mice bearing the human IgG₁ heavy chain locus and the human κ light chain locus (XenoMouse[®] technology; Abgenix; Fremont, California). As shown by FACS analysis, CHIR-5.9 and CHIR-12.12 bind specifically to human CD40 and can prevent

CD40 ligand binding. Both mAbs can compete off CD40-ligand pre-bound to cell surface CD40. Both antibodies are strong antagonists and inhibit *in vitro* CD40 ligand-mediated proliferation of normal B cells, as well as cancer cells from NHL and CLL patients. *In vitro*, both antibodies kill cancer cell lines as well as primary cancer cells from NHL patients by ADCC. Dose-dependent anti-tumor activity was seen in a xenograft human lymphoma model. The binding affinity of CHIR-5.9 to human CD40 is 1.2×10^{-8} M and the binding affinity of CHIR-12.12 to human CD40 is 5×10^{-10} M.

Mouse hybridoma line 131.2F8.5.9 (CMCC#12047) and hybridoma line 153.8E2.D10.D6.12.12 (CMCC#12056) have been deposited with the American Type Culture Collection [ATCC; 10801 University Blvd., Manassas, Virginia 20110-2209 (USA)] under Patent Deposit Number PTA-5542 and PTA-5543, respectively.

The following protocols have been used in the examples described below.

ELISA Assay for Immunoglobulin Quantification

The concentrations of human IgM and IgG were estimated by ELISA. 96-well ELISA plates were coated with 2 µg/ml goat anti-human IgG MAb (The Jackson Laboratory, Bar Harbor, Maine) or with 2 µg/ml goat anti-human IgM MAb 4102 (Bio Source International, California) in 0.05 M carbonate buffer (pH 9.6), by incubation for 16 hours at 4°C. Plates were washed 3 times with PBS-0.05 % Tween-20 (PBS-Tween) and saturated with BSA for 1 hour. After 2 washes the plates were incubated for 2 hour at 37°C with different dilutions of the test samples. After 3 washes, bound Ig was detected by incubation for 2 hour at 37°C with 1 µg/ml peroxidase-labeled goat anti-human IgG MAb or goat anti-human IgM Mab. Plates were washed 4 times, and bound peroxidase activity was revealed by the addition of O-phenylenediamine as a substrate. Human IgG or IgM standards (Caltag, Burlingame, California) was used to establish a standard curve for each assay.

Isolation of the Peripheral Blood Mononuclear Cells (PBMC) from Human Peripheral Blood

20 ml of Ficoll-Paque solution (low endotoxin; Pharmacia) was added per 50 ml polystyrene tube, in 3 tubes, 30 minutes before adding the blood. The Ficoll-Paque

solution was warmed up to room temperature. 3 L of bleach in 1:10 dilution was prepared, and used to wash all the tubes and pipettes contacting the blood. The blood was layered on the top of the Ficoll-Paque solution without disturbing the Ficoll layer, at 1.5 ml blood/1 ml of Ficoll-Paque. The tubes were centrifuged at 1700 rpm for 30 minutes at room temperature with the brake on the centrifuge turned off. As much of the top layer (plasma) as possible was removed, minimizing the vacuum in order to avoid removing the second layer of solution. The second layer, which contains the B and T lymphocytes, was collected using a sterile Pasteur pipette, and place in two 50-ml polystyrene tubes. The collection was diluted with 3 x the volume of cold RPMI with no additives, and the tubes were centrifuged at 1000 RPM for 10 minutes. The media was removed by aspiration, and the cells from both 50-ml tubes were resuspended in a total of 10 ml cold RPMI (with additives) and transferred to a 15-ml tube. The cells were counted using the hemacytometer, then centrifuged at 1000 RPM for 10 minutes. The media was removed and the cells resuspended in 4 ml RPMI. This fraction contained the PBMC.

Isolation of the B cells from PBMC

100 µl of Dynabeads (anti-h CD19) were placed in a 5-ml plastic tube. 3 ml of sterile PBS were added to the beads and mixed, and placed in the magnetic holder, then allowed to sit for 2 minutes. The solution was removed using a Pasteur pipette. 3 ml of sterile PBS were added, mixed, and placed in the magnetic holder, then allowed to sit for 2 minutes. This procedure with sterile PBS was repeated one more time for a total of 3 washes. The PBMC was added into the beads and incubated, while mixing, for 30 minutes at 40°C. The tube containing the PBMC and beads was placed into the magnetic holder for 2 minutes, then the solution was transferred to a new 5-ml tube in the magnetic holder. After 2 minutes, the solution was transferred to a new 15-ml tube. This step was repeated four more times, and the solutions of the first four times were collected in the 15-ml tube, then centrifuged at 1000 RPM for 5 minutes. This step produced the pellet for T-cell separation.

100 µl RPMI (with additives) was added to collect the beads, and the solution was transferred into a 0.7-ml tube. 10 µl of Dynal Detacha Beads were added into the

suspension at room temperature, and it was allowed to rotate for 45 minutes. The suspension was transferred into a new 5-ml tube and 3-ml of RPMI (with additives) were added. The tube was placed in the magnetic holder for 2 minutes. The solution was transferred into a new 5-ml tube in the holder for 2 minutes, then to a 15-ml tube. The previous step was repeated three more times, collecting the solution in the 15-ml tube. The 15-ml tube was centrifuged at 1000 RPM for 10 minutes, and the cells resuspended in 10 ml RPMI. The washing step was repeated 2 more times for a total of 3 washes. The cells were counted before the last centrifugation. This step completed the B-cell purification. Cells were stored in 90% FCS and 10% DMSO and frozen at -80°C.

Isolation of the T Cells

The human T cell Enrichment Column (R&D systems, anti-h CD 3 column kit) was prepared using 20 ml of 1 X column wash buffer by mixing 2 ml of 10 X column wash buffer and 18 ml of sterile distilled water. The column was cleaned with 70% ethanol and placed on top of a 15-ml tube. The top cap of the column was removed first to avoid drawing air into the bottom of the column. Next, the bottom cap was removed, and the tip was cleaned with 70% ethanol. The fluid within the column was allowed to drain into the 15-ml tube. A new sterile 15-ml tube was placed under the column after the column buffer had drained to the level of the white filter. The B-cell depleted PBMC fraction was suspended in 1 ml of buffer and added to the top of the column. The cells were allowed to incubate with the column at room temperature for 10 minutes. The T-cells were eluted from the column with 4 aliquots of 2 ml each of 1 X column wash buffer. The collected T-cells were centrifuged at 1000 RPM for 5 minutes. The supernatant was removed and the cells resuspended in 10 ml RPMI. Cells were counted and centrifuged one more time. The supernatant was removed, completing the T-cell purification. Cells were stored in 90% FCS and 10% DMSO and frozen at -80°C.

For the above procedures, the RPMI composition contained 10% FCS (inactivated at 56°C for 45 min), 1 % Pen/Strep (100 u/ml Penicillin, 0.1 µg/ml Streptomycin), 1% Glutamate, 1% sodium puravate, 50 µM 2-ME.

Flow Cytofluorometric Assay

Ramos cells (106 cells/sample) were incubated in 100 μ l primary antibody (10 μ g/ml in PBS-BSA) for 20 min at 4°C. After 3 washes with PBS-BSA or HBSS- BSA, the cells were incubated in 100 μ l FITC-labeled F(ab')₂ fragments of goat anti-(human IgG) antibodies (Caltag) for 20 min at 4°C. After 3 washes with PBS-BSA and 1 wash with PBS, the cells were resuspended in 0.5-ml PBS. Analyses were performed with a FACSCAN V (Becton Dickinson, San Jose, California).

10 *Generation of Hybridoma Clones*

Splenocytes from immunized mice were fused with SP 2/0 or P 3 x 63Ag8.653 murine myeloma cells at a ratio of 10:1 using 50% polyethylene glycol as previously described by de Boer *et al.* (1988) *J. Immunol. Meth.* 113:143. The fused cells were resuspended in complete IMDM medium supplemented with hypoxanthine (0.1 mM), aminopterin (0.01 mM), thymidine (0.016 mM), and 0.5 ng/ml hIL-6 (Genzyme, Cambridge, Massachusetts). The fused cells were then distributed between the wells of 96-well tissue culture plates, so that each well contained 1 growing hybridoma on average.

After 10-14 days the supernatants of the hybridoma populations were screened for specific antibody production. For the screening of specific antibody production by the hybridoma clones, the supernatants from each well were pooled and tested for anti-CD 40 activity specificity by ELISA first. The positives were then used for fluorescent cell staining of EBV-transformed B cells as described for the FACS assay above. Positive hybridoma cells were cloned twice by limiting dilution in IMDM/FBS containing 0.5 ng/ml hIL-6.

Example 1: Production of Anti-CD40 Antibodies

Several fully human, antagonist anti-CD40 monoclonal antibodies of IgG1 isotype were generated. Transgenic mice bearing the human IgG1 heavy chain locus and the human κ chain locus (Abgenix γ -1 XenoMouse[®] technology (Abgenix; Fremont, California)) were used to generate these antibodies. SF9 insect cells expressing CD40

extracellular domain were used as immunogen. A total of 31 mice spleens were fused with the mouse myeloma SP2/0 cells to generate 895 antibodies that recognize recombinant CD40 in ELISA (Tables 1A and 1B). On average approximately 10% of hybridomas produced using Abgenix XenoMouse® technology may contain mouse lambda light chain instead of human kappa chain. The antibodies containing mouse light lambda chain were selected out. A subset of 260 antibodies that also showed binding to cell-surface CD40 were selected for further analysis. Stable hybridomas selected during a series of subcloning procedures were used for further characterization in binding and functional assays.

Table 1A. A Typical Fusion

Fusion #	anti-CD40 titer		# of wells screened	# of ELISA+	# of cell surface+
	(1:100K)	Fusion Efficiency			
153	3	100%	960	123	33
154	4.67	15%	140	0	0
155	6	~40%	960	3	3
156	3.17	~25%	220	1	0
157	4.67	90%	960	32	6
158	4.4	90%	960	23	8
159	1.17	100%	960	108	18
160	1.78	90%	960	30	5
Total			6120	320	73

Table 1B. Summary of Four Sets of Fusions

# of mice	ELISA-positive hybridomas	Cell surface positive Hybridomas
31	895	260

Table 2. Summary of initial set of data with anti-CD40 IgG1 antibodies

Mother Hybridoma	Hybridoma clones	cell surface binding	Antagonist	ADCC	CDC	CMCC#	V-region DNA sequence
	131.2F5.8.5.1	+++	++	ND	ND	ND	
131.2F5	131.2F5.8.5.9	+++	+++	++	-	12047	Yes
	131.2F5.8.5.11	+++	+++	++	-	12055	Yes
	153.3C5D8D7.8.4.7.1	++	ND	ND	ND	ND	
153.3C5	153.3C5D8D7.8.4.7.8	++	ND	ND	ND	ND	
	153.3C5D8D7.8.4.7.11	+++	+++	+	ND	ND	
	153.1D2.9.1	+++	ND	ND	ND	12067	
153.1D2	153.1D2.9.8	+++	+++	++	-	12057	
	153.1D2.9.12	+++	ND	ND	ND	12068	
	158.6F3.5.1	+++	+++	++	-	12054	Yes
158.6F3	158.6F3.5.7	+++	ND	ND	ND	12061	
	158.6F3.5.10	+++	ND	ND	ND	12062	
	153.8E2D10D6.12.7	+++	ND	ND	ND	12075	
153.8E2	153.8E2D10D6.12.9	+++	ND	ND	ND	12063	
	153.8E2D10D6.12.12	+++	+++	++++	-	12056	Yes
	155.2C2E9F12.2.10.4	+++	+/-	ND	ND	12064	
155.2C2	155.2C2E9F12.2.10.5	+++	ND	ND	ND	12065	
	155.2C2E9F12.2.10.6	+/-	ND	ND	ND	12066	
	166.5E6G12.1	+++	ND	ND	ND	12069	
166.5E6	166.5E6G12.3	+++	ND	ND	ND	12070	
	166.5E6G12.4	+++	+	ND	ND	12071	
177.8C10	177.8C10B3H9	+++	++	ND	ND	ND	
	183.4B3E11.6.1.5	++	ND	ND	ND	ND	
183.4B3	183.4B3E11.6.1.9	++	ND	ND	ND	ND	
	183.4B3E11.6.1.10	+++	++	ND	ND	ND	
	183.2G5D2.8.7	+++	+/-	ND	ND	ND	
183.2G5	183.2G5D2.8.8	+++	ND	ND	ND	ND	
	183.2G5D2.8.9	+++	ND	ND	ND	ND	
	184.6C11D3.2	++	ND	ND	ND	12078	
184.6C11	184.6C11D3.3	++	ND	ND	ND	12080	
	184.6C11D3.6	+/-	+/-	ND	ND	12079	
	185.3E4F12.5.6	+++	ND	ND	ND	12072	
185.3E4	185.3E4F12.5.11	+++	ND	ND	ND	12073	
	185.3E4F12.5.12	+++	+	ND	ND	12074	
	185.1A9E9.6.1	+	ND	ND	ND	ND	
185.1A9	185.1A9E9.6.6	+++	+++	+	ND	ND	
	185.9F11E10.3B5.1	+++	ND	ND	ND	ND	
185.9F11	185.9F11E10.3B5.8	+++	ND	ND	ND	ND	
	185.9F11E10.3B5.12	+++	+++	ND	ND	ND	

Clones from 7 mother hybridomas were identified to have antagonist activity. Based on their relative antagonist potency and ADCC activities, two hybridoma clones were selected. Their names are: 131.2F8.5.9 (5.9) and 153.8E2.D10.D6.12.12 (12.12).

Clones from 7 other hybridomas were identified as having antagonist activity (Table 2 above). Based on their relative antagonistic potency and ADCC activities, two hybridoma clones were selected for further evaluation. They are named 131.2F8.5.9 (5.9) and 153.8E2.D10.D6.12.12 (12.12). The binding profile of these two antibodies to CD40+ lymphoma cell line is shown as a flow cytometric histogram in Figure 1.

Example 2: Polynucleotide and Amino Acid Sequences of Human Anti-CD40

Antibodies

The cDNAs encoding the variable regions of the candidate antibodies were amplified by PCR, cloned, and sequenced. The amino acid sequences for the light chain and heavy chain of the CHIR-12.12 antibody are set forth in Figures 9A and 9B, respectively. See also SEQ ID NO:2 (light chain for mAb CHIR-12.12) and SEQ ID NO:4 (heavy chain for mAb CHIR-12.12). A variant of the heavy chain for mAb CHIR-12.12 is shown in Figure 9B (see also SEQ ID NO:5), which differs from SEQ ID NO:4 in having a serine residue substituted for the alanine residue at position 153 of SEQ ID NO:4. The nucleotide sequences encoding the light chain and heavy chain of the CHIR-12.12 antibody are set forth in Figures 10A and 10B, respectively. See also SEQ ID NO:1 (coding sequence for light chain for mAb CHIR-12.12) and SEQ ID NO:3 (coding sequence for heavy chain for mAb CHIR-12.12). The amino acid sequences for the light chain and heavy chain of the CHIR-5.9 antibody are set forth in Figures 11A and 11B, respectively. See also SEQ ID NO:6 (light chain for mAb CHIR-5.9) and SEQ ID NO:7 (heavy chain for mAb CHIR-5.9). A variant of the heavy chain for mAb CHIR-5.9 is shown in Figure 11B (see also SEQ ID NO:8), which differs from SEQ ID NO:7 in having a serine residue substituted for the alanine residue at position 158 of SEQ ID NO:7.

As expected for antibodies derived from independent hybridomas, there is substantial variation in the nucleotide sequences in the complementarity determining regions (CDRs). The diversity in the CDR3 region of V_H is believed to most significantly determine antibody specificity.

Example 3: Effect of CHIR-5.9 and CHIR-12.12 on the CD40/CD40L Interaction *In vitro*

The candidate antibodies CHIR-5.9 and CHIR-12.12 prevent the binding of CD40 ligand to cell surface CD40 and displace the pre-bound CD40 ligand. Antibodies CHIR-5.9 and CHIR-12.12 were tested for their ability to prevent CD40-ligand binding to CD40 on the surface of a lymphoma cell line (Ramos). Binding of both antibodies (unlabeled) prevented the subsequent binding of PE-CD40 ligand as measured by flow cytometric assays (Figure 2A). In a second set of assays the two antibodies were tested for their ability to replace CD40 ligand pre-bound to cell surface CD40. Both antibodies were effective for competing out pre-bound CD40 ligand, with CHIR-5.9 being slightly more effective than CHIR-12.12 (Figure 2B).

Example 4: CHIR-5.9 and CHIR-12.12 Bind to a Different Epitope on CD40 Than 15B8

The candidate monoclonal antibodies CHIR-5.9 and CHIR-12.12 compete with each other for binding to CD40 but not with 15B8, an IgG₂ anti-CD40 mAb (see International Publication No. WO 02/28904). Antibody competition binding studies using Biacore were designed using CM5 biosensor chips with protein A immobilized via amine coupling, which was used to capture either anti-CD40, CHIR-12.12, or 15B8. Normal association/dissociation binding curves are observed with varying concentrations of CD40-his (data not shown). For competition studies, either CHIR-12.12 or 15B8 were captured onto the protein A surface. Subsequently a CD40-his /CHIR-5.9 Fab complex (100 nM CD40:1 μ M CHIR-5.9 Fab), at varying concentrations, was flowed across the modified surface. In the case of CHIR-12.12, there was no association of the complex observed, indicating CHIR-5.9 blocks binding of CHIR-12.12 to CD40-his. For 15B8, association of the Fab CHIR-5.9 complex was observed indicating CHIR-5.9 does not block binding of 15B8 to CD40 binding site. However, the off rate of the complex dramatically increased (data not shown).

It has also been determined that 15B8 and CHIR-12.12 do not compete for CD40-his binding. This experiment was set up by capturing CHIR-12.12 on the protein A biosensor chip, blocking residual protein A sites with control hIgG₁, binding CD40-his

and then flowing 15B8 over the modified surface. 15B8 did bind under these conditions indicating CHIR-12.12 does not block 15B8 from binding to CD40.

Example 5: Binding Properties of Selected Hybridomas

5 Protein A was immobilized onto CM5 biosensor chips via amine coupling. Human anti-CD40 monoclonal antibodies, at 1.5 µg/ml, were captured onto the modified biosensor surface for 1.5 minutes at 10 µl/min. Recombinant soluble CD40-his was
10 flowed over the biosensor surface at varying concentrations. Antibody and antigen were diluted in 0.01 M HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005% Surfactant P20 (HBS-EP). Kinetic and affinity constants were determined using the Biaevaluation software with a 1:1 interaction model/global fit.

As shown in Table 3 below, there is 121-fold difference in the off rate of CHIR-5.9 and CHIR-12.12 resulting in 24-fold higher affinity for CHIR-12.12.

<u>Antibody</u>	<u>ka (M⁻¹ sec⁻¹)</u>	<u>kd (sec⁻¹)</u>	<u>KD (nM)</u>
Anti-CD40, CHIR-5.9	$(12.35 \pm 0.64) \times 10^5$	$(15.0 \pm 1.3) \times 10^{-3}$	12.15 ± 0.35
Anti-CD40, CHIR-12.12	$(2.41 \pm 0.13) \times 10^5$	$(1.24 \pm 0.06) \times 10^{-4}$	0.51 ± 0.02

Example 6: CHIR-5.9 and CHIR-12.12 are Potent Antagonists for CD40-Mediated Proliferation of Human Lymphocytes from Normal Subjects

20 Engagement of CD40 by CD40 ligand induces proliferation of human B cells. An antagonist anti-CD40 antibody is expected to inhibit this proliferation. Two candidate antibodies (CHIR-5.9 and CHIR-12.12) were tested for their ability to inhibit CD40 ligand-induced proliferation of PBMC from normal human subjects. Formaldehyde-fixed CHO cells transfectant-expressing CD40 ligand (CD40L) were used as a source of CD40

ligand. Human PBMC were cultured for 4 days with the formaldehyde-fixed CHO cells expressing CD40-ligand in the presence of varying concentrations of anti-CD40 mAb CHIR-5.9 or CHIR-12.12. The proliferation was measured by tritiated-thymidine incorporation. Cells were pulsed with tritiated-labeled thymidine at 37°C for 14-18
5 hours.

Both antibodies were found to be very effective for inhibiting CD40 ligand-induced proliferation of human PBMC (Table 4A, mAb CHIR-5.9, Table 4B, mAb CHIR-12.12). The experiment was performed with multiple donors of PBMC (n=12 for CHIR-5.9 and n=2 for CHIR-12.12) to ensure that the observed inhibition was not a
10 peculiarity of cells from a single donor. Follow-up assessments with 4 additional donors of PBMC were carried out for mAb CHIR-12.12 with similar trends observed. A wide range of antibody concentrations (0.01 µg/ml to 100 µg/ml) was used in these assays. Nearly complete inhibition of CD40 ligand-induced proliferation could be achieved at 0.1 µg/ml concentration of antibodies in most cases. Antibody concentration (pM) to inhibit
15 50% of CD-40 ligand-induced lymphocyte proliferation (IC50) for lymphocytes from 6 donors yielded an average IC50 (pM) of 47 (SD=21; donor 1, 24; donor 2, 66; donor 3, 45; donor 4, 84; donor 5, 30; donor 6, 35), which compares favorably with the average IC50 (pM) of 49.65 shown in Table 4B. Based on the current data set, both candidate antibodies seem similar in their potency for inhibition of CD40 ligand-induced
20 proliferation of normal PBMC.

Table 4A. Effect of mAb CHIR-5.9 on CD40L-induced PBMC proliferation.

Exp#		CHO-CD40L	PBMC+	Abs Conc (μ g/ml)	hulgG1		CHIR-5.9		
		alone	CHO-CD40L		CPM	% of inhibition	CPM	% of inhibition	
PBMC-010		1851	121	4436	1	5080	-26	2622	74
		1851	121	4436	0.25	5498	-43	2907	62
		1851	121	4436	0.0625	6029	-65	2619	74
		1851	121	4436	0.0156	5814	-56	1199	131
PBMC-011	Donor#1	2162	178	8222	10	13137	-84	2252	101
		2162	178	8222	1	11785	-61	1438	115
		2162	178	8222	0.1	10758	-43	1249	119
		2162	178	8222	0.01	11322	-53	4705	60
	Donor#2	2216	294	7873	10	16679	-164	2362	103
		2216	294	7873	1	14148	-117	1202	124
		2216	294	7873	0.1	12422	-85	756	133
		2216	294	7873	0.01	13870	-112	6606	24
	Donor#3	2396	241	11021	10	11641	-7	2631	100
		2396	241	11021	1	13528	-30	1450	114
		2396	241	11021	0.1	12176	-14	990	120
		2396	241	11021	0.01	11895	-10	5357	68
	Donor#4	4552	133	15301	10	22098	-64	3768	109
		4552	133	15301	1	19448	-39	2040	125
		4552	133	15301	0.1	18398	-29	1728	128
		4552	133	15301	0.01	22767	-70	9481	55
PBMC-012		777	117	6041	10	7327	-25	2150	76
		777	117	6041	1	6212	-3	1550	87
		777	117	6041	0.1	7006	-19	828	101
		777	117	6041	0.01	7524	-29	1213	94
PBMC-014		1857	73	7889	100	9399	-25	3379	76
		1857	73	7889	20	8120	-4	3870	67
		1857	73	7889	4	8368	-8	2552	90
		1857	73	7889	0.8	9564	-28	1725	103
PBMC-015	Donor#1	3203	127	10485	100	15425	-69	1497	126
		3203	127	10485	20	11497	-14	1611	124
		3203	127	10485	4	11641	-16	1359	128
		3203	127	10485	0.8	12807	-32	1490	126
	Donor#2	3680	175	15145	100	21432	-56	1792	118
		3680	175	15145	20	16998	-16	1779	118
		3680	175	15145	4	17729	-23	1965	117
		3680	175	15145	0.8	17245	-19	2217	115
	Donor#3	2734	152	19775	100	22967	-19	1664	107
		2734	152	19775	20	21224	-9	1848	106
		2734	152	19775	4	20658	-5	1534	108
		2734	152	19775	0.8	18923	5	1262	110
PBMC-016		1118	36	13531	0.1	10928	21	745	103
		1118	36	13531	0.05	11467	17	962	102
		1118	36	13531	0.01	11942	13	3013	85
PBMC-017		962	75	12510	1	13597	-9	258	107

Table 4A. Effect of mAb CHIR-5.9 on CD40L-induced PBMC proliferation (cont.).

Average % inhibition of human PBMC at 100 μ g/ml	-42	107
Average % inhibition of human PBMC at 10 μ g/ml	-69	98
Average % inhibition of human PBMC at 1 μ g/ml	-41	107
Average % inhibition of human PBMC at 0.1 μ g/ml	-28	117
Average % inhibition of human PBMC at 0.01 μ g/ml	-44	64
Average % inhibition of human PBMC	-35	101

% of inhibition:

$100 - (\text{CPM with Abs-PBMC alone} - \text{CHO-CD40L alone}) / (\text{CPM of PBMC} + \text{CHO-CD40L-PBMC alone} - \text{CHO-CD40L alone}) * 100\%$

Table 4B. Effect of mAb CHIR-12.12 on CD40L-induced PBMC proliferation.

Exp#		PBMC alone	CHO-CD40L alone	PBMC+ CHO-CD40L	Abs Conc (μ g/ml)	HulgG1		CHIR-12.12		
						CPM	% of inhibition	CPM	% of inhibition	IC50(nM)
PBMC-025	Donor#1	4051	32	42292	0.1	33354	23	440	110	
		4051	32	42292	0.01	37129	14	8696	88	
		4051	32	42292	0.001	40271	5	32875	25	
		4051	32	42292	0.0001	40034	6	37261	13	24.22
	Donor#2	2260	31	14987	0.1	15767	-6	365	115	
		2260	31	14987	0.01	17134	-17	6734	65	
		2260	31	14987	0.001	20142	-41	16183	-9	
		2260	31	14987	0.0001	17847	-23	16187	-9	65.96
	Donor#1	2039	35	19071	0.1	17136	11	624	109	
		2039	35	19071	0.01	16445	15	6455	74	
		2039	35	19071	0.001	16195	17	17833	7	
		2039	35	19071	0.0001	18192	5	17924	7	45
PBMC-026	Donor#2	2016	64	17834	0.1	17181	4	2078	100	
		2016	64	17834	0.01	16757	7	10946	44	
		2016	64	17834	0.001	18613	-5	17924	-1	
		2016	64	17834	0.0001	17169	4	18569	-5	84
PBMC-028	Donor#1	4288	45	22547	1	18204	24	2098	112	
		4288	45	22547	0.1	20679	10	1827	114	
		4288	45	22547	0.01	22799	-1	6520	88	
		4288	45	22547	0.001	23547	-5	22327	1	
		4288	45	22547	0.0001	24778	-12	24124	-9	30.07
	Donor#2	2148	58	54894	1	48545	12	5199	94	
		2148	58	54894	0.1	45708	17	5091	95	
		2148	58	54894	0.01	51741	6	18890	68	
		2148	58	54894	0.001	52421	5	50978	7	
		2148	58	54894	0.0001	54778	0	52581	4	34.68
PBMC-029	Donor#1	609	69	10054	0.1	11027	-10	2098	85	
		609	69	10054	0.01	10037	0	1827	88	
		609	69	10054	0.001	10222	-2	6520	38	
		609	69	10054	0.0001	11267	-13	22327	-131	28.06
	Donor#2	7737	57	23132	0.1	21254	12	2536	134	
		7737	57	23132	0.01	21726	9	10249	84	
		7737	57	23132	0.001	22579	4	23380	-2	
		7737	57	23132	0.0001	22491	4	23183	0	55.35
PBMC-030	Donor#1	2739	47	53426	0.1	60116	-13	2132	101	
		2739	47	53426	0.01	56411	-6	14297	77	
		2739	47	53426	0.001	59167	-11	55868	-5	
		2739	47	53426	0.0001	59290	-12	60865	-15	35.52
	Donor#2	4310	50	53781	0.1	52881	2	3208	102	
		4310	50	53781	0.01	51741	4	30716	47	
		4310	50	53781	0.001	53072	1	53628	0	
		4310	50	53781	0.0001	58045	-9	54343	-1	102.88
PBMC-032	Donor#1	2458	42	14058	0.1	16579	-22	636	116	40.36
		2458	42	14058	0.01	19250	-45	3358	93	

2458	42	14058	0.001	19852	-50	20639	-57
2458	42	14058	0.0001	19161	-44	18907	-42
Average % inhibition of human PBMC at 0.1 μ g/ml					3		107
Average % inhibition of human PBMC at 0.01 μ g/ml					-1		74
Average % inhibition of human PBMC at 0.001 μ g/ml					-7		0
Average % inhibition of human PBMC at 0.0001 μ g/ml					-8		-17 49.65
% of inhibition: 100-(CPM with Abs-PBMC alone-CHO-CD40L alone)/(CPM of PBMC+CHO-CD40L-PBMC alone-CHO-CD40L alone)*100%							

In addition to B cells, human PBMC also contain natural killer cells that can mediate antibody dependent cytotoxicity (ADCC). To clarify the mechanism of antibody-mediated inhibition of proliferation, assays were performed with B cells purified from human PBMC. Similar to results obtained with PBMC, both antibodies potently inhibited the CD40 ligand-induced proliferation of purified B cells (Table 5, n=3). These data demonstrate that the antagonist activity of the candidate antibodies, and not the mechanism of ADCC, is the cause of proliferation inhibition in these assays.

Table 5. Effect of anti-CD40 antibodies on CD40 ligand-induced proliferation of purified human B cells

Exp#	Donor #	CPM			Abs Conc(μg/ml)	HuIgG1		CHIR-5.9		CHIR-12.12	
		B cells	CHO- CD40L	B cells + CHO- CD40L		CPM	% inhibition	CPM	% inhibition	CPM	% inhibition
B cell-004	1	418	89	3132	100	429	103	271	109	152	114
		418	89	3132	20	3193	-2	316	107	222	111
		418	89	3132	4	3175	-2	144	114	235	110
		418	89	3132	0.8	6334	-122	245	110	63	117
	2	81	73	27240	100	28311	-4	85	100	77	100
		81	73	27240	20	24707	9	65	100	94	100
		81	73	27240	4	23081	15	108	100	68	100
		81	73	27240	0.8	26252	4	87	100	77	100
B cell-005	3	267	75	24552	1	25910	-6	291	100	102	101
		267	75	24552	0.1	28447	-16	259	100	108	101
		267	75	24552	0.01	26706	-9	2957	89	4922	81
Average % inhibition							-3	103		103	

% of inhibition: 100-(CPM with Abs-B cells alone-CHO-CD40L alone)/(CPM of B cell with CHO-CD40L-B cells alone-CHO-CD40L alone)*100%

Example 7: CHIR-5.9 and CHIR-12.12 Do Not Induce Strong Proliferation of Human B Cells from Normal Subjects

CD40 ligand activates normal B cells and B-cell lymphoma cells to proliferate. Binding of some anti-CD40 antibodies (agonist) can provide a similar stimulatory signal

for the proliferation of normal and cancer B cells. Antibodies with strong B cell stimulatory activity are not suitable candidates for therapeutic treatment of B cell lymphomas. The two candidate antibodies were tested for their ability to induce proliferation of B cells from normal volunteer donors. The B cells purified by Ficoll-Hypaque Plus gradient centrifugation from normal donor PBMC were cultured in 96-well plates with varying concentrations of candidate antibodies (range of 0.001 to 100 μ g/ml) for a total of 4 days. In the positive control group, PBMC were cultured with formaldehyde-fixed CHO cells expressing CD40-ligand. The B cell proliferation was measured by incorporation of tritiated-labeled thymidine at 37°C for 14-18 hours. While the CD40 ligand presented on CHO cells induced vigorous proliferation of B cells resulting in an average stimulation index (SI) of 145, the candidate antibodies induced only a weak proliferation with a stimulation index of 2.89 and 5.08 for CHIR-12.12 and CHIR-5.9, respectively (n=3) (Table 6).

Table 6. Proliferation of B cells purified from normal human subjects in response to candidate anti-CD40 mAbs

Exp#	Donor#	B cells	Bcells+CHO-CD40L		Abs conc (µg/ml)	B cells+huIgG1		B cells+CHIR-5.9		B cells+CHIR-12.12		
		CPM	CPM	S. index(1)		CPM	S. index(2)	CPM	S. index (2)	CPM	S. index(2)	
B cell-004 Frozen	1	418	3132	7.49	100	498	1.19	401	0.96	458	1.10	
			418		3132	20	245	0.59	232	0.56	370	0.89
			418		3132	4	241	0.58	232	0.56	211	0.50
			418		3132	0.8	376	0.90	298	0.71	230	0.55
Frozen	2	81	27240	336.30	100	34	0.42	454	5.60	122	1.51	
			81		27240	20	48	0.59	706	8.72	255	3.15
			81		27240	4	41	0.51	567	7.00	367	4.53
			81		27240	0.8	34	0.42	736	9.09	408	5.04
B cell-005	3	267	24552	91.96	1	691	2.59	2101	7.87	1223	4.58	
			267		24552	0.1	686	2.57	2267	8.49	1557	5.83
			267		24552	0.01	808	3.03	2203	8.25	1027	3.85
			267		24552	0.001	567	2.12	846	3.17	826	3.09
Average Stimulation Index (SI)				145.25		1.29		5.08		2.88		
S. index(1): = CPM (B cells+CHO-CD40L)/CPM (B cells alone)												
S. index(2): = CPM (B cells+Abs)/CPM (PBMC alone)												

In addition to B cells, human PBMC contain cell types that bear Fc receptors (FcR) for IgG1 molecules that can provide cross linking of anti-CD40 antibodies bound to CD40 on B cells. This cross-linking could potentially enhance stimulatory activity of anti-CD40 antibodies. To confirm the lack of B cell stimulatory activity of CHIR-5.9 and

CHIR-12.12 antibodies in the presence of cross-linking cells, proliferation experiments were performed with total PBMC containing B cells as well as FcR+ cells. Data from these experiments (Table 7A, mAb CHIR-5.9; Table 7B, mAb CHIR-12.12) confirm that these candidate antibodies even in the presence of FcR-bearing cells in general do not stimulate B cells to proliferate over background proliferation induced by control human IgG1 (n=10). The CD40 ligand induced an average stimulation index (SI) SI of 7.41. The average SI with candidate antibodies were 0.55 and 1.05 for CHIR-12.12 and CHIR-5.9, respectively. Only one of the 10 donor PBMC tested showed some stimulatory response to CHIR-5.9 antibody (donor #2 in Table 7). The lack of stimulatory activity by candidate mAbs was further confirmed by measuring the PBMC proliferation in response to candidate anti-CD40 antibodies immobilized on the plastic

Table 7A. Proliferation of PBMC from normal human subjects in response to mAb CHIR-5.9.

Exp#		PBM C	PBMC+CHO-CD40L		Abs conc (μ g/ml)	PBMC+hulgG1		PBMC+CHIR-5.9	
			CPM	index(1)		CPM	index(2)	CPM	index(2)
PBMC-010			1417 5279	3.73	1	1218	0.86	597 3	4.22
			1417 5279		0.25	1712	1.21	481 5	3.40
			1417 5279		0.062			364 2	2.57
			1417 5279		5	1449	1.02	324 2	2.29
			1417 5279		0.015		0.84		
PBMC-011	Donor#1		2138 8247	3.86	10	3047	1.43	317 7	1.49
			2138 8247		1	2726	1.28	361 7	1.69
			2138 8247		0.1	2026	0.95	201 1	0.94
			2138 8247		0.01	2424	1.13	186 0	0.87
	Donor#2		1156					452	
			2374 1	4.87	10	4966	2.09	3	1.91
			1156					244	
			2374 1		1	2544	1.07	5	1.03
			1156					146	
	Donor#3		2374 1		0.1	2177	0.92	2	0.62
			1156					189	
			2374 1		0.01	4672	1.97	6	0.80
			3229 7956	2.46	10	5035	1.56	211 9	0.66
			3229 7956		1	2826	0.88	109 9	0.34
			3229 7956		0.1	2277	0.71	105 2	0.33
			3229 7956		0.01	3078	0.95	189 9	0.59
	Donor#4		1431					517	
			4198 4	3.41	10	5012	1.19	6	1.23
			1431					470	
			4198 4		1	3592	0.86	2	1.12
			1431					431	
PBMC-014			4198 4		0.1	5298	1.26	9	1.03
			1431					540	
			4198 4		0.01	5758	1.37	0	1.29
			2350 8787	3.74	100	2722	1.16	247 1	1.05
			2350 8787		20	2315	0.99	244 7	1.04
PBMC-015	Donor#1		2350 8787		4	2160	0.92	165 9	0.71
			2350 8787		0.8	2328	0.99	167 1	0.71
			1293					168	
			3284 6	3.94	100	3598	1.10	2	0.51
			1293					156	
	Donor#2		3284 6		20	2751	0.84	2	0.48
			1293					110	
			3284 6		4	3135	0.95	5	0.34
			1293					141	
			3284 6		0.8	4027	1.23	9	0.43
PBMC-015	Donor#2		1912					510	
			6099 1	3.14	100	2999	0.49	4	0.84

	1912					391	
	6099 1		20	4025	0.66	7	0.64
	1912					334	
	6099 1		4	4496	0.74	1	0.55
	1912					413	
	6099 1		0.8	3834	0.63	9	0.68
Donor#3	1982					120	
	2479 6	8.00	100	3564	1.44	4	0.49
	1982					782	
	2479 6		20	1874	0.76		0.32
	1982					634	
	2479 6		4	1779	0.72		0.26
	1982					937	
	2479 6		0.8	2274	0.92		0.38
PBMC-016	1578					103	
	1148 9	13.75	0.1	1255	1.09	6	0.90
	1578					871	
	1148 9		0.05	1284	1.12		0.76
	1578					952	
	1148 9		0.01	1446	1.26		0.83
Average SI of PBMC		5.09			1.06		1.03

index(1): =(PBMC+CHO-CD40L)/PBMC

index(2): =(PBMC+Abs)/PBMC

Table 7B. Proliferation of PBMC from normal human subjects in response to mAb CHIR-12.12.

Exp#		PBMC	PBMC+CHO-CD40L		Abs conc (μ g/ml)	PBMC+hulgG1		PBMC+CHIR-12.12	
			CPM	index(1)		CPM	index(2)	CPM	index(2)
PBMC-025	Donor#1	4051	42292	10.44	0.1	2909	0.72	2451	0.61
		4051	42292		0.01	4725	1.17	8924	2.20
		4051	42292		0.001	8080	1.99	8782	2.17
		4051	42292		0.0001	4351	1.07	4342	1.07
	Donor#2	2260	14987	6.63	0.1	2538	1.12	6741	2.98
		2260	14987		0.01	3524	1.56	8921	3.95
		2260	14987		0.001	3159	1.40	4484	1.98
		2260	14987		0.0001	2801	1.24	2533	1.12
PBMC-026	Donor#1	2085	18313	8.78	0.1	1386	0.66	2761	1.32
		2085	18313		0.01	2871	1.38	3162	1.52
		2085	18313		0.001	2602	1.25	3233	1.55
		2085	18313		0.0001	1709	0.82	1766	0.85
	Donor#2	676	18054	26.71	0.1	660	0.98	2229	3.30
		676	18054		0.01	2864	4.24	1238	1.83
		676	18054		0.001	693	1.03	1507	2.23
		676	18054		0.0001	984	1.46	811	1.20
PBMC-027	Donor#1	2742	13028	4.75	0.1	4725	1.72	2795	1.02
		2742	13028		0.01	4575	1.67	5353	1.95
		2742	13028		0.001	3218	1.17	3501	1.28
		2742	13028		0.0001	5107	1.86	4272	1.56
	Donor#2	1338	11901	8.89	0.1	1633	1.22	1943	1.45
		1338	11901		0.01	1520	1.14	5132	3.84
		1338	11901		0.001	1517	1.13	2067	1.54
		1338	11901		0.0001	1047	0.78	2076	1.55
PBMC-028	Donor#1	4288	22547	5.26	0.1	3686	0.86	2525	0.59
		4288	22547		0.01	3113	0.73	2047	0.48

		4288	22547		0.001	4414	1.03	3515	0.82
		4288	22547		0.0001	2452	0.57	4189	0.98
	Donor#2	2148	54894	25.56	0.1	9127	4.25	5574	2.59
		2148	54894		0.01	4566	2.13	6515	3.03
		2148	54894		0.001	5285	2.46	5919	2.76
		2148	54894		0.0001	4667	2.17	4298	2.00
PBMC-029	Donor#1	609	10054	16.51	0.1	359	0.59	363	0.60
		609	10054		0.01	473	0.78	956	1.57
		609	10054		0.001	461	0.76	1159	1.90
		609	10054		0.0001	625	1.03	558	0.92
	Donor#2	7737	23132	2.99	0.1	4940	0.64	3493	0.45
		7737	23132		0.01	6041	0.78	3644	0.47
		7737	23132		0.001	5098	0.66	5232	0.68
		7737	23132		0.0001	5135	0.66	5241	0.68
PBMC-030	Donor#1	4164	57205	13.74	10	2713	0.65	1046	0.25
		4164	57205		1	3627	0.87	1576	0.38
		4164	57205		0.1	4590	1.10	1512	0.36
		4164	57205		0.01	4384	1.05	2711	0.65
	Donor#2	3324	53865	16.20	10	6376	1.92	4731	1.42
		3324	53865		1	4720	1.42	5219	1.57
		3324	53865		0.1	3880	1.17	5869	1.77
		3324	53865		0.01	3863	1.16	5657	1.70
PBMC-032	Donor#1	1808	15271	8.45	10	2349	1.30	4790	2.65
		1808	15271		1	3820	2.11	5203	2.88
		1808	15271		0.1	2098	1.16	6332	3.50
		1808	15271		0.01	1789	0.99	5005	2.77
Average SI of PBMC				11.92			1.30		1.62

index(1): =CPM of (PBMC+CHO-CD40L)/CPM of PBMC

index(2): =CPM of (PBMC+Abs)/CPM of PBMC

surface of the culture wells (n=2). The average SI with CD40 ligand, CHIR-12.12, and CHIR-5.9 stimulation were 22, 0.67, and 1.2, respectively (Table 8). Taken together these data show that the candidate antiCD40 antibodies do not possess strong B cell stimulatory properties.

Table 8. Proliferation of PBMC from normal human subjects in response to immobilized anti-CD40 antibodies

Exp#	PBMC	PBMC+CHO-CD40L		Abs conc (ug/ml)	PBMC+huIgG1		PBMC+CHIR-5.9		PBMC+CHIR-12.12	
	CPM	CPM	S.index (1)		CPM	S. index(2)	CPM	S.index (2)	CPM	S.index (2)
PBMC-012	225	6808	30.26	10	279	1.24	734	3.26	200	0.89
	225	6808		1	175	0.78	178	0.79	161	0.72
	225	6808		0.1	156	0.69	226	1.00	249	1.11
	225	6808		0.01	293	1.30	232	1.03	254	1.13
Immobilize-004	857	11701	13.65	1000	479	0.56	1428	1.67	384	0.45
	857	11701		100	543	0.63	839	0.98	265	0.31
	857	11701		10	487	0.57	411	0.48	262	0.31
	857	11701		1	632	0.74	372	0.43	376	0.44

Average Stimulation index	21.96	0.81	1.21	0.67
S. index (1): = CPM (PBMC+CHO-CD40L)/CPM (PBMC)				
S. index (2): = CPM (PBMC+mAbs)/CPM (PBMC)				

Example 8: CHIR-5.9 and CHIR-12.12 Are Able to Kill CD40-Bearing Target Cells by ADCC

The candidate antibodies can kill CD40-bearing target cells (lymphoma lines) by the mechanism of ADCC. Both CHIR-5.9 and CHIR-12.12 are fully human antibodies of IgG1 isotype and are expected to have the ability to induce the killing of target cells by the mechanism of ADCC. They were tested for their ability to kill cancer cell lines in *in vitro* assays. Two human lymphoma cell lines (Ramos and Daudi) were selected as target cells for these assays. PBMC or enriched NK cells from 8 normal volunteer donors were used as effector cells in these assays. A more potent ADCC response was observed with CHIR-12.12 compared with CHIR-5.9 against both the lymphoma cancer cell line target cells. Lymphoma cell lines also express CD20, the target antigen for rituximab (Rituxan®), which allowed for comparison of the ADCC activity of these two candidate mAbs with rituximab ADCC activity. For lymphoma cell line target, an average specific lysis of 35%, 59%, and 47% was observed for CHIR-5.9, CHIR-12.12, and rituximab respectively when used at 1 µg/ml concentration (Table 9). The two antibodies did not show much activity in complement dependent cytotoxicity (CDC) assays.

Table 9. Anti-CD40 mAb dependent killing of lymphoma cell lines by ADCC.

Anti-CD40 mAb dependent killing of lymphoma cell lines by ADCC										
Exp#	Effector cell	E:T ratio	Target cells : Human lymphoma cell line (Ramos or Daudi)							
			% lysis IgG1	Abs conc(μ g/ml)	CHIR-5.9		CHIR-12.12		Rituxan	
					% lysis	% lysis - % lysis IgG1	% lysis	% lysis - % lysis IgG1	% lysis	% lysis - % lysis IgG1
ADCC-005	huNK	3	17.05	5	30.75	13.70	65.22	48.17	ND	ND
Alarmor Blue	huNK	3	40.81	5	58.62	17.81	87.87	47.06	ND	ND
ADCC-006	huNK	2	-3.09	10	3.50	6.59	43.71	46.8	34.82	37.91
Alarmor Blue			-8.62	1	-10.10	-1.48	45.13	53.75	37.07	45.69
			-11	0.1	-14.80	-3.80	39.82	50.82	33.61	44.61
			-4.54	0.01	2.53	7.07	50.07	54.61	28.49	33.03
51Cr	huNK	5	1.5	10	32.09	30.59	47.24	45.742	ND	ND
			2.4	1	18.01	15.61	37.42	35.022	ND	ND
			2.5	0.1	14.67	12.17	37.63	35.131	ND	ND
ADCC-009	huNK	10	2.32	5	66.20	63.88	97.70	95.38	86.2	83.88
Calcein AM			0.48	1	67.20	66.72	123.00	122.52	88.2	87.72
			-1.43	0.2	78.40	79.83	118.00	119.43	88.8	90.23
			3.39	0.04	69.10	65.71	109.00	105.61	84.9	81.51
ADCC-011	huNK	8	3.18	1	15.36	12.19	51.59	48.42	22.44	19.27
Calcein AM			4.58	0.01	7.39	2.81	46.80	42.22	14.68	10.10
			5.41	0.002	6.35	0.94	5.10	-0.31	9.58	4.16
			7.03	0.0004	7.76	0.73	5.99	-1.04	5.99	-1.04
ADCC-012	huNK	10	13.34	10	73.31	59.97	117.80	104.46	50.75	37.41
Calcein AM			13.50	1	74.76	61.26	88.64	75.14	65.97	52.47
			12.27	0.01	58.52	46.25	72.88	60.61	50.16	37.89
			13.61	0.005	57.50	43.89	69.45	55.84	39.28	25.67
			11.95	0.001	56.81	44.86	65.17	53.22	33.07	21.12
ADCC-013	PBMC	100	2.54	1	21.03	18.49	37.94	35.40	32.28	29.74
51Cr			2.45	0.1	15.50	13.05	30.82	28.37	27.18	24.73
			2.92	0.01	14.53	11.61	22.59	19.67	12.79	9.87
			2.78	0.001	3.90	1.12	8.99	6.21	3.13	0.35
ADCC-014	PBMC	100	4.64	10	53.54	48.90	56.12	51.48	ND	ND
51Cr			4.64	1	46.84	42.20	43.00	38.36	ND	ND
			4.64	0.1	45.63	40.99	39.94	35.30	ND	ND
			4.64	0.01	7.73	3.09	9.79	5.15	ND	ND
			4.64	0.001	8.83	4.19	10.81	6.17	ND	ND
Average % lysis at 1 μ g/ml concentration of mAbs						35.31		59.03		47.23

* The greater than 100% killing are due to incomplete killing by detergent used for 100% killing control.

Example 9: CD40 Is Present on the Surface of NHL Cells from Lymph Node Biopsy

5 Patients

NHL cells were isolated from biopsied lymph nodes from patients and were preserved in liquid nitrogen until use. Cell viability at the time of analysis exceeded

90%. The cells from two rituximab-sensitive and three rituximab-resistant patients (five patients in total) were stained with either a direct labeled 15B8-FITC or 15B8 plus anti-huIgG₂-FITC and analyzed by Flow cytometry. NHL cells from all the patients were found to express CD40. Table 10 shows that an average of 76% of NHL cells express CD40 (a range of 60-91%).

Table 10.

Patient ID ^a	Patient type ^b	% positive ^c	
		MS81 ^d	15B8 ^e
B	CR	n.d. ^f	91.02
J	CR	n.d.	60.36
H	NR	n.d.	85.08
H	NR	72.24	81.19
K	NR	n.d.	70.69
L	NR	n.d.	66.82

Average % positive

76

^a NHL patients treated with anti-CD20 mAb^b patient response to anti-CD20 mAb; CR=complete responder; NR=nonresponder^c % of cells in lymphocyte gate that stain positive^d MS81, agonist anti-CD40 mAb^e 15B8, antagonist anti-CD40 Mab^f n.d., not done

Example 10: CHIR-5.9 and CHIR-12.12 Do Not Stimulate Proliferation of Cancer Cells from the Lymph Nodes of NHL Patients

CD40 ligand is known to provide a stimulatory signal for the survival and proliferation of lymphoma cells from NHL patients. Binding of some anti-CD40 antibodies (agonist) can provide a similar stimulatory signal for the proliferation of patient cancer cells. Antibodies with strong B cell stimulatory activity are not suitable candidate for therapeutic treatment of B cell lymphomas. The two candidate antibodies were tested for their ability to induce proliferation of NHL cells from 3 patients. The cells isolated from lymph node (LN) biopsies were cultured with varying concentrations of candidate antibodies (range of 0.01 to 300 µg/ml) for a total of 3 days. The cell proliferation was measured by incorporation of tritiated thymidine. Neither of the two candidate mAbs induced any proliferation of cancer cells at any concentration tested (Table 11). Antibodies even in the presence of exogenously added IL-4, a B cell growth

factor, did not induce proliferation of NHL cells (tested in one of the three patient cells). These results indicate that CHIR-5.9 and CHIR-12.12 are not agonist anti-CD40 antibodies and do not stimulate proliferation *in vitro* of NHL cells from patients.

Table 11. Proliferation of cancer cells from LN of NHL patients in response to candidate anti-CD40 mAbs

Donor#	Abs conc(ug/ml)	CPM		S. index	CPM		S. index	CPM		S. index
		Cells + IgG1	cells + CHIR-5.9	CHIR-5.9	cells + CHIR-12.12	CHIR-12.12	CHIR-12.12	cells + MS81	MS81	MS81
PP	0.01	180	203	1.23	133.67	0.74	ND	ND	ND	ND
	0.1	107.5	151.67	1.41	136	1.27	ND	ND	ND	ND
	1	130.67	206.67	1.58	197.33	1.51	179	1.37	1.37	1.37
	10	152.5	245	1.61	137.33	0.90	871.67	5.71	5.71	5.71
	100	137.67	332.33	2.41	157.33	1.14	ND	ND	ND	ND
	300	137.67	254.33	1.85	100.67	0.73	ND	ND	ND	ND
MM	0.01	165	180.33	1.09	124	0.75	ND	ND	ND	ND
	0.1	180.5	149.67	0.83	111.33	0.62	ND	ND	ND	ND
	1	62	109.67	1.77	104.67	1.69	ND	ND	ND	ND
	10	91.5	93.33	1.02	100	1.09	763	8.34	8.34	8.34
	100	123	173	1.41	105.33	0.86	ND	ND	ND	ND
	300	109	183.67	1.69	157	1.44	ND	ND	ND	ND
BD (IL-4)	0.01	1591.5	1623.67	1.02	1422	0.89	ND	ND	ND	ND
	0.1	1405	1281	0.91	1316.33	0.94	ND	ND	ND	ND
	1	1526	1352.33	0.89	1160	0.76	1508.33	0.99	0.99	0.99
	10	1450	1424	0.98	1244	0.86	4146.67	2.86	2.86	2.86
	100	1406.67	1497.67	1.06	1255.33	0.89	ND	ND	ND	ND
	300	1410.33	1466.67	1.04	1233	0.87	ND	ND	ND	ND

5

Example 11: CHIR-5.9 and CHIR-12.12 Can Block CD40 Ligand-Mediated Proliferation of Cancer Cells from Non-Hodgkin's Lymphoma Patients

Engagement of CD40 by CD40 ligand induces proliferation of cancer cells from NHL patients. An antagonist anti-CD40 antibody is expected to inhibit this proliferation.

10 The two candidate anti-CD40 antibodies were tested at varying concentrations (0.01 μg /ml to 100 μg /ml) for their ability to inhibit CD40 ligand-induced proliferation of NHL cells from patients. NHL cells from patients were cultured in suspension over CD40L-expressing feeder in the presence of IL-4. The NHL cell proliferation was measured by ^3H -thymidine incorporation. Both antibodies were found to be very effective for

inhibiting CD40 ligand-induced proliferation of NHL cells (Table 12, n=2). Nearly complete inhibition of CD40 ligand-induced proliferation could be achieved at 1.0 to 10 μ g/ml concentration of antibodies.

Table 12. Inhibition of CD40 ligand-induced proliferation of cancer cells from the LN of NHL patients.

Patient	Abs Conc (ug/ml)	CPM IgG1	CHIR-5.9		CHIR-12.12		Rituximab	
			CPM	% inhibition	CPM	% inhibition	CPM	% inhibition
BD	0.01	29525.5	25369	14	24793	16	29490.3	0
	0.1	29554	20265.33	31	13671	54	29832.7	-1
	1	29486.67	6785.33	77	453	98	26355.3	11
	10	29710	506.33	98	371	99	29427.3	1
	100	29372.33	512.33	98	386.67	99	ND	ND
PP	0.01	23572	23229.33	1	23666	0	25317.3	-7
	0.1	22520	19092.33	15	17197	24	26349.7	-17
	1	23535.67	1442.33	94	802.67	97	26515.7	-13
	10	23101.5	608.67	97	221.33	99	25478.3	-10
	100	23847.33	ND	ND	252	99	ND	ND

% inhibition: 100-(CPM with test Abs/CPM with control mAb) *100%

Example 12: Effect of CHIR-5.9 on Number of Viable NHL Cells When Cultured with CD40-Ligand Bearing Cells

Effects of CHIR-5.9 on the viable NHL cell numbers when cultured with CD40-ligand bearing cells over an extended period of time (days 7, 10, and 14) were investigated. CD40 ligand-mediated signaling through CD40 is important for B cell survival. This set of experiments evaluated the effect of anti-CD40 antibodies on NHL cell numbers at days 7, 10, and 14. NHL cells from five patients were cultured in suspension over CD40L-expressing irradiated feeder cells in the presence of IL-4. The control human IgG and CHIR-5.9 antibodies were added at concentrations of 10 μ g/ml at day 0 and day 7. The viable cells under each condition were counted on the specified day. Cell numbers in the control group (IgG) increased with time as expected. Reduced numbers of cells were recovered from CHIR-5.9-treated cultures compared to control group. The greatest levels of reduction in cell numbers by CHIR-5.9 antibody were observed at day 14 and were on average 80.5% (a range of 49-94%) compared to isotype control (n=5). These data are summarized in Table 13.

Table 13. Effect of anti-CD40 antibody (CHIR-5.9/5.11) on NHL patient cell numbers over prolonged culture period (day 7, 10, and 14)

Patient	Days in culture	Viable cell numbers		% reduction compared to IgG control
		IgG	mAb CHIR-5.9/5.11	
PS	0	100000	1000000	0.00
	7	0	447500	52.14
	10	935000	504100	60.34
	14	127090	525000	48.98
		0		
MT	0	100000	1000000	0.00
	7	0	182500	31.80
	10	267600	191600	71.96
	14	683400	225000	84.48
		145000		
BRF	0	250000	250000	0.00
	7	145000	86667	40.23
	10	207500	65000	68.67
	14	570500	33330	94.16
DP	0	250000	250000	0.00
	7	188330	136670	27.43
	10	235000	128330	45.39
	14	428330	58330	86.38
PP	0	250000	250000	0.00
	7	270000	176670	34.57
	10	311670	128330	58.83
	14	458330	53330	88.36

* % reduction compare to ctrl Abs=100-(test Abs/ctrl Abs)*100

5 Example 13: CHIR-5.9 and CHIR-12.12 Are Able to Kill Cancer Cells from the Lymph Nodes of NHL Patients by ADCC

Both CHIR-5.9 and CHIR-12.12 are fully human antibodies of IgG₁ isotype and were shown to induce the killing of lymphoma cell lines *in vitro* by the mechanism of ADCC (Table 9). They were tested for their ability to kill cancer cells from a single NHL patient in *in vitro* assays. Enriched NK cells from normal volunteer donor either fresh after isolation or after culturing overnight at 37°C were used as effector cells in this assay. Similar results were obtained with both freshly isolated NK cells and NK cells

used after overnight culture. The higher level of ADCC was observed with CHIR-12.12 compared with CHIR-5.9 against the NHL cells from the patient. NHL cells also express CD20, the target antigen for rituximab (Rituxan®), which allowed for comparison of the ADCC activity of these two candidate mAbs with rituximab. Antibody CHIR-12.12 and
5 rituximab show similar level of ADCC activity with CHIR-5.9 scoring lower in this assay. These data are shown in Figures 3A and 3B.

Example 14: CHIR-5.9 and CHIR-12.12 Can Block CD40-Mediated Survival and Proliferation of Cancer Cells from CLL Patients

10 The candidate antibodies can block CD40-mediated survival and proliferation of cancer cells from CLL patients. CLL cells from patients were cultured in suspension over CD40L-expressing formaldehyde-fixed CHO cells under two different conditions: addition of human isotype antibody IgG (control); and addition of either CHIR-5.9 or CHIR-12.12 monoclonal antibody. All antibodies were added at concentrations of 1, 10,
15 and 100 $\mu\text{g/mL}$ in the absence of IL-4. The cell counts were performed at 24 and 48 h by MTS assay. Reduced numbers of cells were recovered from CHIR-5.9- (n=6) and CHIR-12.12- (n=2) treated cultures compared to control group. The greater differences in cell numbers between anti-CD40 mAb-treated and control antibody-treated cultures were seen at the 48-h time point. These data are summarized in Table 14.

Table 14. The effect of candidate antibodies on CD40-induced survival and proliferation of cancer cells from CLL patients measured at 48 h after the culture initiation

Patient#	Ab conc(μ g/ml)	Relative cell numbers			% reduction in cell numbers*	
		IgG1	CHIR-5.9/5.11	CHIR-12.12	CHIR-5.9/5.11	CHIR-12.12
1	1 10 100	269.3 1 101.5 8 130.7 1	25.27	ND	90.62	ND
			33.07	ND	67.44	ND
			40.16	ND	69.28	ND
2	1 10 100	265.5 5 227.5 7 265.9 9	75.8	ND	71.46	ND
			128.5	ND	43.53	ND
			6.4	ND	97.59	ND
3	1 10 100	85.9 70.44 77.65	35.39	ND	58.80	ND
			39.51	ND	43.91	ND
			20.95	ND	73.02	ND
4	1 10 100	80.48 63.01 55.69	15.03	ND	81.32	ND
			19.51	ND	69.04	ND
			3.65	ND	93.45	ND
5	1 10 100	90.63 78.13 63.53	91.66	89.59	-1.14	1.15
			82.28	60.41	-5.31	22.68
			86.47	39.59	-36.11	37.68
6	1 10 100	130.2 1 131.7 7 127.0 8	77.6	71.88	40.40	44.80
			78.13	73.96	40.71	43.87
			76.56	82.29	39.75	35.25

* % reduction compared to control Abs=100-(test Abs/control Abs)*100

5

Example 15: CHIR-5.9 and CHIR-12.12 Show Anti-Tumor Activity in Animal Models

Pharmacology/in vivo efficacy

The candidate mAbs are expected to produce desired pharmacological effects to reduce tumor burden by either/both of two anti-tumor mechanisms, blockade of proliferation/survival signal and induction of ADCC. The currently available human lymphoma xenograft models use long-term lymphoma cell lines that, in contrast to primary cancer cells, do not depend on CD40 stimulation for their growth and survival.

Therefore the component of these mAbs' anti-tumor activity based on blocking the tumor proliferation/survival signal is not expected to contribute to anti-tumor efficacy in these models. The efficacy in these models is dependent on the ADCC, the second anti-tumor mechanism associated with the CHIR-5.9 and CHIR-12.12 mAbs. Two xenograft human lymphoma models based on Namalwa and Daudi cell lines were assessed for anti-tumor activities of candidate mAbs. To further demonstrate their therapeutic activity, these candidate mAbs were evaluated in an unstaged and staged xenograft human lymphoma model based on the Daudi cell line.

10 *Summary of in vivo efficacy data*

When administered intraperitoneally (i.p.) once a week for a total of 3 doses, CHIR-12.12, one of the two candidate mAbs, significantly inhibited the growth of aggressive unstaged B cell lymphoma (Namalwa) in a dose-dependent manner (Figure 4). The second candidate mAb, CHIR-5.9, was tested only at a single dose in this study and was less effective than CHIR-12.12 at the same dose. Interestingly, CHIR-12.12 was found to be more efficacious in this model than rituximab. It is possible that lower efficacy by rituximab could be due to low CD20 expression on the Namalwa lymphoma cell line. The efficacy observed with candidate mAbs has greater importance because only one of the two cancer cell killing mechanisms (ADCC) is operative in the current xenograft lymphoma model. Two killing mechanisms, ADCC and blocking of survival signal, are expected to contribute to anti-tumor activities in human lymphoma patients. This is likely to increase the chance of achieving efficacy in human lymphoma patients. The candidate anti-CD40 mAbs also showed a trend toward tumor growth inhibition in a second B cell lymphoma model (non-validated Daudi model, data not shown). In follow-up studies, the two candidate antibodies were shown to have dose-dependent anti-tumor efficacy in both the unstaged and staged Daudi lymphoma models (Figures 5 and 6, respectively). In the staged Daudi model, the CHIR-12.12 mAb was more efficacious at reducing tumor volume than was a similar dose of Rituxan®.

Xenograft human B cell Lymphoma models

To ensure consistent tumor growth, T cell-deficient nude mice were whole-body irradiated at 3 Gy to further suppress the immune system one day before tumor inoculation. Tumor cells were inoculated subcutaneously in the right flank at 5×10^6 cells per mouse. Treatment was initiated either one day after tumor implantation (unstaged subcutaneous xenograft human B cell lymphoma models, Namalwa and Daudi) or when tumor volume reached 200-400 mm³ (staged Daudi model, usually 15 days after tumor inoculation). Tumor-bearing mice were injected anti-CD40 mAbs intraperitoneally (i.p.) once a week at the indicated doses. Tumor volumes were recorded twice a week. When tumor volume in any group reached 2500 mm³, the study was terminated. Note that in the staged Daudi model, tumor volume data was analyzed up to day 36 due to the death of some mice after that day. Complete regression (CR) was counted until the end of the study. Data were analyzed using ANOVA or Kruskal-Wallis test and corresponding post-test for multi-group comparison.

In the unstaged Namalwa model, anti-CD40 mAb CHIR-12.12, but not Rituxan® (rituximab), significantly ($p = < 0.01$) inhibited the growth of Namalwa tumors (tumor volume reduction of 60% versus 25% for rituximab, $n = 10$ mice/group) (Figure 4). Thus, in this model, anti-CD40 mAb CHIR-12.12 was more potent than rituximab. It is noteworthy that the second candidate mAb, CHIR-5.9, was at least as efficacious as rituximab at a dose $1/10^{\text{th}}$ that of rituximab. Both anti-CD40 mAb CHIR-12.12 and rituxan significantly prevented tumor development in the unstaged Daudi tumor model (14/15 resistance to tumor challenge) (Figure 5).

When these anti-CD40 monoclonal antibodies were further compared in a staged xenograft Daudi model, in which treatment started when the subcutaneous tumor was palpable, anti-CD40 mAb CHIR-12.12 at 1 mg/kg caused significant tumor reduction ($p = 0.003$) with 60% complete regression (6/10), while rituximab at the same dose did not significantly inhibit the tumor growth nor did it cause complete regression (0/10). See Figure 6.

In summary, the anti-CD40 mAb CHIR-12.12 significantly inhibited tumor growth in experimental lymphoma models. At the same dose and regimen, mAb CHIR-12.12 showed better anti-cancer activity than did Rituxan® (rituximab). Further, no

clinical sign of toxicity was observed at this dose and regimen. These data suggest that the anti-CD40 mAb CHIR-12.12 has potent anti-human lymphoma activity *in vitro* and in xenograft models and could be clinically effective for the treatment of lymphoma.

5 Example 16: Pharmacokinetics of CHIR-5.9 and CHIR-12.12

The pharmacokinetics of anti-CD40 mAb in mice was studied after single IV and IP dose administration. Anti-CD40 mAb exhibited high systemic bioavailability after IP administration, and prolonged terminal half-life (>5 days) (data not shown). This pilot study was conducted to aid in the design of pharmacology studies; however, it is of little
10 to no importance for the development activity of this mAb since this fully human mAb does not cross react with mouse CD40.

Example 17: Characterization of Epitope for Monoclonal Antibodies CHIR-12.12 and CHIR-5.9

15 To determine the location of the epitope on CD40 recognized by monoclonal antibodies CHIR-12.12 and CHIR-5.9, SDS-PAGE and Western blot analysis were performed. Purified CD40 (0.5 μ g) was separated on a 4-12% NUPAGE gel under reducing and non-reducing conditions, transferred to PVDF membranes, and probed with monoclonal antibodies at 10 μ g/ml concentration. Blots were probed with alkaline
20 phosphatase conjugated anti-human IgG and developed using the Western Blue^R stabilized substrate for alkaline phosphatase (Promega).

Results indicate that anti-CD40 monoclonal antibody CHIR-12.12 recognizes epitopes on both the non-reduced and reduced forms of CD40, with the non-reduced form of CD40 exhibiting greater intensity than the reduced form of CD40 (Table 15; blots not
25 shown). The fact that recognition was positive for both forms of CD40 indicates that this antibody interacts with a conformational epitope part of which is a linear sequence. Monoclonal antibody CHIR-5.9 primarily recognizes the non-reduced form of CD40 suggesting that this antibody interacts with a primarily conformational epitope (Table 15; blots not shown).

30

Table 15. Domain identification.

	Domain 1	Domain 2	Domain 3	Domain 4
mAb CHIR-12.12	-	+	-	-
mAb CHIR-5.9	-	+	-	-
mAb 15B8	+	-	-	-

To map the antigenic region on CD40, the four extracellular domains of CD40 were cloned and expressed in insect cells as GST fusion proteins. The secretion of the four domains was ensured with a GP67 secretion signal. Insect cell supernatant was analyzed by SDS-PAGE and western blot analysis to identify the domain containing the epitope.

Monoclonal antibody CHIR-12.12 recognizes an epitope on Domain 2 under both reducing and non-reducing conditions (Table 16; blots not shown). In contrast, monoclonal antibody CHIR-5.9 exhibits very weak recognition to Domain 2 (Table 16; blots not shown). Neither of these antibodies recognize Domains 1, 3, or 4 in this analysis.

Table 16. Domain 2 analysis.

	Reduced	Non-reduced
mAb CHIR-12.12	++	+++
mAb CHIR-5.9	+	+

To define more precisely the epitope recognized by mAb CHIR-12.12, peptides were synthesized from the extracellular Domain 2 of CD40, which corresponds to the sequence PCGESEFLDTWNRETHCHQHKYCDPNLGLRVQQKGTSETDTICT (residues 61-104 of the sequence shown in SEQ ID NO:10 or SEQ ID NO:12). SPOTs membranes (Sigma) containing thirty-five 10mer peptides with a 1-amino-acid offset were generated. Western blot analysis with mAb CHIR-12.12 and anti-human IgG beta-

galactosidase as secondary antibody was performed. The blot was stripped and reprobed with mAb CHIR-5.9 to determine the region recognized by this antibody

SPOTs analysis probing with anti-CD40 monoclonal antibody CHIR-12.12 at 10 µg/ml yielded positive reactions with spots 18 through 22. The sequence region covered by these peptides is shown in Table 17.

Table 17. Results of SPOTs analysis probing with anti-CD40 monoclonal antibody CHIR-12.12.

Spot Number	Sequence Region
18	HQHKYCDPNL (residues 78-87 of SEQ ID NO:10 or SEQ ID NO:12)
19	QHKYCDPNLG (residues 79-88 of SEQ ID NO:10 or SEQ ID NO:12)
20	HKYCDPNLGL (residues 80-89 of SEQ ID NO:10 or SEQ ID NO:12)
21	KYCDPNLGLR (residues 81-90 of SEQ ID NO:10 or SEQ ID NO:12)
22	YCDPNLGLRV (residues 82-91 of SEQ ID NO:10 or SEQ ID NO:12)

10

These results correspond to a linear epitope of: YCDPNL (residues 82-87 of the sequence shown in SEQ ID NO:10 or SEQ ID NO:12). This epitope contains Y82, D84, and N86, which have been predicted to be involved in the CD40-CD40 ligand interaction.

SPOTs analysis with mAb CHIR-5.9 showed a weak recognition of peptides represented by spots 20-22 shown in Table 18, suggesting involvement of the region YCDPNLGL (residues 82-89 of the sequence shown in SEQ ID NO:10 or SEQ ID NO:12) in its binding to CD40. It should be noted that the mAbs CHIR-12.12 and CHIR-5.9 compete with each other for binding to CD40 in BIACORE analysis.

15

Table 18. Results of SPOTs analysis probing with anti-CD40 monoclonal antibody CHIR-5.9.

Spot Number	Sequence Region
20	HKYCDPNLGL (residues 80-89 of SEQ ID NO:10 or SEQ ID NO:12)
21	KYCDPNLGLR (residues 81-90 of SEQ ID NO:10 or SEQ ID NO:12)
22	YCDPNLGLRV (residues 82-91 of SEQ ID NO:10 or SEQ ID NO:12)

5 The linear epitopes identified by the SPOTs analyses are within the CD40 B1 module. The sequence of the CD40 B1 module is:

HKYCDPNLGLRVQQKGTSETDTIC (residues 80-103 of SEQ ID NO:10 or SEQ ID NO:12).

10 Within the linear epitope identified for CHIR-12.12 is C83. It is known that this cysteine residue forms a disulphide bond with C103. It is likely that the conformational epitope of the CHIR-12.12 mAb contains this disulfide bond (C83-C103) and/or surrounding amino acids conformationally close to C103.

Example 18: Number of CD20 and CD40 Molecules on Namalwa and Daudi Cells

15 The number of CD20 and CD40 molecules on Namalwa and Daudi cells was determined as outlined in Figure 7, using antibody concentrations of 0.01, 0.1, 1, 10, and 100 $\mu\text{g/ml}$. As can be seen in Figure 7, the average number of CD20 molecules (target for rituximab) is greater on both the Namalwa and Daudi cell lines than is the number of CD40 molecules (target for mAb CHIR-12.12).

20

Example 19: ADCC of mAb CHIR-12.12 and Rituximab Against Daudi Lymphoma Cells

25 The rituximab and candidate mAb CHIR-12.12 were tested *in vitro* for ADCC activity at variable concentrations against lymphoma cell line Daudi as target (T) cells and purified NK cells from healthy human volunteers as effector (E) cells. Freshly

isolated human NK cells were mixed with calcein-labeled Daudi lymphoma cells at an E:T ratio of 10. The cell mixture was incubated for 4 hours at 37°C in the presence of the stated concentrations of either mAb CHIR-12.12 or rituximab. The calcein level released from lysed target cells in the supernatant was measured as Arbitrary Fluorescent Units (AFU). The percent specific lysis was calculated as: $100 \times (\text{AFU test} - \text{AFU spontaneous release}) / (\text{AFU maximal release} - \text{AFU spontaneous release})$, where AFU spontaneous release is the calcein released by target cells in the absence of antibody or NK cells, and AFU maximal release is the calcein released by target cells upon lysis by detergent.

Antibody concentration-dependent Daudi cell lysis was observed (Figure 8; Table 19 below). The maximum specific lysis of target lymphoma cells induced by anti-CD40 mAb was greater compared to the lysis induced by rituximab (63.6% versus 45.9%, n=6; paired *t* test of mAb CHIR-12.12 versus rituximab, p=0.0002). In addition, ED50 for rituximab was on average (n=6) 51.6 pM, 13-fold higher than ED50 for the anti-CD40 mAb CHIR-12.12 for this activity.

Table 19. Comparative ADCC of mAb CHIR-12.12 and rituximab against Daudi lymphoma cells.

NK Cell Donor	Maximal Killing (%)		ED50 (pM)	
	mAb CHIR-12.12	Rituximab	mAb CHIR-12.12	Rituximab
1	50.2	34.9	3.2	14.2
2	83.1	68.6	2.2	27.2
3	64.2	36.9	4.1	66.9
4	53.3	39.5	2.4	47.6
5	74.8	56.6	2.8	24.1
6	56.2	38.9	7.9	129.5
Average	63.6	45.9	3.8	51.6

Example 20: CHIR-12.12 Blocks CD40L-Mediated CD40 Survival and Signaling Pathways in Normal Human B Cells

Soluble CD40 ligand (CD40L) activates B cells and induces various aspects of functional responses, including enhancement of survival and proliferation, and activation of NF κ B, ERK/MAPK, PI3K/Akt, and p38 signaling pathways. In addition, CD40L-mediated CD40 stimulation provides survival signals by reduction of cleaved PARP and induction of the anti-apoptotic proteins, XIAP and Mcl-1, in normal B cells. CD40L-mediated CD40 stimulation also recruits TRAF2 and TRAF3 to bind CD40 cytoplasmic domain.

The following studies demonstrate that CHIR-12.12 directly inhibited all of these stimulation effects on normal human B cells. For example, CHIR-12.12 treatment resulted in increased cleavage of caspase-9, caspase-3, and PARP as well as reduction of XIAP and Mcl-1 in a time- and dose-dependent manner, restoring B cell apoptosis.

Treatment with CHIR-12.12 also inhibited phosphorylation of I κ B kinase (IKK) α and β (NF κ B pathway), ERK, Akt, and p38 in response to CD40L-mediated CD40 stimulation. Further, it was found that CHIR-12.12 did not trigger these apoptotic effects without initial CD40L-mediated CD40 stimulation.

CHIR-12.12 inhibited survival mediated by CD40 ligand by inducing cleavage of PARP.

In these experiments, 0.6×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were stimulated with 1 μ g/ml sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK). CHIR-12.12 (10 μ g/ml) and control IgG were then added. Cells were collected at 0, 20 minutes, 2 hours, 6 hours, 18 hours, and 26 hours. Cleaved caspase-9, cleaved caspase-3, cleaved PARP, and β -actin controls were detected in cell lysates by Western blot.

Briefly, it was observed that CD40L-mediated CD40 stimulation provided survival signals as it did not result in increases of cleaved caspase-9, cleaved caspase-3, or cleaved PARP over time, indicating that the cells were not undergoing apoptosis. However, treatment with CHIR-12.12 resulted in an increase of these cleavage products, indicating that CHIR-12.12 treatment abrogated the effects of CD40L binding on survival

signaling in sCD40L-stimulated normal B cells, restoring B cell apoptosis (data not shown).

CHIR-12.12 inhibited expression of "survival" anti-apoptotic proteins.

5 In these experiments, 0.6×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were stimulated with 1 $\mu\text{g/ml}$ sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK). CHIR-12.12 (10 $\mu\text{g/ml}$) and control IgG were then added. Cells were collected at 0, 20 minutes, 2 hours, 6 hours, 18 hours, and 26 hours. Mcl-1, XIAP, CD40, and β -actin controls were detected in cell lysates by Western blot.

10 Briefly, sCD40L stimulation resulted in sustained expression of Mcl-1 and XIAP over time. However, treatment of the sCD40L-stimulated cells with CHIR 12.12 resulted in a decrease in expression of these proteins overtime (data not shown). Since Mcl-1 and XIAP are "survival" signals capable of blocking the apoptotic pathway, these results demonstrate that CHIR-12.12 treatment removes the blockade against apoptosis in
15 sCD40L-stimulated normal B cells.

CHIR-12.12 treatment inhibited phosphorylation of IKK α (Ser180) and IKK β (Ser 181) in normal B cells.

In these experiments, 1.0×10^6 normal human B cells from healthy donors
20 (percent purity between 85-95%) were stimulated with 1 $\mu\text{g/ml}$ sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK). CHIR-12.12 (10 $\mu\text{g/ml}$) and control IgG were then added. Cells were collected at 0 and 20 minutes. Phosphorylated IKK α (Ser180) and IKK β (Ser 181) and total IKK β controls were detected in cell lysates by Western blot.

Briefly, stimulation by sCD40L resulted in phosphorylation of IKK α (Ser180)
25 and IKK β (Ser 181) over time; however, treatment with CHIR-12.12 abrogated this response to sCD40L stimulation in normal B cells (data not shown).

CHIR-12.12 treatment inhibited survival mediated by CD40 ligand in a dose-dependent manner.

In these experiments, 0.6×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were stimulated with 1 $\mu\text{g/ml}$ sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK). CHIR-12.12 (0.01, 0.1, 0.2, 0.5, 1.0 $\mu\text{g/ml}$) and control IgG were then added. Cells were collected at 24 hours. Cleaved PARP, and β -actin controls were detected in cell lysates by Western blot.

Briefly, CHIR-12.12 treatment resulted in increase of PARP cleavage in sCD40L stimulated cells in a dose-dependent manner and therefore abrogated the survival signaling pathway in sCD40L-stimulated normal B cells (data not shown).

CHIR-12.12 inhibited expression of "survival" anti-apoptotic proteins in a dose-dependent manner.

In these experiments, 0.6×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were stimulated with 1 $\mu\text{g/ml}$ sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK). CHIR-12.12 (0.5, 2, and 10 $\mu\text{g/ml}$) and control IgG were then added. Cells were collected at 22 hours. Mcl-1, XIAP, cleaved PARP, and β -actin controls were detected in cell lysates by Western blot.

Briefly, CHIR-12.12 treatment reduced Mcl-1 and XIAP expression and increased cleaved PARP expression in sCD40L-stimulated cells in a dose-dependent manner, and thus abrogated these blockades to the apoptotic pathway in sCD40L-stimulated normal B cells (data not shown).

CHIR-12.12 did not affect expression of anti-apoptotic proteins, cleaved-PARP, and XIAP, in the absence of soluble CD40L signaling.

In these experiments, 1.0×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were treated with CHIR-12.12 (10 $\mu\text{g/ml}$) and control IgG only (i.e., cells were not pre-stimulated with sCD40L before adding antibody). Cells

were collected at 0, 4, 14, and 16 hours. XIAP, cleaved PARP, and β -actin controls were detected in cell lysates by Western blot.

Briefly, the results show that without sCD40L stimulation, the cells expressed increased concentrations of cleaved PARP, while expression of XIAP remained constant, in both IgG treated control cells and CHIR-12.12 cells (data not shown). These data indicate that CHIR-12.12 does not trigger apoptosis in normal human B cells without CD40L stimulation.

CHIR-12.12 inhibits phosphorylation of IKK α (Ser180) and IKK β (Ser181), Akt, ERK, and p38 in normal B cells.

In these experiments, 1.0×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were serum starved in 1% FBS-containing media and stimulated with 1 μ g/ml sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK). The cultures were treated with CHIR-12.12 (1 and 10 μ g/ml) and control IgG. Cells were collected at 0 and 20 minutes. Phospho-IKK α , phospho-IKK β , total IKK β , phospho-ERK, total ERK, phospho-Akt, total Akt, phospho-p38, and total p38 were detected in cell lysates by Western blot.

Briefly, sCD40L stimulation resulted in increases in IKK α/β phosphorylation, ERK phosphorylation, Akt phosphorylation, and p38 phosphorylation, thus leading to survival and or proliferation of the cells. Treatment of the cells with CHIR-12.12 abrogated the effects of sCD40L stimulation on these signaling pathways in normal B cells (data not shown).

CHIR 12.12 inhibits multiple signaling pathways such as PI3K and MEK /ERK in the CD40 signaling cascade.

In these experiments, 1.0×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were serum starved in 1% FBS-containing media and stimulated with 1 μ g/ml sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK). The cultures were also treated with CHIR-12.12 (1 and 10 μ g/ml), Wortmanin, (a PI3K/Akt

inhibitor; 1 and 10 μ M), LY 294002 (a PI3K/Akt inhibitor; 10 and 30 μ M), and PD 98095 (a MEK inhibitor; 10 and 30 μ g/ml). Cells were collected at 0 and 20 minutes. Phospho-ERK, phospho-Akt, total Akt, phospho-IKK α/β , and total were detected in cell lysates by Western blot.

5 Briefly, the results show that CHIR-12.12 abrogated the phosphorylation of all of these signal transduction molecules, whereas the signal transduction inhibitors showed only specific abrogation of signaling, indicating that CHIR-12.12 likely inhibits upstream of these signal transduction molecules mediated by CD40L stimulation (data not shown).

10 *CHIR-12.12 inhibits the binding of signaling molecules TRAF2 and TRAF3 to the cytoplasmic domain of CD40 in normal B cells.*

In these experiments, 4.0×10^6 normal human B cells from healthy donors (percent purity between 85-95%) were serum starved for four hours in 1% FBS-containing media and stimulated with 1 μ g/ml sCD40L (Alexis Corp., Bingham, Nottinghamshire, UK) for 20 minutes. Cells were collected at 0 and 20 minutes. CD40 was immunoprecipitated using polyclonal anti-CD40 (Santa Cruz Biotechnology, CA), and was probed in a Western blot with anti-TRAF2 mAb (Santa Cruz Biotechnology, CA), anti-TRAF3 mAb (Santa Cruz Biotechnology, CA), and anti-CD40 mAb (Santa Cruz Biotechnology, CA).

20 Briefly, the results show that TRAF2 and TRAF3 co-precipitated with CD40 after sCD40L stimulation. In contrast, treatment with CHIR-12.12 abrogated formation of the CD40-TRAF2/3 signaling complex in sCD40L-stimulated normal B cells. There were no changes in CD40 expression (data not shown).

Without being bound by theory, the results of these experiments, and the results in 25 the examples outlined above, indicate that the CHIR-12.12 antibody is a dual action antagonist anti-CD40 monoclonal antibody having a unique combination of attributes. This fully human monoclonal antibody blocks CD40L-mediated CD40 signaling pathways for survival and proliferation of B cells; this antagonism leads to ultimate cell death. CHIR-12.12 also mediates recognition and binding by effector cells, initiating 30 antibody dependent cellular cytotoxicity (ADCC). Once CHIR-12.12 is bound to effector cells, cytolytic enzymes are released, leading to B-cell apoptosis and lysis. CHIR-12.12

is a more potent anti-tumor antibody than is rituximab when compared in pre-clinical tumor models.

Example 21: Agonist and Antagonist Activity Against Primary Cancer Cell from NHL,

CLL, and NM Patients

In collaboration with clinical investigators, the candidate mAbs is tested for a variety of activities (listed below) against primary cancer cells from NHL and CLL and multiple myeloma patients.

- Agonist effect in proliferation assays (8 NHL patients, 8 CLL patients and 8 MM patients)
- Antagonist effect in proliferation assays (8 NHL patients, 8 CLL patients and 8 MM patients)
- Apoptotic effect by Annexin V assay (3-4 NHL patients, 4 CLL patients, and 4 MM patients)
- Reversing survival signal by Annexin V assay (3 NHL patients, 3 CLL patients and 3 MM patients)
- Complement dependent cytotoxicity (4 NHL patients, 4 CLL patients, and 4MM patients)
- Antibody dependent cytotoxicity (6 NHL patients, 6 CLL patients and 6 MM patients)

Example 22: Identification of Relevant Animal Species for Toxicity Studies

As these two candidate antibodies do not cross-react with rodent CD40, other species must be identified for testing toxicologic effects.

The ability of the two candidate anti-CD40 antibodies to cross-react with animal CD40 is tested by flow cytometric assays. Rat, rabbit, dog, cynomolgus monkeys and marmoset monkeys are tested for this study.

The candidate antibodies show antagonist activity upon binding to CD40 on human B cells. To identify an animal species that has similar response to candidate antibodies, lymphocytes from species that show binding to candidate antibodies are tested in proliferation assays for antagonist activity. The lymphocytes from the species selected

for antagonistic binding of candidate antibodies are further tested for their ability to serve as effector cells for killing CD40-expressing lymphoma cell lines through the mechanism of ADCC. Finally the selected animal species are tested in an IHC study for the tissue-binding pattern of candidate antibodies. The animal species responding to the candidate antibodies in these assays in a manner similar to that observed for human cells are chosen for toxicology studies.

Initial studies indicate that the candidate anti-CD40 mAbs cross-react with cynomolgus monkey CD40.

10 Example 23: Tumor Targeting Profile of CHIR-5.9 and CHIR-12.12

To determine the relative tumor targeting profile of CHIR-12.12 and CHIR-5.9 mAbs, fluorescent-labeled candidate mAbs and isotype control antibodies are administered into tumor-bearing mice. Tumor specimens and normal organs are harvested at different time points after dosing. The accumulation of labeled antibody in tumors and normal organs is analyzed.

Example 24: Mechanism of Action of CHIR-5.9 and CHIR-12.12

To elucidate the mechanism(s) that mediates the tumor growth inhibition by the CHIR-5.9 and CHIR-12.12 mAbs, the following studies are undertaken:

20 Fc-receptor knock-out or blockage model: ADCC is mediated by binding of effector cells such as NK, macrophage, and monocytes to the Fc portion of antibody through Fc receptor. Mice deficient in activating Fc receptors as well as antibodies engineered to disrupt Fc binding to those receptors will block the ADCC mediated tumor growth inhibition. Loss or significantly reduced tumor inhibition in this model will suggest that the tumor growth inhibition by these two candidate mAbs is mainly mediated by ADCC mechanism.

Example 25: Liquid Pharmaceutical Formulation for Antagonist Anti-CD40 Antibodies

The objective of this study was to investigate the effects of solution pH on stability of the antagonist anti-CD40 antibody CHIR-12.12 by both biophysical and biochemical methods in order to select the optimum solution environment for this

antibody. Differential Scanning Calorimetry (DSC) results showed that the conformation stability of CHIR-12.12 is optimal in formulations having pH 5.5-6.5. Based on a combination of SDS-PAGE, Size-Exclusion HPLC (SEC-HPLC), and Cation-Exchange HPLC (CEX-HPLC) analysis, the physicochemical stability of CHIR-12.12 is optimal at about pH 5.0-5.5. In view of these results, one recommended liquid pharmaceutical formulation comprising this antibody is a formulation comprising CHIR-12.12 at about 20 mg/ml formulated in about 10 mM sodium succinate, about 150 mM sodium chloride, and having a pH of about pH 5.5.

10 Materials and Methods

The CHIR-12.12 antibody used in the formulation studies is a human monoclonal antibody produced by a CHO cell culture process. This MAb has a molecular weight of 150 kDa and consists of two light chains and two heavy chains linked together by disulfide bands. It is targeted against the CD40 cell surface receptor on CD40-expressing cells, including normal and malignant B cells, for treatment of various cancers and autoimmune/inflammatory diseases.

The anti-CD40 drug substance used for this study was a CHO-derived purified anti-CD40 (CHIR-12.12) bulk lot. The composition of the drug substance was 9.7 mg/ml CHIR-12.12 antibody in 10 mM sodium citrate, 150 mM sodium chloride, at pH 6.5. The control sample in the study was the received drug substance, followed by freezing at $\leq -60^{\circ}\text{C}$, thawing at RT and testing along with stability samples at predetermined time points. The stability samples were prepared by dialysis of the drug substance against different pH solutions and the CHIR-12.12 concentration in each sample was determined by UV 280 as presented in Table 20.

Table 20. CHIR-12.12 formulations.

Buffer Composition	pH	CHIR-12.12 Concentration (mg/ml)
10 mM sodium citrate, 150 mM sodium chloride	4.5	9.0
10 mM sodium succinate, 150 mM sodium chloride	5.0	9.3
10 mM sodium succinate, 150 mM sodium chloride	5.5	9.2
10 mM sodium citrate, 150 mM sodium chloride	6.0	9.7
10 mM sodium citrate, 150 mM sodium chloride	6.5	9.4

10 mM sodium phosphate, 150 mM sodium chloride	7.0	9.4
10 mM sodium phosphate, 150 mM sodium chloride	7.5	9.5
10 mM glycine, 150 mM sodium chloride	9.0	9.5

Physicochemical stability of the CHIR-12.12 antibody in the various formulations was assayed using the following protocols.

Differential Scanning Calorimetry (DSC)

Conformational stability of different formulation samples was monitored using a MicroCal VP-DSC upon heating 15°C to 90°C at 1°C/min.

SDS-PAGE

Fragmentation and aggregation were estimated using 4-20% Tris-Glycine Gel under non-reducing and reducing conditions. Protein was detected by Coomassie blue staining.

Size Exclusion Chromatograph (SEC-HPLC)

Protein fragmentation and aggregation were also measured by a Water Alliance HPLC with a Tosohaas TSK-GEL 3000SWXL column, 100 mM sodium phosphate, pH 7.0 as mobile phase at a flow rate of 0.7 ml/min.

Cation Exchange Chromatography (CEX-HPLC)

Charge change related degradation was measured using Waters 600s HPLC system with a Dionex Propac WCX-10 column, 50 mM HEPES, pH 7.3 as mobile phase A and 50 mM HEPES containing 500 mM NaCl, pH 7.3 as mobile phase B at a flow rate of 0.5°C/min.

Results and Discussion

Conformational stability study.

Thermal unfolding of CHIR-12.12 revealed at least two thermal transitions, probably representing unfolding melting of the Fab and the Fc domains, respectively. At higher temperatures, the protein presumably aggregated, resulting in loss of DSC signal.

For the formulation screening purpose, the lowest thermal transition temperature was defined as the melting temperature, T_m , in this study. Figure 13 shows the thermal melting temperature as a function of formulation pHs. Formulations at pH 5.5-6.5 provided anti-CD40 with higher conformational stability as demonstrated by the higher thermal melting temperatures.

SDS-PAGE analysis.

The CHIR-12.12 formulation samples at pH 4.5-9.0 were incubated at 40°C for 2 months and subjected to SDS-PAGE analysis (data not shown). Under non-reducing conditions, species with molecular weight (MW) of 23 kDa and 27 kDa were observed in formulations above pH 5.5, and species with MW of 51 kDa were observed in all formulations, but appeared less at pH 5.0-5.5. A species with MW of 100 kDa could be seen at pH 7.5 and pH 9.0.

Under reducing conditions, CHIR-12.12 was reduced into free heavy chains and light chains with MW of 50 kDa and 24 kDa, respectively. The 100 kDa species seemed not fully reducible and increased with increasing solution pH, suggesting non-disulfide covalent association might occur in the molecules. Since there were other species with unknown identities on SDS-PAGE, stability comparison of each formulation is based on the remaining purity of CHIR-12.12. Formulations at pH 5.0-6.0 provided a more stable environment to CHIR-12.12. Few aggregates were detected by SDS-PAGE (data not shown).

SEC-HPLC analysis .

SEC-HPLC analysis detected the intact CHIR-12.12 as the main peak species, an aggregation species as a front peak species separate from the main peak species, a large fragment species as a shoulder peak on the back of the main peak species, and small fragment species were detected post-main peak species. After incubation at 5°C and 25°C for 3 months, negligible amounts of protein fragments and aggregates (<1.0%) were detected in the above formulations and the CHIR-12.12 main peak species remained

greater than 99% purity (data not shown). However, protein fragments gradually developed upon storage at 40°C and more fragments formed at pH 4.5 and pH 6.5-9.0, as shown in Table 21. After incubating the CHIR-12.12 formulations at 40°C for 3 months, about 2-3% aggregates were detected in pH 7.5 and pH 9.0, while less than 1% aggregates were detected in other pH formulations (data not shown). The SEC-HPLC results indicate CHIR-12.12 is more stable at about pH 5.0-6.0.

Table 21. SEC-HPLC results of CHIR-12.12 stability samples under real-time and accelerated storage conditions.

Sample	Main peak %				Fragments %			
	t=0	40°C 1 m	40°C 2 m	40°C 3 m	t=0	40°C 1 m	40°C 2 m	40°C 3 m
Control	99.4	99.2	99.9	99.5	<1.0	<1.0	<1.0	<1.0
pH 4.5	99.4	93.2	86.0	81.3	<1.0	6.4	13.2	18.1
pH 5.0	99.8	98.7	91.3	89.2	<1.0	<1.0	7.8	10.2
pH 5.5	99.8	98.9	91.4	90.6	<1.0	<1.0	7.6	8.8
pH 6.0	99.6	97.7	90.4	87.3	<1.0	1.9	8.2	11.7
pH 6.5	99.3	93.4	89.0	86.9	<1.0	5.6	9.9	12.4
pH 7.0	99.2	93.9	87.4	85.1	<1.0	5.5	11.1	13.5
pH 7.5	99.1	92.8	84.4	81.9	<1.0	6.4	12.9	16.2
pH 9.0	99.3	82.4	61.6	50.6	<1.0	15.4	36.2	47.6

CEX-HPLC analysis.

CEX-HPLC analysis detected the intact CHIR-12.12 as the main peak species, acidic variants eluted earlier than the main peak species, and C-terminal lysine addition variants eluted post-main peak species. Table 22 shows the dependence of the percentages of the remaining main peak CHIR-12.12 species and acidic variants on solution pH. The control sample already contained a high degree of acidic species (~33%), probably due to early-stage fermentation and purification processes. The susceptibility of CHIR-12.12 to higher pH solutions is evidenced by two facts. First, the initial formulation sample at pH 9.0 (t=0) already generated 12% more acidic species than the control. Second, the percentage of acidic species increased sharply with increasing pH. The charge change-related degradation is likely due to deamidation. The

above data indicate that this type of degradation of CHIR-12.12 was minimized at about pH 5.0-5.5.

Table 22. Percentage of peak area by CEX-HPLC for CHIR-12.12 in different pH formulations under real-time and accelerated storage conditions.

Sample	Main peak %					Acidic variants %				
	t=0	5°C 3m	25°C 3 m	40°C 1 m	40°C 2 m	t=0	5°C 3m	25°C 3 m	40°C 1 m	40°C 2 m
Control	49.2	49.8	49.8	49.2	50.3	32.0	33.7	33.7	32.0	33.6
pH 4.5	48.5	49.7	43.7	39.7	30.0	32.5	32.6	38.0	44.2	56.4
pH 5.0	49.6	49.8	48.3	40.6	31.4	32.7	31.8	35.0	44.3	57.1
pH 5.5	50.7	50.3	48.1	40.0	30.2	32.6	31.8	37.8	48.9	63.3
pH 6.0	50.2	49.9	47.9	37.4	23.9	33.1	33.6	38.5	54.9	72.7
pH 6.5	49.4	49.9	42.3	29.7	14.6	33.3	33.6	47.7	65.2	84.6
pH 7.0	49.7	49.9	21.9	-	-	34.4	36.4	64.4	-	-
pH 7.5	49.3	48.3	12.7	-	-	35.5	40.1	79.2	-	-
pH 9.0	41.3	31.8	-	-	-	44.7	59.9	-	-	-

Conclusion

The pH has a significant effect on conformational and physicochemical stabilities of CHIR-12.12. Charge change-related degradation was determined to be the main degradation pathway for CHIR-12.12, which was minimized at pH 5.0-5.5. Based on overall stability data, one recommended liquid pharmaceutical formulation comprising this antibody is a formulation comprising CHIR-12.12 at about 20 mg/ml formulated in about 10 mM sodium succinate, about 150 mM sodium chloride, and having a pH of about pH 5.5.

Example 26: Clinical Studies with CHIR-5.9 and CHIR-12.12

Clinical Objectives

The overall objective is to provide an effective therapy for B cell tumors by targeting them with an anti-CD40 IgG1. These tumors include B-cell lymphoma, Chronic Lymphocytic Lymphoma (CLL), Acute Lymphoblastic Leukemia (ALL), Multiple Myeloma (MM), Waldenstrom's Macroglobulinemia, and Systemic Castleman's Disease.

The signal for these diseases is determined in phase II although some measure of activity may be obtained in phase I. Initially the agent is studied as a single agent, but will be combined with other agents, chemotherapeutics, and other antibodies, as development proceeds.

5. Phase I

- Evaluate safety and pharmacokinetics – dose escalation in subjects with B cell malignancies.
- Choose dose based on safety, tolerability, and change in serum markers of CD40. In general an MTD is sought but other indications of efficacy (depletion of CD40+ B cells, etc.) may be adequate for dose finding.
- Consideration of more than one dose especially for different indications, e.g., the CLL dose may be different than the NHL. Thus, some dose finding may be necessary in phase II.
- Patients are dosed weekly with real-time pharmacokinetic (Pk) sampling. Initially a 4-week cycle is the maximum dosing allowed. The Pk may be highly variable depending on the disease studied, density of CD40 etc.
- This trial(s) is open to subjects with B-cell lymphoma, CLL, and potentially other B cell malignancies.
- Decision to discontinue or continue studies is based on safety, dose, and preliminary evidence of anti-tumor activity.
- Activity of drug as determined by response rate is determined in Phase II.
- Identify dose(s) for Phase II.

Phase II

- Several trials will be initiated in the above-mentioned tumor types with concentration on B-cell lymphoma, CLL, and Multiple Myeloma (MM). Separate trials may be required in low grade and intermediate/high grade NHL as CD40 may have a different function depending on the grade of lymphoma. In low-grade disease, CD40 acts more as a survival factor, preventing apoptosis. In higher-grade disease, interruption of

CD40 signaling may lead to cell death. More than one dose, and more than one schedule may be tested in a randomized phase II setting.

In each disease, target a population that has failed current standard of care:

- 5
 - CLL: patients who were resistant to Campath® and chemotherapy.
 - Low grade NHL: Rituxan® or CHOP-R failures
 - Intermediate NHL: CHOP-R failures
 - Multiple Myeloma: Chemotherapy failures
- 10
 - ✓ Decision to discontinue or continue with study is based on proof of therapeutic concept in Phase II
 - ✓ Determine whether surrogate marker can be used as early indication of clinical efficacy
 - ✓ Identify doses for Phase III

15

Phase III

Phase III will depend on where the signal is detected in phase II, and what competing therapies are considered to be the standard. If the signal is in a stage of disease where there is no standard of therapy, then a single arm, well-controlled study could serve as a pivotal trial. If there are competing agents that are considered standard, then head-to-head studies are conducted.

20

25 Many modifications and other embodiments of the inventions set forth herein will come to mind to one skilled in the art to which these inventions pertain having the benefit of the teachings presented in the foregoing descriptions and the associated drawings. Therefore, it is to be understood that the inventions are not to be limited to the specific embodiments disclosed and that modifications and other embodiments are intended to be included within the scope of the appended claims and list of embodiments disclosed

30

herein. Although specific terms are employed herein, they are used in a generic and descriptive sense only and not for purposes of limitation.


All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All
5 publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Applicant's or agent's file reference	PP20107.004	International application No.	PCT/US2004/
--	-------------	-------------------------------	-------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 18, line 25	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 USA	
Date of deposit 17 September 2003	Accession Number PTA-5542
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 22, line 20; page 75, line 11; page 142 line 9	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indicators are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 


For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

Applicant's or agent's file reference	PP20107.004	International application No.	PCT/US2004/
--	-------------	-------------------------------	-------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 18, line 25	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution <p style="text-align: center;">American Type Culture Collection</p>	
Address of depository institution <i>(including postal code and country)</i> <p style="text-align: center;">10801 University Blvd. Manassas, VA 20110-2209 USA</p>	
Date of deposit <p style="text-align: center;">17 September 2003</p>	Accession Number <p style="text-align: center;">PTA-5543</p>
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
Page 22, line 20; page 75, line 11; page 127, line 15; page 142, line 10	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indicators are not for all designated States)</i>	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

ATCC

10801 University Blvd • Manassas, VA 20110-2209 • Telephone: 703-365-2700 • FAX: 703-365-2746

**BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE****INTERNATIONAL FORM****RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.****To: (Name and Address of Depositor or Attorney)**Chiron Corporation
Attn: Karen Van Note
4560 Horton Street
Emeryville, CA 94608

Deposited on Behalf of: Chiron Corporation

Identification Reference by Depositor:Mouse Hybridoma 131.2F8.5.9: CMCC#12047
Mouse Hybridoma 153.8E2D10D6.12.12: CMCC#12056**Patent Deposit Designation**PTA-5542
PTA-5543

The deposits were accompanied by: a scientific description, a proposed taxonomic description indicated above. The deposits were received September 17, 2003 by this International Depository Authority and have been accepted.

AT YOUR REQUEST: ☒ We will inform you of requests for the strains for 30 years.

The strains will be made available if a patent office signatory to the Budapest Treaty certifies one's right to receive, or if a U.S. Patent is issued citing the strains, and ATCC is instructed by the United States Patent & Trademark Office or the depositor to release said strains.

If the cultures should die or be destroyed during the effective term of the deposit, it shall be your responsibility to replace them with living cultures of the same.

The strains will be maintained for a period of at least 30 years from date of deposit, or five years after the most recent request for a sample, whichever is longer. The United States and many other countries are signatory to the Budapest Treaty.

The viability of the cultures cited above was tested September 23, 2003. On that date, the cultures were viable.

International Depository Authority: American Type Culture Collection, Manassas, VA 20110-2209 USA.

Signature of person having authority to represent ATCC:
Marie Harris, Patent Specialist, ATCC Patent DepositoryDate: October 2, 2003cc: Lisa Alexander
Ref: Docket or Case No.: P20107.001

B249

Received Nov-03-03 03:20pm

From-510 923 4765

To-Intellectual Property Page 002

THAT WHICH IS CLAIMED:

1. A human monoclonal antibody that is capable of specifically binding to a human CD40 antigen expressed on the surface of a human CD40-expressing cell, said monoclonal antibody being free of significant agonist activity, wherein said monoclonal antibody exhibits increased anti-tumor activity relative to an equivalent amount of the monoclonal chimeric anti-CD20 monoclonal antibody IDEC-C2B8, wherein said anti-tumor activity is assayed in a staged nude mouse xenograft tumor model using the Daudi human B cell lymphoma cell line.

2. The human monoclonal antibody of claim 1, wherein said antibody is selected from the group consisting of:

a) the monoclonal antibody CHIR-12.12;
b) the monoclonal antibody produced by the hybridoma cell line 12.12, deposited with the ATCC as Patent Deposit No. PTA-5543;
c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

d) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

e) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 12.12;

f) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

g) a monoclonal antibody that competes with the monoclonal antibody CHIR-12.12 in a competitive binding assay;

h) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-g), wherein said antibody is recombinantly produced; and

i) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-h), wherein said fragment retains the capability of specifically binding to said human CD40 antigen.

3. The monoclonal antibody of claim 1, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

4. A hybridoma cell line capable of producing the monoclonal antibody of claim 1.

5. A method for treating a cancer characterized by expression of CD40, comprising administering to a human patient an effective amount of a human anti-CD40 monoclonal antibody of claim 1.

6. The method of claim 5, wherein said cancer is selected from the group consisting of a non-Hodgkins lymphoma, chronic lymphocytic leukemia, multiple myeloma, B cell lymphoma, high-grade B cell lymphoma, intermediate-grade B cell lymphoma, low-grade B cell lymphoma, B cell acute lymphoblastic leukemia, myeloblastic leukemia, and Hodgkin's disease.

7. A human monoclonal antibody that is capable of specifically binding to a human CD40 antigen expressed on the surface of a human CD40-expressing cell, said monoclonal antibody being free of significant agonist activity, whereby, when said monoclonal antibody binds to the CD40 antigen expressed on the surface of said cell, the growth or differentiation of said cell is inhibited, wherein said antibody is selected from the group consisting of:

a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in
5 SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in
10 SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group
15 consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87
20 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

25 j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen.

30

8. The antigen-binding fragment of claim 7, wherein said fragment is selected from the group consisting of a Fab fragment, an F(ab')₂ fragment, an Fv fragment, and a single-chain Fv fragment.

5 9. The monoclonal antibody of claim 7, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10⁻⁶ M to about 10⁻¹² M.

10 10. An isolated nucleic acid molecule comprising a polynucleotide that encodes an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, and SEQ ID NO:8.

15 11. A hybridoma cell line capable of producing a human monoclonal antibody having specificity for a human CD40 antigen expressed on the surface of a human CD40-expressing cell, whereby said monoclonal antibody is free of significant agonist activity, whereby, when said monoclonal antibody binds to the CD40 antigen expressed on the surface of said cell, the growth or differentiation of said cell is inhibited, and wherein said monoclonal antibody is selected from the group consisting of:

20 a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

25 c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

30 d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ

ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay; and,

j) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of a)-i), wherein said fragment retains the capability of specifically binding to said human CD40 antigen.

12. A method for inhibiting growth or differentiation of a normal human B cell, comprising contacting said B cell with an effective amount of a human anti-CD40 monoclonal antibody selected from the group consisting of:

a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in

SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains

the capability of specifically binding to said human CD40 antigen; said antibody or fragment thereof being free of significant agonist activity, and whereby when said antibody or fragment thereof binds to said CD40 antigen on said B cell, the growth or differentiation of said B cell is inhibited.

13. The method of claim 12, wherein said monoclonal antibody or fragment thereof binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

14. The method of claim 12, wherein said fragment is selected from the group consisting of a Fab fragment, an $F(ab')_2$ fragment, an Fv fragment, and a single-chain Fv fragment.

15. A method for inhibiting proliferation of a normal human B cell, wherein said proliferation is augmented by the interaction of a CD40 ligand with a CD40 antigen expressed on the surface of said B cell, said method comprising contacting said B cell with an effective amount of a human anti-CD40 monoclonal antibody selected from the group consisting of:

a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen; said antibody or fragment thereof being free of significant agonist activity, and whereby when said antibody or fragment thereof binds to said CD40 antigen on said B cell, the growth or differentiation of said B cell is inhibited.

16. The method of claim 15, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

17. The method of claim 15, wherein said fragment is selected from the group consisting of a Fab fragment, an $F(ab')_2$ fragment, an Fv fragment, and a single-chain Fv fragment.

18. A method for inhibiting antibody production by B cells in a human patient, comprising administering to a human patient an effective amount of a human anti-CD40 monoclonal antibody or fragment thereof selected from the group consisting of:

- a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;
- b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;
- c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;
- d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ

ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen;

said antibody or fragment thereof being free of significant agonist activity, and whereby when said antibody or fragment thereof binds to said CD40 antigen on said B cell, the growth or differentiation of said B cell is inhibited.

19. The method of claim 18, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

20. The method of claim 18, wherein said fragment is selected from the group consisting of a Fab fragment, an $F(ab')_2$ fragment, an Fv fragment, and a single-chain Fv fragment.

21. A method for inhibiting growth of cancer cells of B cell lineage, comprising contacting said cancer cells with an effective amount of a human anti-CD40 monoclonal antibody selected from the group consisting of:

a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

5 b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

10 d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

20 f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

25 h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

30 j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen; said antibody or fragment thereof being free of significant agonist activity, and whereby
5 when said antibody or fragment thereof binds to said CD40 antigen on said B cell, the growth or differentiation of said B cell is inhibited.

22. The method of claim 21, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

23. The method of claim 21, wherein said fragment is selected from the group consisting of a Fab fragment, an $F(ab')_2$ fragment, an Fv fragment, and a single-chain Fv fragment.

24. The method of claim 21, wherein the cancer is selected from the group consisting of non-Hodgkins lymphoma, chronic lymphocytic leukemia, multiple myeloma, B cell lymphoma, high-grade B cell lymphoma, intermediate-grade B cell lymphoma, low-grade B cell lymphoma, B cell acute lymphoblastic leukemia, myeloblastic leukemia, and Hodgkin's disease.

25. A method for treating an autoimmune disease, comprising administering to a human patient an effective amount of a human anti-CD40 monoclonal antibody selected from the group consisting of:

- a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;
- b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;
- c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen; said antibody or fragment thereof being free of significant agonist activity, and whereby when said antibody or fragment thereof binds to said CD40 antigen on said B cell, the growth or differentiation of said B cell is inhibited.

26. The method of claim 25, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

27. The method of claim 25, wherein said fragment is selected from the group consisting of a Fab fragment, an F(ab')₂ fragment, an Fv fragment, and a single-chain Fv fragment.

5 28. The method of claim 25, wherein said autoimmune disease is selected from the group consisting of systemic lupus erythematosus, autoimmune thrombocytopenic purpura, Rheumatoid arthritis, multiple sclerosis, ankylosing spondylitis, myasthenia gravis, and pemphigus vulgaris.

10 29. A method for treating a cancer characterized by expression of CD40, comprising administering to a human patient an effective amount of a human anti-CD40 monoclonal antibody selected from the group consisting of:

a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;
b) the monoclonal antibody produced by the hybridoma cell line 5.9 or
15 12.12;
c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID
20 NO:8;

d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID
25 NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

30 f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

5 i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

10 k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen; said antibody or fragment thereof being free of significant agonist activity, and whereby when said antibody or fragment thereof binds to said CD40 antigen on said B cell, the growth or differentiation of said B cell is inhibited.

15

30. The method of claim 29, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

20 31. The method of claim 29, wherein said fragment is selected from the group consisting of a Fab fragment, an $F(ab')_2$ fragment, an Fv fragment, and a single-chain Fv fragment.

25 32. A method for identifying an antibody that has antagonist activity toward CD40-expressing cells, comprising performing a competitive binding assay with a monoclonal antibody selected from the group consisting of:

a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

30 c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ

ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in
5 SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group
10 consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87
15 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12; and

i) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-h), wherein said antibody is recombinantly produced; and

20 j) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of a)-i), wherein said fragment retains the capability of specifically binding to said human CD40 antigen.

33. An antagonist anti-CD40 monoclonal antibody that specifically binds
25 Domain 2 of CD40.

34. The monoclonal antibody of claim 33, wherein said antibody is a human antibody.

30 35. The monoclonal antibody of claim 34, wherein said antibody is free of significant agonist activity.

36. The monoclonal antibody of claim 33, wherein said antibody has the binding specificity of an antibody selected from the group consisting of the antibody produced by hybridoma cell line 5.9 and the antibody produced by hybridoma cell line 12.12.

37. The monoclonal antibody of claim 33, wherein said antibody is selected from the group consisting of the antibody produced by hybridoma cell line deposited with the ATCC as Patent Deposit No. PTA -5542 and hybridoma cell line deposited with the ATCC as Patent Deposit No. PTA-5543.

38. The monoclonal antibody of claim 33, wherein said antibody has the binding specificity of monoclonal antibody CHIR-12.12 or CHIR-5.9.

39. The monoclonal antibody of claim 33, wherein said antibody binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12.

40. The monoclonal antibody of claim 33, wherein said antibody is selected from the group consisting of:

a) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

b) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

c) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 12.12;

d) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

e) a monoclonal antibody that competes with the monoclonal antibody CHIR-12.12 in a competitive binding assay; and

5 f) a monoclonal antibody that is an antigen-binding fragment of the CHIR-12.12 monoclonal antibody or the foregoing monoclonal antibodies in preceding items (a)-(e), where the fragment retains the capability of specifically binding to the human CD40 antigen.

10 41. A method for inhibiting a CD40 ligand-mediated CD40 signaling pathway in a human CD40-expressing cell, said method comprising contacting said cell with an effective amount of a human anti-CD40 monoclonal antibody selected from the group consisting of:

a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

15 b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

20 d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

25 e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

30

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

5 h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

j) the monoclonal antibody of preceding item a) or a monoclonal antibody of
10 any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen.

15 42. The method of claim 41, wherein said monoclonal antibody binds to said human CD40 antigen with an affinity (K_D) of at least about 10^{-6} M to about 10^{-12} M.

43. The method of claim 41, wherein said fragment is selected from the group consisting of a Fab fragment, an $F(ab')_2$ fragment, an Fv fragment, and a single-chain Fv
20 fragment.

44. The method of claim 41, wherein said human CD40-expressing cell is a normal human B cell or a malignant human B cell and said CD40 signaling pathway is B cell survival.

25 45. A pharmaceutical composition comprising a human monoclonal antibody that is capable of specifically binding to a human CD40 antigen expressed on the surface of a human CD40-expressing cell, said monoclonal antibody being free of significant agonist activity, wherein said antibody is selected from the group consisting of:

30 a) the monoclonal antibody CHIR-5.9 or CHIR-12.12;

b) the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

c) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:6, the sequence shown in
5 SEQ ID NO:7, the sequence shown in SEQ ID NO:8, both the sequence shown in SEQ ID NO:6 and SEQ ID NO:7, and both the sequence shown in SEQ ID NO:6 and SEQ ID NO:8;

d) a monoclonal antibody comprising an amino acid sequence selected from the group consisting of the sequence shown in SEQ ID NO:2, the sequence shown in
10 SEQ ID NO:4, the sequence shown in SEQ ID NO:5, both the sequence shown in SEQ ID NO:2 and SEQ ID NO:4, and both the sequence shown in SEQ ID NO:2 and SEQ ID NO:5;

e) a monoclonal antibody having an amino acid sequence encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group
15 consisting of the sequence shown in SEQ ID NO:1, the sequence shown in SEQ ID NO:3, and both the sequence shown in SEQ ID NO:1 and SEQ ID NO:3;

f) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 5.9 or 12.12;

g) a monoclonal antibody that binds to an epitope comprising residues 82-87
20 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

h) a monoclonal antibody that binds to an epitope comprising residues 82-89 of the human CD40 sequence shown in SEQ ID NO:10 or SEQ ID NO:12;

i) a monoclonal antibody that competes with the monoclonal antibody CHIR-5.9 or CHIR-12.12 in a competitive binding assay;

25 j) the monoclonal antibody of preceding item a) or a monoclonal antibody of any one of preceding items c)-i), wherein said antibody is recombinantly produced; and

k) a monoclonal antibody that is an antigen-binding fragment of a monoclonal antibody of any one of preceding items a)-j), wherein said fragment retains the capability of specifically binding to said human CD40 antigen.

30

46. The pharmaceutical composition of claim 45, wherein said composition is a liquid pharmaceutical formulation comprising a buffer in an amount to maintain the pH of the formulation in a range of about pH 5.0 to about pH 7.0.

5 47. The pharmaceutical composition of claim 46, wherein said formulation further comprises an isotonizing agent in an amount to render same composition near isotonic.

10 48. The pharmaceutical composition of claim 47, wherein said isotonizing agent is sodium chloride, said sodium chloride being present in said formulation at a concentration of about 50 mM to about 300 mM.

49. The pharmaceutical composition of claim 48, wherein said sodium chloride is present in said formulation at a concentration of about 150 mM.

15 50. The pharmaceutical composition of any one of claims 46 to 49, wherein said buffer is selected from the group consisting of succinate, citrate, and phosphate buffers.



20 51. The pharmaceutical composition of claim 50, wherein said formulation comprises said buffer at a concentration of about 1 mM to about 50 mM.

52. The pharmaceutical composition of claim 51, wherein said buffer is sodium succinate or sodium citrate at a concentration of about 5 mM to about 15 mM.

25 53. The pharmaceutical composition of any one of claims 46 to 52, wherein said formulation further comprises a surfactant in an amount from about 0.001% to about 1.0%.

54. The pharmaceutical composition of claim 53, wherein said surfactant is polysorbate 80, which is present in said formulation in an amount from about 0.001% to about 0.5%.

FIGURE 1A



Fluorescence intensity

FIGURE 1B





FIGURE 2A

**FIGURE 2B**

4/16




FIGURE 3A

5/16




FIGURE 3B

6/16




FIGURE 4

7/16




FIGURE 5

8/16




FIGURE 6

[illegible]

	Daudi		Namalwa	
Exp.	CD40	CD20	CD40	CD20
E090403	14403.0	93676.5	3296.4	6200.1
E091003	13214.9	108438.5	3081.5	4788.2
E091103	13702.6	100509.1	3165.7	3988.3
E091203	13278.9	128158.3	3164.9	4618.0
Average	13,649.9	107,695.6	3,177.1	4,898.7
Stdev	546.7	14915.9	88.8	933.4

FIGURE 7

10/16




FIGURE 8

FIGURE 9ACHIR 12.12 light chain:

leader:

MALPAQLLGLLMLWVSGSSG

variable:

DIVMTQSPLSLTVTPGEPASISCRSSQSLLYSNGYNYLDWYLQKPGQSPQVLISLGSNRASG
VPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQARQTPFTFGPGTKVDIR

constant:

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK
DSTYLSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC**FIGURE 9B**CHIR-12.12 heavy chain:

leader:

MEFGLSWVFLVAILRGVQC

variable:

QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMHWVRQAPGKGLEWVAVISYEESNRYHAD
SVKGRFTISRDNISKITLYLQMNSLRTEDTAVYYCARDGGIAAPGPDYWGQGTLVTVSS

constant:

ASTKGPSVFPLAPASKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL
YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVV
SVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSV
MHEALHNHYTQKSLSLSPGK

alternative constant region:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL
YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVV
SVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSV
MHEALHNHYTQKSLSLSPGK

12/16

FIGURE 10A

DNA sequence of light chain of CHIR-12.12:

5'atggcgctccctgctcagctcctggggctgctaagtctctgggtctctggatccagtggggatattgtgatgactcagctccactctc
cctgaccgtcacccttgagagccggcctccatctcctgcagggtccagtcagagcctcctgtatagtaatggatacaactatttgattg
gtacctgcagaagccagggcagtcctccacaggtcctgatctcttgggttctaatacgggcctccgggtccctgacaggttcagtgga
gtggatcaggcacagattttactgaaaatcagcagagtgagggtgaggtgttgggttattactgcatgcaagctcgacaaact
ccattcactttcggccctgggaccaaagtggatatcagacgaactgtggctgcaccatctgtcttcattctcccgccatctgatgagcagt
tgaatctggaactgcctctgtgtgtgcctgtgaataacttctatcccagagaggccaaagtacagtggaaggtggataacgccctcc
aatcgggtaactcccaggagagtgacacagagcaggacgaaggacagcacctacagcctcagcagcacctgacgtgagcaa
agcagactacgagaaacacaaagtctacgcctgcgaagtcacccatcagggcctgagctcgccctgcacaaagagcttaacaggg
gagagtgttag3'

FIGURE 10B

DNA sequence of heavy chain of CHIR-12.12 (including introns):

5'atggagtttgggctgagctgggttttctgttctattttaagaggtgtccagtgacaggtgcagttggtggagcttgggggaggcgt
ggccagcctgggaggtccctgagactctcctgtgcagcctctggattcaccttcagtagctatggcatgcactgggtccgccaggctc
caggcaaggggctggagtgggtggcagttatatcatatgaggaaagtaatagatccatgcagactccgtgaaggggccgattcacca
tctccagagacaattccaagatcacgctgtatctgcaaatgaacagcctcagaactgaggacacggctgtgtattactgtgcgagagat
gggggtatagcagcacctgggcctgactactggggccagggaacctgtgtcaccgtctctcagcaagtaccaaggggccatccgt
cttccccctggcgcccgctagcaagagcacctctgggggcacagcgccctgggctgctgtcaaggactacttccccgaaccgg
tgacggtgtcgtggaactcaggcgccctgaccagcgcgctgcacaccttccggctgtctacagtcctcaggactctactccctcag
cagcgtggtgaccgtgccctccagcagcttgggacccagacctacatctgcaacgtgaatcacaaagcccagcaacaccaaggtgg
acaagagagttggtgagaggccagcacaggaggagggtgtctgctggaagccagggtcagcgctcctgacctggacgcatccc
gctatgcagtcctcagtcaggcagcaaggcaggccccgtctgcctcttccaccggaggcctctgcccggccactcatgctcagg
gagagggtcttctggcttttccccaggctctgggcaggcacaggctaggtgcccctaaccaggccctgcacacaaaggggcaggt
gctgggctcagacctgccaagagccatatccgggaggacctgcccctgacctaaagccaccccaaggccaaactctccactccc
tcagctcggacaccttctctctccagattccagtaactcccaattctctctgcagagcccaaatcttgtacaaaactcacacatgc
ccaccgtgcccaggtaagccagcccaggcctcgcctccagctcaaggcgggacaggtgccctagagtagctgcacccaggac
aggccccagccgggtgctgacacgtccacctccatctcttctcagcacctgaactcctggggggaccgtcagttctcttctcccc
aaaacccaaggacacctcatgatctccggacctgaggtcacatgcgtggtggtggacgtgagccacgaagacctgaggtca
agttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtaacagcacgtaccgtgt
ggtcagcgtctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaagggtctccaacaaagccctcccagccc
ccatcgagaaaaccatctccaaagccaaaggtgggacccgtgggtgagggccacatggacagaggccggctcggcccaccc
tctgccctgagagtgaccgtgtaccaacctgttcctacagggcagccccgagaaccacaggtgtacacctgcccccatccggg
gaggagatgaccaagaaccagggtcagcctgacctgctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaa
tgggcagccggagaacaactacaagaccacgcctcccgtgctggactccgacggctccttctctatagcaagctcaccgtggac
aagagcaggtggcagcagggaacgtctctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctcc
ctgtctccgggttaaatga3'

FIGURE 11ACHIR-5.9 light chain:

leader:

MALLAQLLGLLMLWVPGSSG

variable:

AIVMTQPPLSSPVTLGQPASISCRSSQSLVHSDGNTYLNWLQQRPGQPPRLLIYKFFRRLSG
VPDRFSGSGAGTDFTLKISRVEAEDVGVIYCMQVTQFPHTFGQGRLEIK

constant:

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNMFYPREAKVQWKVDNALQSGNSQESVTEQDSK
DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC**FIGURE 11B**CHIR-5.9 heavy chain:

leader:

MGSTAILALLLAVLQGVCA

variable:

EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMPGKGLEWMGIIYPGDS DTRYSP
SFQGQVTISADKSISTAYLQWSSLKASDTAMYICARGTAAGR DYYYYYGM D VWGQGT TTVTVS
S

constant:

ASTKGPSVFPLAPASKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL
YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVV
SVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSV
MHEALHNHYTQKSLSLSPGK

alternative constant region:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL
YSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVF
LFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVV
SVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSL
TCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSV
MHEALHNHYTQKSLSLSPGK

FIGURE 12A

Coding sequence for short isoform of human CD40:

```
1 atggttcgtc tgcctctgca gtgcgtctc tggggctgct tgctgaccgc tgtccatcca
61 gaaccaccca ctgcatgcag agaaaaacag tacctaataa acagtcagtg ctgttctttg
121 tgccagccag gacagaaact ggtgagtgcac tgcacagagt tcaactgaaac ggaatgcctt
181 cttgcggtg aaagcgaatt cctagacacc tggaacagag agacacactg ccaccagcac
241 aaatactgcg accccaacct agggcttcgg gtccagcaga agggcacctc agaaacagac
301 accatctgca cctgtgaaga aggtggcac tgtacgagtg aggcctgtga gagctgtgtc
361 ctgcaccgct catgctcgcc cggctttggg gtcaagcaga ttgctacagg ggtttctgat
421 accatctgcg agccctgccc agtcggcttc ttctccaatg tgatcatctgc ttctgaaaaa
481 tgtcaccctt ggacaaggct ccaggatcg gctgagagcc ctggtggtga tccccatcat
541 cttcgggatc ctgtttgcca tcctcttggt gctggtcttt atcaaaaagg tggccaagaa
601 gccaaccaat aa
```

FIGURE 12B

Encoded short isoform of human CD40:

```
1 mvrplqcvl wgclltavhp epptacrekq ylinsqccsl cqpqqlvsd cteftetecl
61 pcgesefldt wnrethchqh kyedpnlgrr vqqkgsetd tictceegwh ctseacescv
121 lhrscspgfg vkqiatgvsd ticepcpvgf fsnvssafek chpwtrspgs aespqgdphh
181 lrdpvchplg aglyqkqqe anq
```

FIGURE 12C


Coding sequence for long isoform of human CD40:

```
1 atggttcgtc tgcctctgca gtgcgtcctc tggggctgct tgctgaccgc tgtccatcca
61 gaaccaccca ctgcatgcag agaaaaacag tacctaataa acagtcagtg ctgttctttg
121 tgccagccag gacagaaact ggtgagtgac tgcacagagt tcaactgaaac ggaatgcctt
181 ccttgccgtg aaagcgaatt cctagacacc tggaacagag agacacactg ccaccagcac
241 aaatactgcg accccaacct agggcttcgg gtccagcaga agggcacctc agaaacagac
301 accatctgca cctgtgaaga aggctggcac tgtacgagtg aggcctgtga gagctgtgtc
361 ctgcaccgct catgctcgcc cggctttggg gtcaagcaga ttgctacagg ggtttctgat
421 accatctgcg agccctgccc agtcggcttc ttctccaatg tgtcatctgc ttctgaaaaa
481 tgcaccctt ggacaagctg tgagacaaa gacctggtg tgcaacaggc aggcacaaac
541 aagactgatg ttgtctgtgg tccccaggat cggtgagag ccctggtggt gatccccatc
601 atcttcggga tcctgtttgc catcctcttg gtgctggtct ttatcaaaaa ggtggccaag
661 aagccaacca ataaggcccc ccacccaag caggaacccc aggagatcaa ttttcccgac
721 gatcttctg gctccaacac tgctgtcca gtgcaggaga ctttacatgg atgccaaccg
781 gtcaccagg aggatggcaa agagagtgc atctcagtc aggagagaca gtga
```

FIGURE 12D

Encoded long isoform of human CD40:

```
1 mvrplqcvl wgclltavhp epptacrekq ylinsqccsl cpggqlvds cteftetcl
61 pcgesefldt wnrethchqh kyedpnlglr vqkgtsetd tictceegwh ctseacescv
121 lhrscspgfg vkqiatgvsd ticepcpvgf fsnvssafek chpwtscetk dlvvqqagtn
181 ktdvvcgpqd rlralvvipi ifgilfaill vlvfikkvak kptnkaphpk qepqeinfpd
241 dlpgsntaap vqetlhgcqp vtqedgkesr isvqerq
```

FIGURE 13

SEQUENCE LISTING

<110> Chen, Bao-Lu
 Hurst, Deborah
 Lee, Sang Hoon
 Long, Li
 Lu, Xiaofeng
 Luqman, Mohammad
 Yabannavar, Asha
 Zaror, Isabel

<120> Antagonist Anti-CD40 Monoclonal
 Antibodies and Methods for Their Use

<130> PP20107.004 (282916)

<150> 60/565,710

<151> 2004-04-27

<150> 60/525,579

<151> 2003-11-26

<150> 60/517,337

<151> 2003-11-04

<160> 12

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 720

<212> DNA

<213> Artificial Sequence

<220>

<223> Coding sequence for light chain of 12.12 human
 anti-CD40 antibody

<221> CDS

<222> (1)...(720)

<400> 1

atg gcg ctc cct gct cag ctc ctg ggg ctg cta atg ctc tgg gtc tct	48
Met Ala Leu Pro Gln Leu Leu Gly Leu Met Leu Trp Val Ser	
1 5 10 15	
gga tcc agt ggg gat att gtg atg act cag tct cca ctc tcc ctg acc	96
Gly Ser Ser Gly Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Thr	
20 25 30	
gtc acc cct gga gag ccg gcc tcc atc tcc tgc agg tcc agt cag agc	144
Val Thr Pro Gly Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser	
35 40 45	
ctc ctg tat agt aat gga tac aac tat ttg gat tgg tac ctg cag aag	192
Leu Leu Tyr Ser Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys	
50 55 60	
cca ggg cag tct cca cag gtc ctg atc tct ttg ggt tct aat cgg gcc	240
Pro Gly Gln Ser Pro Gln Val Leu Ile Ser Leu Gly Ser Asn Arg Ala	
65 70 75 80	
tcc ggg gtc cct gac agg ttc agt ggc agt gga tca ggc aca gat ttt	288
Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe	

85										90					95					
aca	ctg	aaa	atc	agc	aga	gtg	gag	gct	gag	gat	gtt	ggg	gtt	tat	tac	336				
Thr	Leu	Lys	Ile	Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr					
			100					105					110							
tgc	atg	caa	gct	cga	caa	act	cca	ttc	act	ttc	ggc	cct	ggg	acc	aaa	384				
Cys	Met	Gln	Ala	Arg	Gln	Thr	Pro	Phe	Thr	Phe	Gly	Pro	Gly	Thr	Lys					
		115					120					125								
gtg	gat	atc	aga	cga	act	gtg	gct	gca	cca	tct	gtc	ttc	atc	ttc	ccg	432				
Val	Asp	Ile	Arg	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro					
	130					135					140									
cca	tct	gat	gag	cag	ttg	aaa	tct	gga	act	gcc	tct	gtt	gtg	tgc	ctg	480				
Pro	Ser	Asp	Glu	Gln	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Val	Cys	Leu					
145					150				155						160					
ctg	aat	aac	ttc	tat	ccc	aga	gag	gcc	aaa	gta	cag	tgg	aag	gtg	gat	528				
Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp					
			165					170					175							
aac	gcc	ctc	caa	tcg	ggt	aac	tcc	cag	gag	agt	gtc	aca	gag	cag	gac	576				
Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp					
		180					185					190								
agc	aag	gac	agc	acc	tac	agc	ctc	agc	agc	acc	ctg	acg	ctg	agc	aaa	624				
Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys					
		195				200					205									
gca	gac	tac	gag	aaa	cac	aaa	gtc	tac	gcc	tgc	gaa	gtc	acc	cat	cag	672				
Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln					
	210					215					220									
ggc	ctg	agc	tcg	ccc	gtc	aca	aag	agc	ttc	aac	agg	gga	gag	tgt	tag	720				
Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys	*					
225					230				235											

<210> 2

<211> 239

<212> PRT

<213> Artificial Sequence

<220>

<223> Light chain of 12.12 human anti-CD40 antibody

<400> 2

Met	Ala	Leu	Pro	Ala	Gln	Leu	Leu	Gly	Leu	Leu	Met	Leu	Trp	Val	Ser
1				5					10					15	
Gly	Ser	Ser	Gly	Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Leu	Ser	Leu	Thr
			20					25					30		
Val	Thr	Pro	Gly	Glu	Pro	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser
		35				40					45				
Leu	Leu	Tyr	Ser	Asn	Gly	Tyr	Asn	Tyr	Leu	Asp	Trp	Tyr	Leu	Gln	Lys
	50				55				60						
Pro	Gly	Gln	Ser	Pro	Gln	Val	Leu	Ile	Ser	Leu	Gly	Ser	Asn	Arg	Ala
65					70				75					80	
Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe
			85					90					95		
Thr	Leu	Lys	Ile	Ser	Arg	Val	Glu	Ala	Glu	Asp	Val	Gly	Val	Tyr	Tyr
			100				105					110			
Cys	Met	Gln	Ala	Arg	Gln	Thr	Pro	Phe	Thr	Phe	Gly	Pro	Gly	Thr	Lys
		115				120					125				
Val	Asp	Ile	Arg	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro

130	135	140
Pro Ser Asp Glu Gln Leu	Lys Ser Gly Thr Ala	Ser Val Val Cys Leu
145	150	155
Leu Asn Asn Phe Tyr Pro	Arg Glu Ala Lys Val	Gln Trp Lys Val Asp
165	170	175
Asn Ala Leu Gln Ser Gly	Asn Ser Gln Glu Ser	Val Thr Glu Gln Asp
180	185	190
Ser Lys Asp Ser Thr Tyr	Ser Leu Ser Ser Thr	Leu Thr Leu Ser Lys
195	200	205
Ala Asp Tyr Glu Lys His	Lys Val Tyr Ala Cys	Glu Val Thr His Gln
210	215	220
Gly Leu Ser Ser Pro Val	Thr Lys Ser Phe Asn	Arg Gly Glu Cys
225	230	235

<210> 3
 <211> 2016
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Coding sequence for heavy chain of 12.12 human
 anti-CD40 antibody (with introns)

<400> 3

atg	gag	ttt	ggg	ctg	agc	tgg	gtt	ttc	ctt	gtt	gct	att	tta	aga	ggt	48
gtc	cag	tgt	cag	gtg	cag	ttg	gtg	gag	tct	ggg	gga	ggc	gtg	gtc	cag	96
cct	ggg	agg	tcc	ctg	aga	ctc	tcc	tgt	gca	gcc	tct	gga	ttc	acc	ttc	144
agt	agc	tat	ggc	atg	cac	tgg	gtc	cgc	cag	gct	cca	ggc	aag	ggg	ctg	192
gag	tgg	gtg	gca	gtt	ata	tca	tat	gag	gaa	agt	aat	aga	tac	cat	gca	240
gac	tcc	gtg	aag	ggc	cga	ttc	acc	atc	tcc	aga	gac	aat	tcc	aag	atc	288
acg	ctg	tat	ctg	caa	atg	aac	agc	ctc	aga	act	gag	gac	acg	gct	gtg	336
tat	tac	tgt	gcg	aga	gat	ggg	ggt	ata	gca	gca	cct	ggg	cct	gac	tac	384
tgg	ggc	cag	gga	acc	ctg	gtc	acc	gtc	tcc	tca	gca	agt	acc	aag	ggc	432
cca	tcc	gtc	ttc	ccc	ctg	gcg	ccc	gct	agc	aag	agc	acc	tct	ggg	ggc	480
aca	gcg	gcc	ctg	ggc	tgc	ctg	gtc	aag	gac	tac	ttc	ccc	gaa	ccg	gtg	528
acg	gtg	tcg	tgg	aac	tca	ggc	gcc	ctg	acc	agc	ggc	gtg	cac	acc	ttc	576
ccg	gct	gtc	cta	cag	tcc	tca	gga	ctc	tac	tcc	ctc	agc	agc	gtg	gtg	624
acc	gtg	ccc	tcc	agc	agc	ttg	ggc	acc	cag	acc	tac	atc	tgc	aac	gtg	672
aat	cac	aag	ccc	agc	aac	acc	aag	gtg	gac	aag	aga	gtt	ggt	gag	agg	720
cca	gca	cag	gga	ggg	agg	gtg	tct	gct	gga	agc	cag	gct	cag	cgc	tcc	768
tgc	ctg	gac	gca	tcc	cgg	cta	tgc	agt	ccc	agt	cca	ggg	cag	caa	ggc	816
agg	ccc	cgt	ctg	cct	ctt	cac	ccg	gag	gcc	tct	gcc	cgc	ccc	act	cat	864
gct	cag	gga	gag	ggt	ctt	ctg	gct	ttt	tcc	cca	ggc	tct	ggg	cag	gca	912
cag	gct	agg	tgc	ccc	taa	ccc	agg	ccc	tgc	aca	caa	agg	ggc	agg	tgc	960
tgg	gct	cag	acc	tgc	caa	gag	cca	tat	ccg	gga	gga	ccc	tgc	ccc	tga	1008
cct	aag	ccc	acc	cca	aag	gcc	aaa	ctc	tcc	act	ccc	tca	gct	cgg	aca	1056
cct	tct	ctc	ctc	cca	gat	tcc	agt	aac	tcc	caa	tct	tct	ctc	tgc	aga	1104
gcc	caa	atc	ttg	tga	caa	aac	tca	cac	atg	ccc	acc	gtg	ccc	agg	taa	1152
gcc	agc	cca	ggc	ctc	gcc	ctc	cag	ctc	aag	gcg	gga	cag	gtg	ccc	tag	1200
agt	agc	ctg	cat	cca	ggg	aca	ggc	ccc	agc	cgg	gtg	ctg	aca	cgt	cca	1248
cct	cca	tct	ctt	cct	cag	cac	ctg	aac	tcc	tgg	ggg	gac	cgt	cag	tct	1296
tcc	tct	tcc	ccc	caa	aac	cca	agg	aca	ccc	tca	tga	tct	ccc	gga	ccc	1344
ctg	agg	tca	cat	gcg	tgg	tgg	tgg	acg	tga	gcc	acg	aag	acc	ctg	agg	1392
tca	agt	tca	act	ggt	acg	tgg	acg	gcg	tgg	agg	tgc	ata	atg	cca	aga	1440
caa	agc	cgc	ggg	agg	agc	agt	aca	aca	gca	cgt	acc	gtg	tgg	tca	gcg	1488
tcc	tca	ccg	tcc	tgc	acc	agg	act	ggc	tga	atg	gca	agg	agt	aca	agt	1536
gca	agg	tct	cca	aca	aag	ccc	tcc	cag	ccc	cca	tcg	aga	aaa	cca	tct	1584
cca	aag	cca	aag	gtg	gga	ccc	gtg	ggg	tgc	gag	ggc	cac	atg	gac	aga	1632
ggc	cgg	ctc	ggc	cca	ccc	tct	gcc	ctg	aga	gtg	acc	gct	gta	cca	acc	1680
tct	gtc	cct	aca	ggg	cag	ccc	cga	gaa	cca	cag	gtg	tac	acc	ctg	ccc	1728
cca	tcc	cgg	gag	gag	atg	acc	aag	aac	cag	gtc	agc	ctg	acc	tgc	ctg	1776
gtc	aaa	ggc	ttc	tat	ccc	agc	gac	atc	gcc	gtg	gag	tgg	gag	agc	aat	1824
ggg	cag	ccg	gag	aac	aac	tac	aag	acc	acg	cct	ccc	gtg	ctg	gac	tcc	1872
gac	ggc	tcc	ttc	ttc	ctc	tat	agc	aag	ctc	acc	gtg	gac	aag	agc	agg	1920
tgg	cag	cag	ggg	aac	gtc	ttc	tca	tgc	gtg	atg	cat	gag	gct	ctg	1968	

cac aac cac tac acg cag aag agc ctc tcc ctg tct ccg ggt aaa tga 2016

<210> 4

<211> 469

<212> PRT

<213> Artificial Sequence

$\langle 220 \rangle$

<223> Heavy chain of 12.12 human anti-CD40 antibody

<400> 4

Met 1	Glu	Phe	Gly	Leu 5	Ser	Trp	Val	Phe	Leu 10	Val	Ala	Ile	Leu	Arg	Gly
Val	Gln	Cys	Gln 20	Val	Gln	Leu	Val	Glu 25	Ser	Gly	Gly	Gly	Val 30	Val	Gln
Pro	Gly	Arg	Ser 35	Leu	Arg	Leu	Ser 40	Cys	Ala	Ala	Ser	Gly 45	Phe	Thr	Phe
Ser	Ser 50	Tyr	Gly	Met	His	Trp 55	Val	Arg	Gln	Ala 60	Pro	Gly	Lys	Gly	Leu
Glu 65	Trp	Val	Ala	Val 70	Ile	Ser	Tyr	Glu	Glu	Ser 75	Asn	Arg	Tyr	His	Ala 80
Asp	Ser	Val	Lys 85	Gly	Arg	Phe	Thr	Ile	Ser 90	Arg	Asp	Asn	Ser 95	Lys	Ile
Thr	Leu	Tyr	Leu 100	Gln	Met	Asn	Ser 105	Leu	Arg	Thr	Glu	Asp 110	Thr	Ala	Val
Tyr	Tyr	Cys 115	Ala	Arg	Asp	Gly 120	Gly	Ile	Ala	Ala	Pro	Gly 125	Pro	Asp	Tyr
Trp	Gly 130	Gln	Gly	Thr	Leu	Val 135	Thr	Val	Ser	Ser	Ala 140	Ser	Thr	Lys	Gly
Pro 145	Ser	Val	Phe	Pro 150	Leu	Ala	Pro	Ala	Ser	Lys 155	Ser	Thr	Ser	Gly	Gly 160
Thr	Ala	Ala	Leu 165	Gly	Cys	Leu	Val	Lys	Asp 170	Tyr	Phe	Pro	Glu 175	Pro	Val
Thr	Val	Ser	Trp 180	Asn	Ser	Gly	Ala 185	Leu	Thr	Ser	Gly	Val 190	His	Thr	Phe
Pro	Ala 195	Val	Leu	Gln	Ser	Ser 200	Gly	Leu	Tyr	Ser	Leu 205	Ser	Ser	Val	Val
Thr	Val 210	Pro	Ser	Ser	Ser 215	Leu	Gly	Thr	Gln	Thr 220	Tyr	Ile	Cys	Asn	Val
Asn 225	His	Lys	Pro	Ser 230	Asn	Thr	Lys	Val	Asp	Lys 235	Arg	Val	Glu	Pro	Lys 240
Ser	Cys	Asp	Lys 245	Thr	His	Thr	Cys	Pro	Pro 250	Cys	Pro	Ala	Pro	Glu	Leu 255
Leu	Gly	Gly	Pro 260	Ser	Val	Phe	Leu	Phe 265	Pro	Pro	Lys	Pro	Lys 270	Asp	Thr
Leu	Met 275	Ile	Ser	Arg	Thr	Pro	Glu 280	Val	Thr	Cys	Val 285	Val	Val	Asp	Val
Ser	His 290	Glu	Asp	Pro	Glu	Val 295	Lys	Phe	Asn	Trp	Tyr 300	Val	Asp	Gly	Val
Glu 305	Val	His	Asn	Ala 310	Lys	Thr	Lys	Pro	Arg	Glu 315	Gln	Tyr	Asn	Ser	
Thr	Tyr	Arg	Val 325	Val	Ser	Val	Leu	Thr	Val 330	Leu	His	Gln	Asp	Trp	Leu
Asn	Gly	Lys	Glu 340	Tyr	Lys	Cys	Lys	Val 345	Ser	Asn	Lys	Ala	Leu 350	Pro	Ala
Pro	Ile 355	Glu	Lys	Thr	Ile	Ser 360	Lys	Ala	Lys	Gly	Gln 365	Pro	Arg	Glu	Pro
Gln	Val 370	Tyr	Thr	Leu	Pro	Pro 375	Ser	Arg	Glu	Glu 380	Met	Thr	Lys	Asn	Gln
Val 385	Ser	Leu	Thr	Cys 390	Leu	Val	Lys	Gly	Phe	Tyr 395	Pro	Ser	Asp	Ile	Ala
Val	Glu	Trp	Glu 405	Ser	Asn	Gly	Gln	Pro	Glu 410	Asn	Asn	Tyr	Lys	Thr	Thr
Pro	Pro	Val	Leu 420	Asp	Ser	Asp	Gly	Ser 425	Phe	Phe	Leu	Tyr	Ser 430	Lys	Leu

Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 435 440 445
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 450 455 460
 Leu Ser Pro Gly Lys
 465

<210> 5

<211> 469

<212> PRT

<213> Artificial Sequence

<220>

<223> Heavy chain of variant of 12.12 human anti-CD40
 antibody

<400> 5

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Arg Gly
 1 5 10 15
 Val Gln Cys Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln
 20 25 30
 Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Ser Tyr Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Val Ala Val Ile Ser Tyr Glu Glu Ser Asn Arg Tyr His Ala
 65 70 75 80
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Ile
 85 90 95
 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Thr Glu Asp Thr Ala Val
 100 105 110
 Tyr Tyr Cys Ala Arg Asp Gly Gly Ile Ala Ala Pro Gly Pro Asp Tyr
 115 120 125
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly
 130 135 140
 Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly
 145 150 155 160
 Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val
 165 170 175
 Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe
 180 185 190
 Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val
 195 200 205
 Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val
 210 215 220
 Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys
 225 230 235 240
 Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu
 245 250 255
 Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
 260 265 270
 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
 275 280 285
 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val
 290 295 300
 Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser
 305 310 315 320
 Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
 325 330 335
 Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala
 340 345 350
 Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro
 355 360 365
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln
 370 375 380

```

Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
385                      390                      395                      400
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
                      405                      410                      415
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
                      420                      425                      430
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
435                      440                      445
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
450                      455                      460
Leu Ser Pro Gly Lys
465

```

<210> 6
 <211> 239
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Light chain of 5.9 human anti-CD40 antibody

```

<400> 6
Met Ala Leu Leu Ala Gln Leu Leu Gly Leu Leu Met Leu Trp Val Pro
1                      5                      10                      15
Gly Ser Ser Gly Ala Ile Val Met Thr Gln Pro Pro Leu Ser Ser Pro
20                      25                      30
Val Thr Leu Gly Gln Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser
35                      40                      45
Leu Val His Ser Asp Gly Asn Thr Tyr Leu Asn Trp Leu Gln Gln Arg
50                      55                      60
Pro Gly Gln Pro Pro Arg Leu Leu Ile Tyr Lys Phe Phe Arg Arg Leu
65                      70                      75                      80
Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ala Gly Thr Asp Phe
85                      90                      95
Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr
100                      105                      110
Cys Met Gln Val Thr Gln Phe Pro His Thr Phe Gly Gln Gly Thr Arg
115                      120                      125
Leu Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro
130                      135                      140
Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu
145                      150                      155                      160
Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp
165                      170                      175
Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp
180                      185                      190
Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys
195                      200                      205
Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln
210                      215                      220
Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
225                      230                      235

```

<210> 7
 <211> 474
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Heavy chain of 5.9 human anti-CD40 antibody

```

<400> 7
Met Gly Ser Thr Ala Ile Leu Ala Leu Leu Leu Ala Val Leu Gln Gly
1                      5                      10                      15

```

```

Val Cys Ala Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
      20      25      30
Pro Gly Glu Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe
      35      40      45
Thr Ser Tyr Trp Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu
      50      55      60
Glu Trp Met Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser
      65      70      75      80
Pro Ser Phe Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser
      85      90      95
Thr Ala Tyr Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met
      100      105      110
Tyr Tyr Cys Ala Arg Gly Thr Ala Ala Gly Arg Asp Tyr Tyr Tyr Tyr
      115      120      125
Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
      130      135      140
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ala Ser Lys
      145      150      155      160
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
      165      170      175
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
      180      185      190
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
      195      200      205
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
      210      215      220
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
      225      230      235      240
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
      245      250      255
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
      260      265      270
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
      275      280      285
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
      290      295      300
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
      305      310      315      320
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
      325      330      335
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
      340      345      350
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
      355      360      365
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
      370      375      380
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      385      390      395      400
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      405      410      415
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
      420      425      430
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      435      440      445
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      450      455      460
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
      465      470

```

<210> 8

<211> 474

<212> PRT

<213> Artificial Sequence

<220>

<223> Heavy chain of variant of 5.9 human anti-CD40
antibody

<400> 8

```

Met Gly Ser Thr Ala Ile Leu Ala Leu Leu Leu Ala Val Leu Gln Gly
 1          5          10          15
Val Cys Ala Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
          20          25          30
Pro Gly Glu Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe
          35          40          45
Thr Ser Tyr Trp Ile Gly Trp Val Arg Gln Met Pro Gly Lys Gly Leu
          50          55          60
Glu Trp Met Gly Ile Ile Tyr Pro Gly Asp Ser Asp Thr Arg Tyr Ser
          65          70          75          80
Pro Ser Phe Gln Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser
          85          90          95
Thr Ala Tyr Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met
          100          105          110
Tyr Tyr Cys Ala Arg Gly Thr Ala Ala Gly Arg Asp Tyr Tyr Tyr Tyr
          115          120          125
Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
          130          135          140
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
          145          150          155          160
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
          165          170          175
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
          180          185          190
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
          195          200          205
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
          210          215          220
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
          225          230          235          240
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
          245          250          255
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
          260          265          270
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
          275          280          285
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
          290          295          300
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
          305          310          315          320
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
          325          330          335
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
          340          345          350
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
          355          360          365
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
          370          375          380
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
          385          390          395          400
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
          405          410          415
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
          420          425          430
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
          435          440          445
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
          450          455          460
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
          465          470

```


<210> 9
 <211> 612
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)...(612)

<221> misc_feature
 <222> (0)...(0)
 <223> Coding sequence for short isoform of human CD40

```

<400> 9
atg gtt cgt ctg cct ctg cag tgc gtc ctc tgg ggc tgc ttg ctg acc 48
Met Val Arg Leu Pro Leu Gln Cys Val Leu Trp Gly Cys Leu Leu Thr
1 5 10 15

gct gtc cat cca gaa cca ccc act gca tgc aga gaa aaa cag tac cta 96
Ala Val His Pro Glu Pro Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu
20 25 30

ata aac agt cag tgc tgt tct ttg tgc cag cca gga cag aaa ctg gtg 144
Ile Asn Ser Gln Cys Cys Ser Leu Cys Gln Pro Gly Gln Lys Leu Val
35 40 45

agt gac tgc aca gag ttc act gaa acg gaa tgc ctt cct tgc ggt gaa 192
Ser Asp Cys Thr Glu Phe Thr Glu Thr Glu Cys Leu Pro Cys Gly Glu
50 55 60

agc gaa ttc cta gac acc tgg aac aga gag aca cac tgc cac cag cac 240
Ser Glu Phe Leu Asp Thr Trp Asn Arg Glu Thr His Cys His Gln His
65 70 75 80

aaa tac tgc gac ccc aac cta ggg ctt cgg gtc cag cag aag ggc acc 288
Lys Tyr Cys Asp Pro Asn Leu Gly Leu Arg Val Gln Gln Lys Gly Thr
85 90 95

tca gaa aca gac acc atc tgc acc tgt gaa gaa ggc tgg cac tgt acg 336
Ser Glu Thr Asp Thr Ile Cys Thr Cys Glu Glu Gly Trp His Cys Thr
100 105 110

agt gag gcc tgt gag agc tgt gtc ctg cac cgc tca tgc tcg ccc ggc 384
Ser Glu Ala Cys Glu Ser Cys Val Leu His Arg Ser Cys Ser Pro Gly
115 120 125

ttt ggg gtc aag cag att gct aca ggg gtt tct gat acc atc tgc gag 432
Phe Gly Val Lys Gln Ile Ala Thr Gly Val Ser Asp Thr Ile Cys Glu
130 135 140

ccc tgc cca gtc ggc ttc ttc tcc aat gtg tca tct gct ttc gaa aaa 480
Pro Cys Pro Val Gly Phe Phe Ser Asn Val Ser Ser Ala Phe Glu Lys
145 150 155 160

tgt cac cct tgg aca agg tcc cca gga tcg gct gag agc cct ggt ggt 528
Cys His Pro Trp Thr Arg Ser Pro Gly Ser Ala Glu Ser Pro Gly Gly
165 170 175

gat ccc cat cat ctt cgg gat cct gtt tgc cat cct ctt ggt gct ggt 576
Asp Pro His His Leu Arg Asp Pro Val Cys His Pro Leu Gly Ala Gly
180 185 190

ctt tat caa aaa ggt ggc caa gaa gcc aac caa taa 612
Leu Tyr Gln Lys Gly Gly Gln Glu Ala Asn Gln *
195 200

```

<210> 10
 <211> 203
 <212> PRT
 <213> Homo sapiens

<400> 10
 Met Val Arg Leu Pro Leu Gln Cys Val Leu Trp Gly Cys Leu Leu Thr
 1 5 10 15
 Ala Val His Pro Glu Pro Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu
 20 25 30
 Ile Asn Ser Gln Cys Cys Ser Leu Cys Gln Pro Gly Gln Lys Leu Val
 35 40 45
 Ser Asp Cys Thr Glu Phe Thr Glu Thr Glu Cys Leu Pro Cys Gly Glu
 50 55 60
 Ser Glu Phe Leu Asp Thr Trp Asn Arg Glu Thr His Cys His Gln His
 65 70 75 80
 Lys Tyr Cys Asp Pro Asn Leu Gly Leu Arg Val Gln Gln Lys Gly Thr
 85 90 95
 Ser Glu Thr Asp Thr Ile Cys Thr Cys Glu Glu Gly Trp His Cys Thr
 100 105 110
 Ser Glu Ala Cys Glu Ser Cys Val Leu His Arg Ser Cys Ser Pro Gly
 115 120 125
 Phe Gly Val Lys Gln Ile Ala Thr Gly Val Ser Asp Thr Ile Cys Glu
 130 135 140
 Pro Cys Pro Val Gly Phe Phe Ser Asn Val Ser Ser Ala Phe Glu Lys
 145 150 155 160
 Cys His Pro Trp Thr Arg Ser Pro Gly Ser Ala Glu Ser Pro Gly Gly
 165 170 175
 Asp Pro His His Leu Arg Asp Pro Val Cys His Pro Leu Gly Ala Gly
 180 185 190
 Leu Tyr Gln Lys Gly Gly Gln Glu Ala Asn Gln
 195 200

<210> 11
 <211> 834
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)...(834)
 <221> misc_feature
 <222> (0)...(0)
 <223> Coding sequence for long isoform of human CD40

<400> 11
 atg gtt cgt ctg cct ctg cag tgc gtc ctc tgg ggc tgc ttg ctg acc 48
 Met Val Arg Leu Pro Leu Gln Cys Val Leu Trp Gly Cys Leu Leu Thr
 1 5 10 15
 gct gtc cat cca gaa cca ccc act gca tgc aga gaa aaa cag tac cta 96
 Ala Val His Pro Glu Pro Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu
 20 25 30
 ata aac agt cag tgc tgt tct ttg tgc cag cca gga cag aaa ctg gtg 144
 Ile Asn Ser Gln Cys Cys Ser Leu Cys Gln Pro Gly Gln Lys Leu Val
 35 40 45
 agt gac tgc aca gag ttc act gaa acg gaa tgc ctt cct tgc ggt gaa 192
 Ser Asp Cys Thr Glu Phe Thr Glu Thr Glu Cys Leu Pro Cys Gly Glu
 50 55 60
 agc gaa ttc cta gac acc tgg aac aga gag aca cac tgc cac cag cac 240

```

Ser Glu Phe Leu Asp Thr Trp Asn Arg Glu Thr His Cys His Gln His
65              70              75              80

aaa tac tgc gac ccc aac cta ggg ctt cgg gtc cag cag aag ggc acc 288
Lys Tyr Cys Asp Pro Asn Leu Gly Leu Arg Val Gln Gln Lys Gly Thr
              85              90              95

tca gaa aca gac acc atc tgc acc tgt gaa gaa ggc tgg cac tgt acg 336
Ser Glu Thr Asp Thr Ile Cys Thr Cys Glu Glu Gly Trp His Cys Thr
              100              105              110

agt gag gcc tgt gag agc tgt gtc ctg cac cgc tca tgc tcg ccc ggc 384
Ser Glu Ala Cys Glu Ser Cys Val Leu His Arg Ser Cys Ser Pro Gly
              115              120              125

ttt ggg gtc aag cag att gct aca ggg gtt tct gat acc atc tgc gag 432
Phe Gly Val Lys Gln Ile Ala Thr Gly Val Ser Asp Thr Ile Cys Glu
              130              135              140

ccc tgc cca gtc ggc ttc ttc tcc aat gtg tca tct gct ttc gaa aaa 480
Pro Cys Pro Val Gly Phe Phe Ser Asn Val Ser Ser Ala Phe Glu Lys
145              150              155              160

tgt cac cct tgg aca agc tgt gag acc aaa gac ctg gtt gtg caa cag 528
Cys His Pro Trp Thr Ser Cys Glu Thr Lys Asp Leu Val Val Gln Gln
              165              170              175

gca ggc aca aac aag act gat gtt gtc tgt ggt ccc cag gat cgg ctg 576
Ala Gly Thr Asn Lys Thr Asp Val Val Cys Gly Pro Gln Asp Arg Leu
              180              185              190

aga gcc ctg gtg gtg atc ccc atc atc ttc ggg atc ctg ttt gcc atc 624
Arg Ala Leu Val Val Ile Pro Ile Ile Phe Gly Ile Leu Phe Ala Ile
              195              200              205

ctc ttg gtg ctg gtc ttt atc aaa aag gtg gcc aag aag cca acc aat 672
Leu Leu Val Leu Val Phe Ile Lys Lys Val Ala Lys Lys Pro Thr Asn
              210              215              220

aag gcc ccc cac ccc aag cag gaa ccc cag gag atc aat ttt ccc gac 720
Lys Ala Pro His Pro Lys Gln Glu Pro Gln Glu Ile Asn Phe Pro Asp
225              230              235              240

gat ctt cct ggc tcc aac act gct gct cca gtg cag gag act tta cat 768
Asp Leu Pro Gly Ser Asn Thr Ala Ala Pro Val Gln Glu Thr Leu His
              245              250              255

gga tgc caa ccg gtc acc cag gag gat ggc aaa gag agt cgc atc tca 816
Gly Cys Gln Pro Val Thr Gln Glu Asp Gly Lys Glu Ser Arg Ile Ser
              260              265              270

gtg cag gag aga cag tga
Val Gln Glu Arg Gln *
              275

```

```

<210> 12
<211> 277
<212> PRT
<213> Homo sapiens

```

```

<400> 12
Met Val Arg Leu Pro Leu Gln Cys Val Leu Trp Gly Cys Leu Leu Thr
1          5          10          15
Ala Val His Pro Glu Pro Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu
20          25          30

```

Ile	Asn	Ser	Gln	Cys	Cys	Ser	Leu	Cys	Gln	Pro	Gly	Gln	Lys	Leu	Val
	35						40					45			
Ser	Asp	Cys	Thr	Glu	Phe	Thr	Glu	Thr	Glu	Cys	Leu	Pro	Cys	Gly	Glu
	50					55					60				
Ser	Glu	Phe	Leu	Asp	Thr	Trp	Asn	Arg	Glu	Thr	His	Cys	His	Gln	His
65					70				75					80	
Lys	Tyr	Cys	Asp	Pro	Asn	Leu	Gly	Leu	Arg	Val	Gln	Gln	Lys	Gly	Thr
			85					90						95	
Ser	Glu	Thr	Asp	Thr	Ile	Cys	Thr	Cys	Glu	Glu	Gly	Trp	His	Cys	Thr
		100					105						110		
Ser	Glu	Ala	Cys	Glu	Ser	Cys	Val	Leu	His	Arg	Ser	Cys	Ser	Pro	Gly
	115						120					125			
Phe	Gly	Val	Lys	Gln	Ile	Ala	Thr	Gly	Val	Ser	Asp	Thr	Ile	Cys	Glu
	130					135					140				
Pro	Cys	Pro	Val	Gly	Phe	Phe	Ser	Asn	Val	Ser	Ser	Ala	Phe	Glu	Lys
145					150				155					160	
Cys	His	Pro	Trp	Thr	Ser	Cys	Glu	Thr	Lys	Asp	Leu	Val	Val	Gln	Gln
			165						170					175	
Ala	Gly	Thr	Asn	Lys	Thr	Asp	Val	Val	Cys	Gly	Pro	Gln	Asp	Arg	Leu
		180					185						190		
Arg	Ala	Leu	Val	Val	Ile	Pro	Ile	Ile	Phe	Gly	Ile	Leu	Phe	Ala	Ile
	195					200						205			
Leu	Leu	Val	Leu	Val	Phe	Ile	Lys	Lys	Val	Ala	Lys	Lys	Pro	Thr	Asn
	210					215					220				
Lys	Ala	Pro	His	Pro	Lys	Gln	Glu	Pro	Gln	Glu	Ile	Asn	Phe	Pro	Asp
225					230				235					240	
Asp	Leu	Pro	Gly	Ser	Asn	Thr	Ala	Ala	Pro	Val	Gln	Glu	Thr	Leu	His
			245						250					255	
Gly	Cys	Gln	Pro	Val	Thr	Gln	Glu	Asp	Gly	Lys	Glu	Ser	Arg	Ile	Ser
			260					265					270		
Val	Gln	Glu	Arg	Gln											
	275														